

ASSESSMENT OF NUTRIENTS LEVELS IN GROUNDWATER WITHIN THE LOWER PRA BASIN OF GHANA

M. K. DORLEKU*, A. O. AFFUM, C. K. TAY AND D. NUKPEZAH

(M. K. D. & C. K. T.: CSIR- Water Research Institute, P.O Box AH 38 Achimota; A. O. A.: National Nuclear Research Institute, Ghana Atomic Energy Commission P.O Box LG 80, Legon; D. N.: Institute of Environmental Studies and Sanitation, University of Ghana, Legon, Accra, Ghana).

Corresponding author's email: mdorleku@hotmail.com

Abstract

The level of concentration of nitrates, phosphates, and sulphates was determined in groundwater within the Pra Basin to ascertain the degree of anthropogenic influence via the application of fertilizer and other agro-chemical to farmlands by farmers. Sixty-five (65) boreholes in forty-five (45) communities were analysed between January and April 2012 to cover the dry season and between June and October 2012 to cover the wet season. The results show that nitrate for both seasons range between 0.011 mg/l and 5.01 mg/l and therefore within the WHO (2006) guideline value of 10 mg/l, whilst that of phosphate range between 0.01 mg/l and 3.08 mg/l and therefore outside the USEPA guideline value of 2.5 mg/l. Sulphate recorded a range between 1.60 and 96.2 mg/l and therefore within the WHO (2006) guideline value of 250 mg/l. The concentration levels of the two nutrients of the parameters, nitrates and phosphate were correlated against borehole depth and the results show a weak positive and negative correlation for nitrate and phosphate respectively. There was thus, an indication that the levels of all parameters, nitrate, phosphate and sulphate are not yet a threat to groundwater in the basin.

Keywords: Groundwater, Nutrients, Fertilizer application, Agrochemicals, Biogeological processes

Introduction

Globally, the safety and quality of water are essential, especially to human health, agriculture, aquaculture, industry and all forms of life. Over the last few years, groundwater resource has become the principal source of domestic water supply in many countries of the world including Ghana (Hynds *et al.* 2014). The preference of groundwater over surface water as a source of drinking water is based on the fact that groundwater excluded from the atmosphere would be less susceptible to pollution (Kortatsi, 2006; Smedley *et al.*, 1995). This is true to an extent. However,

groundwater in hard-rock aquifers, particularly in mining areas, is known to be vulnerable to quality problems that may have serious impact on human health as a result of carbonate deficient rocks that give rise to poorly buffered waters (Smedley *et al.*, 1995). It has been reported that many public health surveys and water quality analyses have shown that groundwater is not immune to a wide range of contaminants including nutrients such as nitrates and phosphates (Asamoah & Amarin, 2011).

Nutrients are essential for healthy plant and animal populations and provide a range of

benefits, including increased food production for a growing global population. However, too much nutrient can have adverse effects on water quality, drinking water sources, recreation, and aquatic life (USGS, 2010). Effective crop production depends on an adequate supply of nutrients to the crops through fertilizer application to achieve maximum yield. However, soil nutrients need to be managed properly to meet the fertility requirements of crops without adversely affecting the quality of our valuable water resources. On a typical dairy farm, all nutrients are cycled from the soil to crops, to animals, and finally back to the soil as manure. However, nutrient recycling on almost all farms is not a closed-loop system and nutrients leave the farm in various ways and lost into the environment through soil erosion, phosphorus runoff, nitrogen leaching, and manure ammonia volatilization which not only make nutrient cycling inefficient but also impose negative impacts on water quality. (CAFÉ, 2017).

The inorganic contaminant of greatest concern in groundwater is the nitrate ion, (NO_3^-) which commonly occurs in both rural and suburban aquifer (Baird & Cann, 2008). Excessive nitrate in drinking water can lead to methemoglobinemia also known as “blue-baby syndrome,” in which oxygen levels in the blood become too low, sometimes with fatal results (Baird & Cann, 2008; USGS, 2010). Concentrations of dissolved phosphorus in groundwater typically are low because phosphorus tends to sorb to soil and aquifer sediments and is not readily transported in groundwater (Holman *et al.* 2008) The principal sources of phosphorus to groundwater systems include overlying soils, dissolution of minerals that contain phosphate in aquifer sediments, agricultural fertilizer, animal waste, and leaking septic systems or infiltration of wastewater (Fuhrer *et al.* 1999). The Pra Riv-

er and its tributaries constitute a major source of water supply to communities within the basin. The major tributaries are perennial and constitute all-year-round reliable water source and recharge resource for groundwater (WRC, 2010). The Pra Basin is one of the most extensively and intensively used river basin areas in Ghana in terms of settlement, agriculture, logging and mining (WRC, 2010). The basin contains most of the large cocoa growing areas in the Eastern, Ashanti, and Central regions. Tree cash crop cultivation other than cocoa includes mainly oil palm with a lot of large commercial plantations (WRC, 2010). Also, food crops such as cassava, yam, cocoyam, plantain, as well as fruits such as banana, oranges, etc, are produced together with cocoa for subsistence with the use of inorganic fertilizers and other agrochemicals (Tay *et al.*, 2013; Tay, 2015). Knowledge on the possible future threat posed by nutrient pollution such as nitrate and phosphate as a result of the use of inorganic fertilizers and agrochemicals for purposes of farming within the basin is limited. However, Ahialey *et al.* (2010); Tay *et al.*, (2013) did some work on general routine water quality within the basin. It is for this reason that this paper seeks to identify the level and source of nutrient pollution on the groundwater as well as any possible health hazards associated with it.

Experimental

Study area

The Lower Pra Basin lies within $05^{\circ}0'00''$ and $06^{\circ}0'00''$ N and $01^{\circ}0'00''$ and $02^{\circ}0'00''$ (Fig.1). The climate is sub-equatorial wet, with two rainy seasons (May-July and September-November) (Dickson & Benneh 2004). The mean annual rainfall is relatively high, about 1,500mm but is also very variable, ranging between 1300 mm and 1900 mm. The basin is

warm and moist throughout most of the year. The relative humidity is between 70% - 80% throughout the year (Dickson & Benneh 2004). In the drier seasons, temperatures are around 26°C in August and 30°C in March (WRC, 2012). It is made up of four major tributaries namely the Ofin, Oda, Anum and Birim. These tributaries merge into the lower Pra which then flows into the Gulf of Guinea through its estuary at Shama (GWSC, 1989). The basin lies completely within the forest ecological zone of Ghana and has a moist semi-deciduous

forest with valuable timber species (Dickson & Benneh 2004). Owing to the development of the cocoa industry, the initial forest has been modified to a secondary forest consisting of shrubs, soft woody plants and climbers (Dickson & Benneh, 2004). The basin is principally dominated by the forest ochrosols, and to a lesser extent, forest ochrosol-oxysol intergrade (Dickson & Benneh, 2004; GWSC, 1989). The ochrosols are highly coloured soils with little leaching characteristics (Dickson & Benneh, 2004).




Fig.1: Map showing sampling communities and borehole locations (Map of Ghana Insert) within the different geological settings.

Geology and soil

The basin is predominantly underlain by granite rocks of the Cape Coast and Dixcove granitoids (Fig.1). Various portions of the basin are also underlain by metasediments of Birrimian and Tarkwain both of the pre-cambrian origin (Kesse, 1985). The basin is principally dominated by the forest ochrosol-oxysol intergrades, which also contains more nutrients and are alkaline. They have rather variable drainage properties and therefore require protection against erosion. In the lower Pra nearer the coast, further classification includes the acid glosol and coastal savannah soil groups (GWSC, 1989).

Aquifer characteristics

Borehole depths of the study area are generally shallow ranging between 30 to 96 metres, with a mean value of 44.42m. Borehole yield is generally low and largely variable, ranging from 0.4 to 51.7 m³h⁻¹ and mean value of 4.55 m³h⁻¹ with schist and granite aquifers having relatively higher yields (Tay *et al.*, 2013, Dorleku 2013). The fractures in the rocks are generally open. The granite and schist rocks are exposed whilst; the Birimian and Takwian rocks have thick overburdens. The soils characteristically developed over the same kind of highly weathered parent material with lateritic to clayey topsoil layer and thickness which range from 4 to 14 m, which may, however, extend further in some areas. The static water levels of the boreholes generally range from 0.4 to 22.4 m with a mean of 3.7 m (Tay *et al.*, 2013).

Land Uses

The Pra Basin, in general, has considerable potential for development in agriculture, forestry, tourism, and mining, and provides

livelihoods for many through these. Over 63% of the population is engaged in the Agriculture and related sectors of the economy. Both commercial and subsistence agricultural activities are practiced, with cocoa being the main commercial crop (WRC, 2012). In the districts where large scale cocoa farming is practiced within the basin, annual returns in terms of family income are high and the standard of living is appreciable. In the Benso area of the Western region commercial oil palm plantation is prevalent (WRC, 2012). It was observed that small-scale palm oil industries are situated around Assin North and Benso areas because of large oil palm plantations (Dorleku, 2013).

Sampling and analysis

Water samples were collected from 65 boreholes in about 45 communities. The samples were collected two times for each season, January and April 2012 for the dry season and June and October for the wet season into previously sterile 1 Litre plastic bottles. About 500 litres of water was pumped out of the boreholes and the bottles rinsed with the borehole water just before each sample was taken. Temperature, pH, and conductivity were measured in the field using Hach Sens ion 156 meter. The samples were transported to the CSIR-WRI laboratory in Accra at a temperature of about 4°C in an ice chest with ice cubes and kept in the refrigerator until analysed. Nitrate was determined by hydrazine reduction method using T60 UV spectrometer at wavelength of 520 nm (APHA, 1998). Phosphate and sulphate were determined by stannous chloride and turbidimetric method at a wavelength of 690 nm and 420 nm respectively (APHA, 1998).

Quality Control

To ensure the reproducibility of the data, duplicate analysis was carried out for every batch of five samples. Blanks were analysed together with samples to check for

contamination. Duplicates of 0.5, 1.0, and 10 mg/l control standards from high-purity salts of nitrate, phosphate and sulphate respectively were prepared and analysed together with the samples to check for accuracy.

TABLE 1A
Seasonal nutrient concentration in individual boreholes in the basin

Sample Source	pH Sample ID	Dry		Temp °C		EC µS/cm		PO4-P mg/l		SO4 ²⁻ mg/l		NO ₃ -N mg/l	
				Dry	Wet	Wet	Dry	Wet	Dry	Wet	Dry	Wet	DryWet
Sabina	BH1	4.9	6.45	26.9	28.2	270	180	0.200	0.368	18.8	19.1	0.392	0.111
Ayitey	BH2	5.3	5.94	27.6	31.8	200	180	0.180	0.373	8.9	9.2	0.335	0.408
Nkrafo	BH3	5.2	6.87	27.1	30.1	360	260	0.060	0.278	4.7	31.3	0.192	0.011
Nkrafo	BH4	5.9	6.33	26.3	27.6	180	150	0.030	0.473	10.9	11.4	1.96	0.327
Obirikwaku	BH5	6.2	6.05	25.7	29.5	260	210	0.180	0.416	18.4	18.5	0.435	0.178
Odumasi	BH6	5.8	5.56	26.2	27.5	150	110	0.320	0.290	11.8	12.1	2.78	1.37
OdumasiCamp	BH7	6.2	5.67	26.9	27.7	300	200	0.410	0.303	6.2	30.9	0.666	0.353
Obobakokrowa	BH8	6.9	5.24	25.6	28.7	290	210	0.160	0.221	43.7	6.5	0.481	0.482
Dwedaama	BH9	6.4	6.26	25.7	27.3	240	180	0.560	0.234	35.0	35.0	0.243	0.071
Dwedaama	BH10	5.3	5.37	26.7	26.3	90	70	0.250	0.316	14.0	14.4	0.482	0.34
Worakase Habitat	BH11	5.8	4.73	26.5	25.7	80	80	0.540	0.293	7.4	7.3	2.03	1.55
Brofoyedru Habitat	BH12	5.2	4.47	28.2	25.6	120	100	0.570	0.501	17.9	17.6	0.759	0.05
Assin Nyankumase	BH13	6.9	4.90	29.2	27.6	470	70	0.160	0.496	55.6	56.9	3.22	0.073
Assin Nyankumase	BH14	7.3	5.09	28.5	28.5	180	380	0.351	0.288	18.1	18.9	0.582	1.29
Assin Nyankumase	BH15	7.9	4.69	28.1	27.8	70	160	0.643	0.342	18.7	18.1	5.01	1.45
Akonfudi	BH16	6.5	5.07	29.3	26.8	290	230	0.030	0.208	10.9	28.7	0.301	0.341
Akunfudi	BH17	6.3	5.14	29.1	27.2	500	360	0.180	0.298	18.4	25.6	0.677	0.391
Assin Breku(SDA)	BH18	6.0	6.41	30.5	26.4	140	260	0.410	0.257	4	43.2	0.753	0.053
Assin Breku(Gyidi)	BH19	6.3	5.53	28.7	27.3	290	230	0.210	0.373	43.7	6.14	0.986	0.518
Assin Breku	BH20	5.7	5.70	28.3	28	500	480	0.380	0.345	6	5.69	0.272	0.053
Techiman No 1	BH21	6.1	5.12	29.1	26.9	60	60	0.400	0.278	31.7	12.8	0.188	0.451
Kwame Ankra	BH22	6.5	5.35	28.0	26.2	143	110	0.260	0.429	35.0	9.36	0.372	0.295
Ninkyiso	BH23	5.8	4.92	28.2	26.9	210	170	0.250	0.308	14.0	12.4	0.755	0.295
Amoakokrom	BH24	5.8	4.96	28.8	27.3	320	240	1.200	0.334	24.1	23.9	0.622	0.251
Nyamebekyere	BH25	5.6	5.63	27.2	26.8	170	130	1.900	0.010	7.4	3.6	0.732	0.462
Jerusalem	BH26	6.1	5.03	30.4	26.6	80	70	3.080	0.380	31.8	7.5	0.918	0.758
Antoabasa	BH27	5.9	5.34	30.2	28.3	170	140	0.490	0.632	21.9	31.6	0.702	0.197
Antoabasa	BH28	5.9	5.08	29.2	27.6	220	170	0.270	0.506	18.3	26.6	0.482	0.352
Bedidua	BH29	6.5	5.15	28.8	28.3	220	170	0.480	0.458	10.3	18.1	0.469	0.114
Anum	BH30	6.0	—	28.3	—	270	—	0.340	—	16.7	—	0.198	—
Kyeikrom	BH31	5.6	5.70	31.6	31	150	120	0.340	0.300	3.6	5.2	0.37	0.43
Adukrom	BH32	5.9	—	27.1	—	140	—	0.340	—	22.2	—	0.717	—
Subrisu	BH33	6.0	—	27.0	—	240	—	0.140	—	24.2	—	0.28	0.114

TABLE 1B
Seasonal nutrient concentrations in individual boreholes continued.

Nsukyir	BH34	5.9	5.4	26.5	27.3	170	170	0.670	0.733	22.2	22.0	0.499	0.205
Denyase Domeabra	BH35	5.6	5.98	28.0	27.1	220	160	0.740	0.78	9.9	15.5	0.328	0.123
Twifo Mampong	BH36	6.0	4.24	28.6	29.6	279	220	0.669	0.532	23.8	77.9	0.671	0.255
Twifo Mampong	BH37	6.4	5.90	30.7	29.1	1140	590	0.558	0.92	13.6	77.8	0.36	0.16
Akwa Yaw	BH38	6.6		29.3		130		0.161		10.4		0.961	
Bremang	BH39	6.3	6.18	29.8	31.2	310	260	0.271	0.109	22.6	76.2	0.339	0.079
Bremang	BH40	6.0	6.06	29.2	31.4	240	200	0.268	0.938	3.6	44.6	0.412	0.142
Twifo Agona	BH41	5.9	4.96	28.8	29.3	190	140	1.990	0.175	7.5	22.4	0.205	0.091
Somnyamekordru	BH42	5.3	6.24	26.8	28.1	710	410	0.669	0.866	13.6	13.8	0.242	0.146
Somnyamekordru	BH43	6.3	5.95	27.4	28.4	824	130	0.558	0.137	10.4	10.4	0.369	0.609
Atu Krom	BH44	5.6	4.95	29.5	28.1	350	240	0.429	0.156	58.7	58.2	0.255	0.163
Subresu	BH45	6.1	5.29	28.1	28.4	190	200	0.156	0.115	62.1	61.9	0.294	0.071
Gromsa	BH46	6.9	4.40	28.9	27.1	190	150	0.271	0.509	75.7	23.6	0.346	0.23
Anyinase Ankase	BH47	6.4	4.66	28.5	26.3	100	90	0.268	0.488	9.1	9.7	0.387	0.141
Semchem	BH48	6.6	4.88	29.9	29.6	560	200	0.362	0.491	91	77.5	0.349	0.388
Semchem	BH49	6.1	4.56	27.2	27.6	400	300	0.437	0.506	31.9	31.2	0.304	0.151
Mampong	BH50	5.3	5.02	26.4	27.6	110	100	0.502	0.707	1.6	32.7	0.718	0.451
Essamang	BH51	5.8	5.6	26.7	27.6	60	50	0.210	0.121	32.2	14.4	2.24	0.038
Achim/Akyem	BH52	6.5	6.5	26.8	26.4	423	417	0.37	0.34	26.9	23.9	0.33	0.32
Achim/Akyem	BH53	6.7	6.5	27.2	26.6	371	368	0.33	0.32	27.2	25.2	0.15	0.17
Achim/Akyem	BH54	6.3	6.1	26.1	27.2	453	449	0.32	0.33	35.4	32.9	2.08	2.01
Ayiem	BH55	5.7	5.9	28.5	28.5	140	137	0.82	0.75	76.3	77.1	0.59	0.62
Botawjwina	BH56	6.7	6.5	29.2	28.8	615	605	0.29	0.35	62.4	69.4	0.26	0.32
Botawjwina	BH57	6.4	6.2	28.7	28.8	923	906	0.33	0.24	96.2	79.1	0.54	0.49
Benso	BH58	5.6	6.3	27.8	27.5	156	521	0.44	0.42	13.9	78.9	0.28	2.61
Benso	BH59	5.2	5.4	27.3	28.1	412	153	0.42	0.37	72.3	14.9	2.71	0.50
Benso	BH60	6.1	5.6	29.2	27.4	527	403	0.57	0.38	73.9	70.3	0.40	
Adum Banso	BH61	5.6	5.8	26.5	27.8	411	402	0.34	0.33	52.4	50.4	0.34	0.44
Adum Dominase	BH62	5.5	5.7	28.2	28.2	343	339	0.44	0.40	21.6	21.6	0.63	0.67
Adum Dominase	BH63	5.9	5.7	28.7	27.8	93.4	90.9	0.40	0.39	14.2	13.9	0.73	0.69
Adum Trebuom	BH64	6.1	6.0	27.1	26.6	196	188	0.53	0.58	24.2	22.5	0.51	0.55
Adum Trebuom	BH65	6.4	6.6	28.3	26.9	333	327	0.41	0.40	22.2	20.2	0.36	0.30
Min Value		4.9	4.24	25.6	25.6	60	50	0.030	0.010	1.6	3.6	0.15	0.011
Max Value		7.9	6.87	31.6	31.8	1140	906	3.080	0.938	96.2	79.1	5.01	2.61

TABLE 2
Mean Dry Season statistical summary of data in mg/l- unless specified

n=65	MIN	MAX	Mean	Median	SD	WHO
pH (pH units)	4.90	7.90	6.06	6.00	0.54	6.50-8.50
EC ($\mu\text{s}/\text{cm}$)	60.0	1140	291	240	208	-
$\text{NO}_3^- \text{-N}$	0.15	5.01	0.75	0.47	0.85	10
$\text{PO}_4^{3-} \text{-P}$	0.03	3.08	0.47	0.36	0.20	-
SO_4^{2-}	1.6	96.2	27.0	18.8	22.7	250

TABLE 3
Mean Wet Season statistical summary of data in mg/l-unless specified.

n=65	MIN	MAX	Mean	Median	SD	WHO
pH (pH unit)	4.24	6.87	5.5	5.59	0.63	6.50-8.50
EC ($\mu\text{s}/\text{cm}$)	50	906	236	188	159	-
NO_3^- -N	0.011	2.61	0.43	0.32	0.49	10
PO_4^{3-} -P	0.010	0.938	0.40	0.37	0.20	-
SO_4^{2-}	3.60	79.1	30.1	22.4	23.2	250.

TABLE 4
Statistical summary of data from the Basin

n= 65	Min.	Max.	Mean	Median	Stdev.	P-Value	W.H.O(2004)
Temp($^{\circ}\text{C}$)	25.6	31.8	28	27.9	1.33	–	–
PH (pHunits)	4.24	7.90	5.80	5.79	0.58	–	6.5-8.0
EC ($\mu\text{s}/\text{cm}$)	55	1140	264	214	183	–	–
NO_3^- -N	0.08	5.01	0.59	0.39	0.67	0.011	10
PO_4^{3-} -P	0.01	3.08	0.44	0.36	0.34	0.225	-
SO_4^{2-}	1.6	96.2	28.5	20.6	22.9	0.443	250




Fig. 2: Box plot of nutrient concentrations of boreholes in the Pra basin

Results and Discussion

Table 1 represents the mean seasonal nutrient concentrations in groundwater within the Lower Pra Basin. Tables 2 and 3 represent the statistical summary of dry and wet seasons

respectively whilst Table 4 shows the summary of statistical data from the basin.

The results show that the mean concentration of the parameters as compared to the WHO(2006) guideline values are low and do not pose an immediate threat to human health.




Fig. 3: % stacked Histogram of dry and wet seasons for Phosphate in the basin.




Fig. 4: % stacked Histogram of dry and wet seasons for Nitrate in the basin.




Fig. 5: Bar graph of mean wet and dry season nutrient concentrations.

The concentration of each parameter as related to each other is illustrated by the box plot in fig 2. It can be seen from the box plot that sulphate is largely variable and have relatively large range as compared to the other two (which are nutrients).




Fig. 6: Scatter plot showing correlation between borehole depth and NO₃ concentration.




Fig. 7: Scatter plot showing correlation between borehole depth and PO₄³⁻P concentration.

Nitrate

Nitrate is often a major health concern in the developed countries as a result of well-documented pollution of groundwater from the use of nitrate-based fertilizers and a wide spectrum of synthetic pesticides for improving crop yield, leading to methaemoglobinaemia and stomach cancers (Edmunds & Smedley, 1996; WHO, 2011). Nitrate is very soluble and although some may be used by plants, much of the dissolved nitrate escapes unused into deeper parts of the soil and groundwater and can be toxic to humans at concentrations between 10 to 15 mg/l (Egboka *et al.*, 1989). Dry season values range from 0.15 to 5.01 mg/l with the wet season recording values between 0.011 and 2.61 mg/l (Tables 2 and 3). None of the boreholes recorded values above the W.H.O (2006) value of 10 mg/l. However, the highest value of 5.01 mg/l was recorded in the dry season at Assin Nyankumasi (BH 15). Field observations within the study area indicate that the borehole located at Assin Nyankumase (BH15) is sited very close to a public latrine, which might have contributed to the relatively high nitrate value. The highest mean wet season value (2.60 mg/l) was recorded at Benso (BH58), where there are commercial oil palm plantations and therefore might have been influenced by fertilizer usage. Ahiale et al, (2010), recorded values between 0.089 mg/l and 9.03 mg/l within the basin. The mean and median values recorded for the dry season were 0.75 and 0.47 mg/l respectively whilst that of the wet season recorded values of 0.43 and 0.32 mg/l respectively (tables 2 and 3) and further shown by the bar graph (fig 4). Mean dry season values were generally higher than wet season values, Fig 5. The relative lower values for the wet season might have been the result of rainfall precipitation and recharging of aquifers by freshwater culminating in the

dilution of concentration levels. There is a significant difference between the dry and wet season values ($p < 0.05$), Table 4. There was, however, a very weak positive correlation ($R^2 = 0.122$) between borehole depth and nitrate concentration, Fig 6, indicating that nitrate concentration in the aquifers might be as a result of natural biogeological processes, rather than from anthropogenic sources i.e fertilizer and synthetic pesticide inputs. Concentrations of nitrate generally decrease with good depth (USGS, 2010). This is because (1) recharge of deep, old groundwater most likely occurred when anthropogenic nitrogen inputs were relatively low; (2) the amount of time required for microbial processes such as denitrification to take place increases as water moves downward through the aquifer; and (3) deep groundwater is commonly a mixture of water that has moved along flow paths that originate in areas of differing land uses and nitrogen inputs (USGS, 2010). Aquifers within the basin are not under threat from agriculture activities.

Phosphate

Phosphorus is a nutrient required by all organisms for the basic processes of life. Weathering of certain minerals particularly apatite provides a natural source of dissolved phosphorus in groundwater (Hitchon *et al.*, 1999). Phosphorus in natural waters is usually found in the form of phosphates (PO_4^{3-}) (UNEP GEM, 2006). Orthophosphates applied to agricultural or residential cultivated land as fertilizers are carried into surface waters with storm runoff which eventually ends up in groundwater (APHA, 1998). Dry season values ranged from 0.03 to 3.08 mg/l whilst the wet season recorded values between 0.010 and 0.938 mg/l. The W.H.O has not given any guideline value for phosphate, but

the U.S.EPA recommended a safe value of 2.5 mg/l. Only one borehole (BH26) with phosphate concentration 3.08 mg/l in the dry season recorded value above the U.S.EPA recommended a value of 2.5 mg/l. Mean and median values recorded for the dry season were 0.47 and 0.36 mg/l respectively and that of wet season was 0.40 and 0.37 mg/l respectively. There is no significant difference between dry and wet season values ($p > 0.05$) (Table 4 and Fig 5). Hence natural geologic sources might have a greater influence on concentrations in groundwater than anthropogenic sources. Concentrations of dissolved phosphorus (DP) in groundwater typically are low because phosphorus tends to sorb to soil and aquifer sediments and is not readily transported into groundwater (Holman et al. 2008). There is a very weak negative correlation between borehole depth and phosphate concentration, ($R^2 = 0.025$), Fig 7 within the Pra basin. United State Geological Survey (USGS), 2010; Dubrovsky et al. 2010 reported that dissolved phosphorus concentrations in groundwater showed no correlation to fertilizer and manure use in agricultural areas in California and other regions; more so, there was similarity of concentrations in both shallow and deep groundwater and therefore concluded that natural geologic sources of phosphorus may have a greater influence on concentrations in groundwater than do human-derived sources from activities on the land surface. Hence natural geologic factors might have contributed to phosphorus levels in the aquifers within the Pra basin.

Sulphate

Sulphate is an abundant ion in the earth's crust and high concentrations may be present in winter due to leaching of gypsum, sodium sulphate, and some shales (UNEP-GEM,

2006). As a result of oxidation of pyrites and gypsum mine drainage may contain high concentrations of sulphate, leading to groundwater contamination. Fertilizers and natural sources, such as volcanoes also contribute to sulphate presence in groundwater (MPCA, 1999). The dry season recorded 27.0 and 18.8 mg/l for mean and median respectively whilst that of the wet season was 30.1 and 22.4 mg/l respectively. Higher mean wet season value is in agreement with literature as stated earlier. Sulphate values for both seasons ranged between 1.6 to 96.2 mg/l within the basin, and thus below the W.H.O (2006) permissible limit of 250mg/l (Table 4). There is no significant difference however, between the dry season and wet season values with $P > 0.05$, (Table 4). Acid mine drainage might not yet be contributing significantly to the aquifers within the basin.

Conclusion

Nitrate values recorded in the basin were low, with mean dry and wet season values of 0.75 and 0.43 mg/l respectively, far below the W.H.O recommended a value of 10.0 mg/l for drinking water. The contribution of synthetic fertilizers and other anthropogenic activities does not seem to pose a threat to groundwater in the basin. There was a very weak positive correlation between borehole depth and nitrate concentration, indicating that nitrate concentration in the aquifers might be predominantly as a result of natural geological processes. Mean phosphorus value for both seasons was 0.44 mg/l, even though one borehole recorded an elevated mean dry season value of 3.08 mg/l which is above the USEPA recommended value of 2.5 mg/l. There is a very weak negative correlation between borehole depth and phosphate concentrations. Hence natural geologic sources might have a greater

influence on concentrations in groundwater than anthropogenic sources. Phosphorus does not pose a threat to aquifers in the basin. Even though the wet season recorded higher values for sulphate than the dry season, there was no significant difference between them. Sulphate values for both seasons ranged between 1.6 to 96.2 mg/l in the basin, all below the W.H.O permissible limit of 250 mg/l. Geological processes such as leaching of certain rocks might be the contributing factor in the aquifers rather than fertilizer sources.

References

- AHIALEY, E. K., SERFOH-ARMAH, Y., & KORTATSI, B. K. (2010) Hydrochemical analysis of groundwater in the Lower Pra Basin. *J. Water Resource and Protection* **2**, 864 – 871.
- APHA (1998) “*Standard Method for the Examination of Water and Wastewater*,” 20th Edition, American Public Health Association, Washington D.C. 1998.
- ASAMOAH, D. N. & AMORIN, R. (2011) Assessment of the quality of bottled/sachet water in the Tarkwa-Nsuaem Municipality(TM) of Ghana. *Research Journal of Applied Science Engineering and Technology* **3**, 377 – 385.
- BAIRD, C. & CANN, M. (2008) *Environmental Chemistry* W.H. Freeman and Company Fourth edition, New York Press. 301 – 302.
- CAFÉ, (2017) The Centre for Agriculture, food and the Environment, University of Massachusetts Amherst. <https://ag.umass.edu/crops-dairy-livestock-equine/fact-sheets/nutrients-water-quality>. Accessed on 4th April, 2017.
- DICKSON, K. B., & BENNEH, G. (2004) *A New Geography of Ghana*. Fifth impression, Revised edit. Longman, London
- DORLEKU, M. (2013) ‘Groundwater Contamination With Toxic Metals Through Smallscale Mining Within the Lower Pra Basin’, Ghana. MPhil. Thesis, University of Ghana, Legon - Accra, Ghana.
- DUBROVSKY, N. M., BUROW, K. R., CLARK, G. M., *et al.*, (2010) The quality of our Nation’s waters—Nutrients in the Nation’s streams and groundwater, 1992–2004: US Geol Surv Circ 1350, 174 p
- EGBOKA, B. C., NWANKWOR, G. I., ORAJAKA, I. P. *et al.*, (1989) Principles and Problems of Environmental Pollution of Groundwater Resources with Case Examples from Developing Countries. *Environmental Health Perspectives* **83**, 39 – 68
- FUHRER, G. J., GILLIOM, R. J., HAMILTON, P. A., *et al.*, (1999) The quality of our nation’s water: nutrients and pesticides. US Geol Surv Circ 1225, 82p.
- HITCHON, B., PERKINS, H. E. & GUNTER, W. D. (1999) *Introduction to Ground Water Geochemistry*. Geoscience Publishing Ltd., Sherwood Park, Alberta, Canada. pp 310.
- HOLMAN, I. P., WHELAN, M. J., HOWDEN, N. J. K., *et al.*, (2008) Phosphorus in groundwater – an overlooked contributor to eutrophication? *Hydrological Processes* **22**, 5121 – 5127.
- HYNDS, P. D., THOMAS, M. K., & PINTAR, K. D. M. (2014) Contamination of groundwater systems in the US and Canada by Enteric Pathogens, 1990–2013: a review and pooled-analysis. *PLoS ONE* **9** (5), e93301. doi:10.1371/journal.pone.0093301.
- KESSE, G. O. (1985) *The mineral and rock resources of Ghana*. A.A/ Balkema, Rotterdam/Boston.
- KORTATSI, B. K. (2006) Concentration of Trace Metals in Boreholes in the Ankobra Basin, Ghana. *West African Journal of Applied Ecology* **10**, 73 – 87.
- MPCA (1999) Minnesota Pollution Control Agency, Environmental Outcomes Division

- MUNN, M. D., NOLAN, B. T., PUCKETT, L. J., *et al.*, (2010) The quality of our Nation's waters — Nutrients in the Nation's streams and groundwater, 1992 – 2004: US Geol Surv Circ 1350, 174 p.
- PRÜSS-ÜSTÜN, A., BOS, R., GORE, F. & BARTRAM, J. (2008) *Safer Water, Better Health: costs, benefits and sustainability of interventions to protect and promote health*. Geneva: World Health Organization.
- SMEDLEY, P. L., EDNUNDS, W. M., WEST, J. M., *et al.*, (1995) Vulnerability of Shallow Groundwater Quality due to Natural Geochemical Environment. Health Problems related to groundwater in the Obuasi and Bolgatanga areas, *Ghana British Geological Survey*, 130 pp (WC/95/043 Vol. 2) (unpublished)
- TAY, C., HAYFORD, E., HODGSON, I. O., *et al.*, (2013) "Hydrochemical appraisal of groundwater evolution within the Lower Pra Basin, Ghana: a hierarchical cluster analysis (HCA) approach". *Environ Earth Sci* DOI 10.1007/s12665-014-3644-4.
- TAY, C. (2015) Hydrogeochemical Processes Influencing Groundwater Quality Within The Lower Pra Basin, Ghana. PhD Thesis, University of Ghana, Legon-Accra, Ghana
- UNITED NATIONS ENVIRONMENT PROGRAMME (UNEP) (2006) Global Environment Monitoring System/Water Programme.
- USGS (2010) Nutrients in the Nation's Streams and Groundwater, 1992 – 2004. National Water-Quality Assessment Program, *U.S. Geological Survey*, Reston, Virginia
- WHO (2006). *Guidelines for drinking-water quality*, 3rd ed. Vol.1. Recommendations, World Health Organization, Geneva.
- WHO (2011) Nitrate and Nitrite in Drinking-water Background document for development of WHO *Guidelines for Drinking-water Quality*. World Health Organization, Geneva.
- WATER RESOURCES COMMISSION (WRC), (2012) Pra River Basin-Integrated Water Resources Management Plan, June 2012.

Received 01 Nov 18; revised 02 Jul 19.