




Clinical indications for plain abdominal radiographs: A survey study among radiographers



Authors:

Lynn Burrell¹ 
Razana Williams¹ 
Wilma ten Ham-Baloyi² 

Affiliations:

¹Department of Radiography,
Faculty of Health Sciences,
Nelson Mandela University,
Gqeberha, South Africa

²Department of Nursing
Science, Nelson Mandela
University, Gqeberha,
South Africa

Corresponding author:

Wilma ten Ham-Baloyi,
Wilma.tenHam-Baloyi@
nmmu.ac.za

Dates:

Received: 07 Dec. 2022
Accepted: 17 May 2023
Published: 29 Aug. 2023

How to cite this article:

Burrell, L., Williams, R. & ten
Ham-Baloyi, W., 2023,
'Clinical indications for plain
abdominal radiographs: A
survey study among
radiographers', *Health SA
Gesondheid* 28(0), a2289.
[https://doi.org/10.4102/
hstag.v28i0.2289](https://doi.org/10.4102/hstag.v28i0.2289)

Copyright:

© 2023. The Authors.
Licensee: AOSIS. This work
is licensed under the
Creative Commons
Attribution License.

Read online:



Scan this QR
code with your
smart phone or
mobile device
to read online.

Background: Abdominal pain is a common complaint in the Emergency Department. Radiographers' knowledge and practices regarding clinical indications for performing abdominal examinations are crucial in the results radiographs produced.

Aim: To determine the knowledge and practices of radiographers regarding the clinical indications for performing radiographic examinations of the abdomen.

Setting: Four public hospitals in the Eastern Cape province, South Africa.

Methods: A cross-sectional study was conducted, using a convenience, all-inclusive sample of $n = 85$ radiographers. A hard copy self-administered questionnaire was distributed between February and June 2020. Descriptive (mean and standard deviations) and inferential (chi² test) statistics were generated using IBM® SPSS® version 26.0 software package.

Results: Knowledge of clinical indications had a mean of 59.41. All four demographics (age, years of experience, attended a short course and attended pattern recognition course) were significantly associated with overall knowledge. Additionally, short course attendance was significantly associated with most practice items, and two knowledge items (which views are done for perforation; and which view(s) demonstrate a stab abdomen). Pattern recognition was significantly associated with one knowledge item (which views are regarded as an acute abdomen).

Conclusion: Despite the lack of short courses and pattern recognition courses, radiographers' knowledge of clinical indications was good (>50%). Continuous training, accessible protocols for performing clinical indications for plain abdominal radiographic examinations for radiographers, audit, feedback and reminders to enhance protocol adherence are recommended.

Contribution: The study findings could be used to enhance knowledge and practices regarding clinical indications for plain abdominal radiographic examinations among radiographers.

Keywords: clinical indications; diagnostic radiography; radiographic examinations; abdominal radiographs; radiographers; knowledge; practices; radiographic image.

Introduction

Plain abdominal radiography is the initial imaging modality of choice when it comes to making a diagnosis for the abdomen (Tomizawa et al. 2017). It is defined as a means of obtaining imaging of the internal structures found in the abdominal cavity, by passing x-rays through them and recording the shadows cast by these structures (Lau 2007). Despite advancements in radiology such as computed radiography and ultrasound, plain abdominal radiography, using x-rays, still plays a significant role in the imaging protocol of the abdomen (Bell 2022). Patel, Moran and Nakada (2018) state that plain abdominal radiography is readily available and has the additional benefits of a lower radiation dose and lower cost factor to the patient.

Abdominal problems are common clinical complaints in the Emergency Department (ED) and account globally for 8% of all patient visits to the ED (Meltzer et al. 2017). These abdominal problems are wide and range from surgical to non-surgical conditions and from self-resolving to life-threatening conditions (Patterson, Kashyap & Dominique 2021; Velissaris et al. 2017). The clinical indications for abdominal radiographic examinations are normally grouped into suspected bowel obstruction, perforation, suspected foreign body, moderate to severe undifferentiated abdominal pain and renal tract calculi follow-up (Rink & Wessels 2021). However, plain imaging

radiographs of the abdomen and chest are commonly requested for acute medical emergencies in patients who present with non-specific abdominal signs and symptoms (Fox 2017).

Gleadle, Li and Yong (2017) state that there are many diseases that could cause acute abdominal pain or acute abdomen and each of these has a specific imaging protocol to follow. Thus, the radiographer needs to have the necessary knowledge about the clinical indications as well as the protocols in order to assist in the diagnosis of the patient (European Society of Radiology 2019). Furthermore, the most appropriate abdominal protocol should be followed in order to demonstrate the pathology in question and, by applying the knowledge to practice, mismanagement and treatment delays are minimised, enhancing patient safety (Gleadle et al. 2017). It is important for the radiographer to follow the distinct protocol for each indication so that pathology is not overlooked. Radiographs allow the physician to see the pathology presented and when an incorrect protocol is followed, it leads to delay in treatment and risks that could include severe untreated pain, additional complications and even death of the patient, which compromises patient safety (Vom & Williams 2017). Furthermore, a protocol allows radiographers to accept a more autonomous role that cultivates critical thinking, reflection and research-informed decision making when performing or justifying performance of radiographic examinations (Vom & Williams 2017). Various protocols for abdominal radiography (indications) exist, including the Australian Diagnostic Pathways, The British Royal College, The American College of radiology, The French National Authority for Health and The Royal College of Radiologist Guidelines, which differ slightly from each other. Radiographers in South African public hospitals follow the provincial Department of Health's protocol that lists the following indications: perforated abdomen, stab abdomen, subphrenic abscess and renal calculus (Department of Health 2010). However, as observed by the first author, there was evidence of radiographers in the study setting not adhering to this protocol by conducting inadequate projections when performing radiographic examinations of the abdomen. Non-adherence to protocols regarding plain abdominal examinations by radiographers could be a result of limited knowledge and practices, which ultimately impacts patients negatively thus affecting the safety and treatment plan of the patient (McFadden et al. 2018). Although studies have been conducted regarding the radiographers' perspectives, knowledge and practices on x-ray examinations in Europe (McFadden et al. 2018; Wit, Vroonland & Bijwaard 2022), little is known on the knowledge and practices of radiographers with regard to the clinical indications for performing radiographic examinations of the abdomen, as evidenced by a paucity of articles in this regard found globally, including South Africa. This study, therefore, aims to determine the knowledge and practices of radiographers with regard to the clinical indications for performing radiographic examinations of the abdomen at public hospitals in the Eastern Cape, South Africa.

Methods

This quantitative study used a cross-sectional survey design and was conducted by the first author under the supervision of the second and third authors. The study was part of a bigger study that was conducted to develop evidence-based recommendations regarding abdominal radiographic imaging practices for radiographers.

In the Eastern Cape, there are five radiology departments in five public hospitals, respectively. The main study was conducted at radiology departments in four public hospitals in the Eastern Cape, South Africa. The four hospitals were selected as they perform a high number of abdominal examinations. Radiographers employed in these hospitals have been trained in plain abdominal radiographs and the clinical indications for these and, while qualified, receive refresher short courses on performing abdominal examinations and pattern recognition as part of their continuous professional development.

To obtain as large as possible sample size, an all-inclusive, convenience sampling method was used to select the radiographers for the study.

Survey questionnaire

The data collection instrument was a self-administered questionnaire, developed based on a literature review and aligned with the provincial Department of Health's protocol (Department of Health 2010). Kindly note that this protocol may differ from other, international protocols. The questionnaire consisted of the following three sections:

- Section A – demographics, including age, years of experience, attendance at short courses and pattern recognition training (four questions).
- Section B – knowledge of radiographers with regard to clinical indications when performing abdominal radiographic examinations (eight multiple-choice questions).
- Section C – practices of radiographers with regard to the clinical indications when performing abdominal radiographic examinations (nine questions: five 5-point Likert Scale questions and four multiple-choice questions)

As part of the process to validate the questionnaire, a pilot test was conducted in a radiography department in one purposively selected hospital as this was the hospital with the lowest number of staff, consisting of 12 radiographers. The data obtained from the pilot study were used within the main study to enhance the sample size as no changes were made to the questionnaire after the pilot study was conducted.

After permission for the study was obtained, the managers at the hospitals (who acted as the gatekeepers) were informed of the study. After they agreed with the terms of the study, an information session at each of the hospitals

was held during a weekly staff meeting so as to include all potential participants and not disrupt care. Participants were issued a consent form, which was signed and returned to the first author. Participants were then given a questionnaire which, once completed, was to be placed in a sealed box the researcher provided in the manager's office and collected within 2 weeks. The manager sent a weekly reminder regarding completion of the questionnaires to participants.

To avoid contamination of data, participants were encouraged not to discuss the answers with each other. Similarly, they were encouraged not to use any other sources to inform their answers, including books or accessing information through electronic devices. Data were collected from April to June 2020. During data collection, COVID-19 protocols were adhered to and an agreement was made with the managers not to disrupt patient care during data collection.

Data processing and statistical analysis

The completeness of the self-administered questionnaire was checked before capturing the data using IBM®SPSS® version 26.0 software package. Descriptive (e.g. mean and standard deviation) and inferential (Chi² test) statistics were generated using IBM® SPSS® version 26.0 software package.3.0 The significance level was determined at $p < 0.05$. The scores for knowledge and practices were calculated as percentages, in the following manner:

Knowledge score = (Number of correct responses to knowledge items 1 to 8)/8 × 100.

An overall practice score could not be determined as there were no true and false responses, which are required to determine overall practice scores.

Rigour

The data collection tool's rigour was enhanced through the pilot study and review by the statistician and basing the questionnaire on existing protocol (Department of Health 2010). Additionally, the questionnaire was reviewed by an expert in radiography to ensure unleading, simple, neutral questions were asked, and participants were assured the questionnaire was anonymous to mitigate performance and response bias.

Think about using anonymous surveys.

Ethical considerations

Permission for the study was provided by the Faculty of Health Sciences at the relevant University (H19-HEA-RAD-008) and by the Provincial Department of Health (EC_201908_002). Permission was granted formally in writing from the CEOs at the different hospitals and verbally from the head of each radiology department. Informed consent was obtained from respondents, and the questionnaire was anonymous.

Results

A total of 85 self-administered questionnaires were completed out of 118 questionnaires issued, resulting in a response rate of 72%. The demographic characteristics of the respondents are presented in Table 1.

As outlined in Table 1, almost half (42.2%) of participants were found to be in the age category of 30–39 years of age and the majority (67.0%) had between 5 and 15 years of work experience. Most participants did not attend a short course (92.9%) nor any pattern recognition training (85.9%).

As the sample size of each of the four hospitals was small and unequal, it was not feasible for hospitals to be compared and thus the results concern all radiographers ($n = 85$) across the four hospitals.

Knowledge and practices

Overall, the best-answered statement in terms of knowledge was B4, regarding which views are done for constipation, with a correct response rate of 100% ($n = 85$). The worst answered statement in the knowledge section was B1, regarding which views are done for diarrhoea, with a correct response rate of 1.2% ($n = 1$) (see Table 2).

In terms of practices, question C1, stating 'I always read the history of the patient before I do the requested examination' received the highest number of correct responses at 94.1% ($n = 80$). Question C6, regarding abdominal views to do when a patient is unable to stand, received the lowest number of correct responses at 18.8% ($n = 16$) (see Table 2).

Demographics and knowledge and practices

Chi-square tests were used to investigate the association between the variables such as age, years of experience,

TABLE 1: Demographic characteristics of the respondents ($N = 85$).

Demographics	Population size (n)	%
Age (in years)		
20–29	29	34.1
30–39	36	42.4
40–49	14	16.5
50–59	6	7.1
Years of experience		
1–4	14	16.5
5–9	31	36.5
10–15	26	30.5
≥ 16	13	15.3
No response	1	1.2
Short course attendance (on performing abdominal examinations)		
No	79	92.9
Yes	4	4.7
No response	2	2.4
Pattern recognition training		
No	73	85.9
Yes	12	14.1

%, percentage; n , number.

TABLE 2: Knowledge and practices of clinical indications of the abdomen.

Variable	Correct		Incorrect	
	<i>n</i>	%	<i>n</i>	%
Knowledge items				
B1. Which view(s) are regarded as an acute abdomen (Answer D: Erect cxr,† erect axr‡ & supine axr)	74	87.1	11	12.9
B2. Which views will be done for renal pathology (Answer B: Erect & supine axr)	6	7.1	79	92.9
B3. Which views will be done for a perforation (Answer D: Erect axr & erect cxr)	72	84.7	13	15.3
B4. Which views will be done for constipation (Answer A: Supine axr)	85	100	0	0
B5. Which views will be done for diarrhoea (Answer B: Erect axr)	1	1.2	84	98.8
B6. Which views will be done for abdominal pain (Answer A: Acute axr series)	43	50.6	42	49.4
B7. Which views do you do for a stab chest (Answer C: Erect cxr)	73	85.9	12	14.1
B8. Which views will demonstrate stab abdomen (Answer D: Erect axr & erect cxr)	50	58.8	32	37.6
Practice statements				
C1. I always read the history of the patient before I do the requested examination (Agree)	80	94.1	4	4.7
C2. Patients must be referred back to the doctor if no history indicated (Agree)	75	88.3	9	10.6
C3. Doctors must be called in when forms are not filled in correctly (Agree)	74	87.1	10	11.8
C4. I perform additional views when pathology is noted (Agree)	77	90.6	6	7.1
C5. I follow incorrect protocols indicated on forms by doctors (Disagree)	76	89.4	8	9.3
C6. Abdominal views to do when a patient is unable to stand (decubitus axr C)	16	18.8	64	75.3
C7. When doctors request incorrect projections, do you (Do what department's protocol says about listed indication & ignore doctors' requests C)	51	60	32	37.4
C8. If a patient is stabbed left side of the abdomen, additional projections you would do (left decubitus A)	51	60	32	37.4
C9. Do you use or follow abdominal protocols afterhours (Yes B)	48	56.5	34	40

Note: Most participants had an average to high knowledge score ($n = 71$; 83.5%), with a mean knowledge score of 59.412, which was in the category of average to high (> 50).

%, percentage; *n*, number.

†, chest x-ray; ‡, abdomen x-ray.

attendance at short courses, pattern recognition training and knowledge and practices (see Table 3, Table 4).

As outlined in Table 3, significant correlations were found between short course attendance and two of the eight knowledge items. These were B3, regarding which views are done for perforation (0.002), and B8, regarding which views demonstrate a stab abdomen (0.002). Similarly, significant correlations were found between short course attendance and eight of the nine practice items (C1, C2, C3, C5, C6, C7, C8 and C9). In addition, significant correlations were found between pattern recognition and one of the eight knowledge items, that is, B1 (Which view(s) are regarded as an acute abdomen) and two practice items: C3 and C5.

Table 4 outlines the overall knowledge score versus demographics items. As outlined, all four demographic items – namely, age (p -value is 0.003), years of experience (p -value is 0.002), short course (p -value is 0.049) and pattern recognition (p -value is 0.029) – were significantly correlated with the overall knowledge score.

Discussion

This study aimed to determine the knowledge and practices of radiographers with regard to the clinical indications for performing radiographic examinations of the abdomen at public hospitals in the Eastern Cape. The study results revealed that although the majority of respondents indicated to have received minimal training in terms of short course attendance and pattern recognition related to abdominal radiographic examinations, overall knowledge of respondents regarding clinical indications for plain abdominal radiographic examinations can be described as above average. Most

participants had an average to high knowledge score ($n = 71$; 83.5%), with a mean knowledge score of 59.412, which was in the category of average to high (> 50). Although average to high knowledge scores is generally regarded a good thing, there is scope for further increasing knowledge levels to enhance practice levels and subsequently service levels and improved patient outcomes (Lundvall, Dahlström & Dahlgren 2021). For example, enhanced knowledge of radiographers regarding clinical indications for plain abdominal radiographic examinations has been found to assist with optimisation of image quality and, subsequently, an accurate diagnosis and appropriate treatment (Alsleem et al. 2019).

The knowledge of respondents regarding clinical indications for performing radiographic examinations of the abdomen varied. The knowledge question regarding which view is done for constipation was answered the best, whereas which view is done for diarrhoea was answered worst, which could be contributed to the fact that the projection required for constipation is easier to obtain, namely, a supine view of the abdomen, which requires less moving and discomfort for a patient, as opposed to the erect radiograph considered as recommended practice for diarrhoea. The latter may therefore not be practised and, although a recommended practice, not remembered well, affecting knowledge scores (Geng et al. 2018). Furthermore, there seems to be a lack of adherence to protocols especially in the out of hours' period, which would infer an inconsistent approach to imaging, and this could lead to health inequalities for patients presenting in the out of hours' period to those during normal working hours. Also, ideally, all participants should have indicated that they always read clinical history as in order to justify the examination valid clinical indications must be included and read prior to exposure as it affects patient safety, as confirmed elsewhere

TABLE 3: Knowledge and practice items versus demographic items ($n = 85$).

Variable	Age		Years of experience		Short course attendance		Pattern recognition	
	Value	df^{\dagger}	Value	df^{\ddagger}	Value	df^{\ddagger}	Value	df^{\ddagger}
Knowledge items B1. Which view(s) are regarded as an acute abdomen	2.443	6	6.694	8	7.792	4	17.520	4
		0.927		0.422		0.201		0.041*
B2. Which views will be done for renal pathology	6.781	6	10.501	8	9.132	4	4.723	4
		0.338		0.230		0.085		0.329
B3. Which views will be done for a perforation	2.368	6	6.600	8	46.468	4	7.242	4
		0.912		0.506		0.002*		0.148
B4. Which views will be done for constipation	13.323	6	5.604	4	0.77	2	6.807	2
		0.071		0.165		1.00		0.141
B5. Which views will be done for diarrhoea	10.808	9	9.445	12	4.132	6	1.145	6
		0.275		0.627		0.437		0.947

Table 3 continues on the next page →

TABLE 3 (Continues...): Knowledge and practice items versus demographic items (n = 85).

Variable	Age		Years of experience		Short course attendance		Pattern recognition	
	Value	df†	Value	df	Value	df	Value	df
B6. Which views will be done for abdominal pain	8.407	9	8.704	12	5.146	6	5.223	6
	α^{\ddagger} 0.504	α 0.687	α^{\ddagger} 0.307	α 0.428	α^{\ddagger} 0.336	α 1.000	α^{\ddagger} 0.831	α 0.365
B7. Which views do you do for a stab chest	13.403	9	12.286	12	6.145	6	1.189	6
	α^{\ddagger} 0.126	α 0.307	α^{\ddagger} 0.900	α 0.002*	α^{\ddagger} 0.009*	α 0.115	α^{\ddagger} 0.121	α 0.115
B8. Which views will demonstrate a stab abdomen	10.612	12	8.711	16	59.021	8	2.203	8
	α^{\ddagger} 0.545	α 0.900	α^{\ddagger} 0.335	α 0.049*	α^{\ddagger} 0.009*	α 0.039*	α^{\ddagger} 0.001*	α 0.039*
Practice statements								
C1. I always read the history of the patient before I do the requested examination	9.634	12	15.395	16	46.671	16	12.82	8
	α^{\ddagger} 0.608	α 0.335	α^{\ddagger} 0.551	α 0.009*	α^{\ddagger} 0.009*	α 0.115	α^{\ddagger} 0.121	α 0.115
C2. Patients must be referred back to the doctor if no history is indicated	9.384	12	13.447	16	47.530	16	12.271	8
	α^{\ddagger} 0.633	α 0.551	α^{\ddagger} 0.340	α 0.001*	α^{\ddagger} 0.001*	α 0.039*	α^{\ddagger} 0.001*	α 0.039*
C3. Doctors must be called in when forms are not filled in correctly	14.114	12	15.991	16	64.983	16	17.615	8
	α^{\ddagger} 0.283	α 0.340	α^{\ddagger} 0.001*	α 0.001*	α^{\ddagger} 0.001*	α 0.039*	α^{\ddagger} 0.001*	α 0.039*

Table 3 continues on the next page →

TABLE 3 (Continues...): Knowledge and practice items versus demographic items ($n = 85$).

Variable	Age		Years of experience		Short course attendance		Pattern recognition	
	Value	df	Value	df	Value	df	Value	df
C4. I perform additional views when pathology is noted	5.306	12	12.470	16	21.383	16	5.681	8
	α 0.961		α 0.585		α 0.108		α 0.387	
C5. I follow incorrect protocols indicated on forms by doctors	12.181	12	18.107	16	43.751	16	95.094	8
	α 0.395		α 0.213		α 0.032*		α 0.001*	
C6. Abdominal views to do when a patient is unable to stand	12.843	9	6.116	12	36.780	12	2.292	6
	α 0.168		α 0.884		α 0.001*		α 0.788	
C7. When doctors request incorrect projections, what would you do?	5.252	9	9.349	12	85.566	12	5.168	6
	α 0.829		α 0.602		α 0.000*		α 0.379	
C8. If a patient is stabbed in the left side of the abdomen, additional views you would do	3.221	9	8.532	12	85.527	12	12.659	6
	α 0.961		α 0.648		α 0.000*		α 0.082	
C9. Do you use/ follow abdominal protocol after hours	9.029	9	9.977	12	56.529	12	4.796	6
	α 0.438		α 0.491		α 0.001*		α 0.309	

†, degree of freedom; ‡, Significance ($p < 0.05$); *, indicates significance.

TABLE 4: Overall knowledge score versus demographic items.

Demographic items	Type III Sum of squares	Degree of freedom (df)	Mean Square	F	Significance
Corrected model	6637.999	19	349.368	2.038	0.018*
Intercept	48523.604	1	48523.604	282.998	0.000*
Age	2608.104	3	869.368	5.070	0.003*
Years of experience	2816.847	3	938.949	5.476	0.002*
Short course	1083.374	2	541.687	3.159	0.049*
Pattern recognition	851.953	1	851.953	4.969	0.029*

*, indicates significance.

(Ebben et al. 2018). Consistent adherences to protocols and recommended practices, crucial to equal and safe patient care can be enhanced through educational strategies, combined with audit and feedback as well as reminders to reinforce protocol adherence (Ebben et al. 2018).

Overall knowledge scores were significantly correlated with age, years of experience, short course attendance and pattern recognition training. Age as well as years of experience have been significantly associated with knowledge among radiographers elsewhere (Stami et al. 2018), and similar to this study could be attributed to the relatively young age of the respondents and medium years of experience (Ugwoke Eze, Uche Eze & Idogwu 2017). Furthermore, the importance of training – in this case attendance at short courses and pattern recognition training – is crucial to develop radiographers' knowledge, professional skills and clinical competency, as recommended elsewhere (Tay & Kaur 2021). Additionally, possible knowledge gaps contributed by age and years of experience can be rectified in short courses and pattern recognition training (Moolman, Mulla & Mdletshe 2020). Attendance at short courses could enhance both knowledge and practices regarding abdominal examinations, as this study found these variables to be significantly correlated with the attendance of short courses, as found in a similar study by Farajollahi et al. (2014).

There is thus a need for sustained education and training regarding clinical indications for abdominal radiographic examinations among radiographers in this study. Additionally, it is recommended that all protocols on abdominal radiographic practices be available to radiographers. Regular in-service training should be provided regarding the protocols, as well as audits and feedback to enhance protocol adherence. Training on the protocols combined with protocol adherence strategies may result in consistently better knowledge and practices regarding clinical indications for plain abdominal radiographic examinations among radiographers, reducing inadequate visualisation of pathology that results in delayed treatment and misdiagnosis, enhancing patient safety and outcomes (Alsleem et al., 2019; Moolman et al. 2020).

This study was limited as the use of convenience sampling, the relatively small sample size and some of the knowledge items related to practices such as using erect AXR for diarrhoea and using acute AXR series for abdominal

pain, which may not be common global practices, affects representativeness, hence limiting the generalisation of the study findings to a larger, global population. There is a need for further exploration of the reasons for the significant relationship between all demographic items and knowledge, which was not investigated in this study. Furthermore, although a pilot study was done to validate the questionnaire, questions and instructions could have been misunderstood by some respondents. Further adjustment and testing of the questionnaire are therefore recommended. Finally, as the study was limited to radiographers in public hospitals in the Eastern Cape only and random sampling because of the small population was not possible, the study should be repeated in radiology departments in public hospitals in South Africa.

Conclusion

The results from this study revealed that although the majority of radiographers in radiology departments in four public hospitals in the Eastern Cape, South Africa, indicated to have received minimal training in terms of short course attendance and pattern recognition related to abdominal radiographic examinations, overall knowledge of respondents regarding clinical indications for plain abdominal radiographic examinations can be described as above average ($n = 71$; 83.5%), with a mean knowledge score of 59.412 (> 50). A significant association between all four demographics (age, years of experience, attended a short course and attended pattern recognition course) and overall knowledge was found. To enhance practices, there is a need for continuous training of radiographers regarding clinical indications for plain abdominal radiographic examinations based on protocols and guidelines, which should be made available to all radiographers as well as audits, feedback and reminders to enhance protocol adherence. Findings of this study could be used to obtain a better understanding of the level of knowledge and practices regarding clinical indications for plain abdominal radiographic examinations among radiographers. Recommendations to expand this study were provided and its findings could be used as the basis for the development of a national policy or strategic plan regarding clinical indications for plain abdominal radiographic examinations in radiology departments.

Acknowledgements

Competing interests

The authors declare that they have no financial or personal relationships that may have inappropriately influenced them in writing this article.

Authors' contributions

L.B., R.W., W.T.H.B. all contributed toward the study conception and design. L.B. was responsible for the data collection and analysis. W.T.H.B. drafted the article and L.B., R.W. were responsible for the critical revision of the article.

Funding information

This research received no specific grant from any funding agency in the public, commercial or not-for-profit sectors.

Data availability

The data that support the findings of this study is available from the corresponding author, W.t.H-B., upon reasonable request.

Disclaimer

The views and opinions expressed in this article are those of the authors and do not necessarily reflect the official policy or position of any affiliated agency of the authors.

References

- Alsleem, H., Davidson, R., Al-Dhafiri, B., Alsleem, R. & Ameer, H., 2019, 'Evaluation of radiographers' knowledge and attitudes of image quality optimisation in paediatric digital radiography in Saudi Arabia and Australia: A survey-based study', *Journal of Medical Radiation Sciences* 66(4), 229–237. <https://doi.org/10.1002/jmrs.366>
- Bell, J.B., 2022, *Abdominal radiography*, viewed 05 May 2022, from <https://radiopaedia.org/articles/abdominal-radiography>.
- Department of Health, 2010, *Routine diagnostic radiography protocols*, Eastern Cape Department of Health, Bhisho.
- Ebben, R.H., Siqeca, F., Madsen, U.R., Vloet, L.C. & Van Achterberg, T., 2018, 'Effectiveness of implementation strategies for the improvement of guideline and protocol adherence in emergency care: A systematic review', *BMJ Open* 8(11), e017572. <https://doi.org/10.1136/bmjopen-2017-017572>
- European Society of Radiology (ESR), 2019, 'Patient safety in medical imaging: A joint paper of the European Society of Radiology (ESR) and the European Federation of Radiographer Societies (EFRS)', *Radiography* 25(2), e26–38. <https://doi.org/10.1186/s13244-019-0721-y>
- Farajollahi, A.R., Fouladi, D.F., Ghojzadeh, M. & Movafaghi, A., 2014, 'Radiographers' professional knowledge regarding parameters and safety issues in plain radiography: A questionnaire survey', *The British Journal of Radiology* 87(1040), 20140090. <https://doi.org/10.1259/bjr.20140090>
- Fox, J.C., 2017, *Clinical emergency radiology*, Cambridge University Press, Cambridge.
- Geng, W.Z., Fuller, M., Osborne, B. & Thoirs, K., 2018, 'The value of the erect abdominal radiograph for the diagnosis of mechanical bowel obstruction and paralytic ileus in adults presenting with acute abdominal pain', *Journal of Medical Radiation Sciences* 65(4), 259–266. <https://doi.org/10.1002/jmrs.299>
- Gleadle, J., Li, J. & Yong, T., 2017, *Clinical investigations at a glance*, John Wiley & Sons, New York.
- Lau, P., 2007, 'Plain radiography', in R. Schmidt & W. Willis (eds.), *Encyclopaedia of pain*, pp. 1836–1841, Springer, Berlin, Heidelberg.
- Lundvall, L.L., Dahlström, N. & Dahlgren, M.A., 2021, 'Radiography students' learning during clinical placements: Developing professional knowing in practice', *Vocations and Learning* 14(3), 439–457. <https://doi.org/10.1007/s12186-021-09269-1>
- McFadden, S., Roding, T., De Vries, G., Benwell, M., Bijwaard, H. & Scheurleer, J., 2018, 'Digital imaging and radiographic practise in diagnostic radiography: An overview of current knowledge and practice in Europe', *Radiography* 24(2), 137–141. <https://doi.org/10.1016/j.radi.2017.11.004>
- Meltzer, A.C., Pines, J.M., Richards, L.M., Mullins, P. & Mazer-Amirshahi, M., 2017, 'US emergency department visits for adults with abdominal and pelvic pain (2007–13): Trends in demographics, resource utilization and medication usage', *The American Journal of Emergency Medicine* 35(12), 1966–1969. <https://doi.org/10.1016/j.ajem.2017.06.019>
- Moolman, N., Mulla, F. & Mdletshe, S., 2020, 'Radiographer knowledge and practice of paediatric radiation dose protocols in digital radiography in Gauteng', *Radiography* 26(2), 117–121. <https://doi.org/10.1016/j.radi.2019.09.006>
- Patel, S.R., Moran, M.E. & Nakada, S.Y., 2018, *The history of technologic advancements in urology*, Springer International Publishing, New York, NY.
- Patterson, J.W., Kashyap, S. & Dominique, E., 2021, *Acute abdomen*, viewed 10 May 2022, from <https://www.ncbi.nlm.nih.gov/books/NBK459328/>.
- Rink, L. & Wessels, I., 2021, *Reference module in biomedical sciences*, Elsevier, Amsterdam.
- Stami, T., Ritin, F. & Dominique, P., 2018, 'Demographic predictors of emotional intelligence among radiation therapists', *Journal of Medical Radiation Sciences* 65(2), 114–122. <https://doi.org/10.1002/jmrs.277>
- Tay, Y.X. & Kaur, B., 2021, 'Competency-based education for training of diagnostic radiographers in percutaneous nephrolithotomy procedure in the operating theatre-an initial experience', *Journal of Medical Imaging and Radiation Sciences* 52(4), S8–S10. <https://doi.org/10.1016/j.jmir.2021.05.010>
- Tomizawa, M., Shinozaki, F., Hasegawa, R., Shirai, Y., Motoyoshi, Y., Sugiyama, T. et al., 2017, 'Abdominal ultrasonography for patients with abdominal pain as a first-line diagnostic imaging modality', *Experimental and Therapeutic Medicine* 13(5), 1932–1936. <https://doi.org/10.3892/etm.2017.4209>
- Ugwoke Eze, C., Uche Eze, C. & Idogwu, H.A., 2017, 'Evaluation of engagement in reflective practice by radiographers in Enugu Metropolis, southeast Nigeria', *South African Radiographer* 55(1), 33–38.
- Velissaris, D., Karanikolas, M., Pantzaris, N., Kipourgos, G., Bampalis, V., Karanikola, K. et al., 2017, 'Acute abdominal pain assessment in the emergency department: The experience of a Greek university hospital', *Journal of Clinical Medicine Research* 9(12), 987. <https://doi.org/10.14740/jocmr3206w>
- Vom, J. & Williams, I., 2017, 'Justification of radiographic examinations: What are the key issues?', *Journal of Medical Radiation Sciences* 64(3), 212–219. <https://doi.org/10.1002/jmrs.211>
- Wit, F., Vroonland, C.C.J.J. & Bijwaard, H., 2022, 'The radiographer's perspective on x-ray examinations in potentially pregnant patients; results of a focus group study among Dutch radiographers', *Radiography* 28(2), 372–377. <https://doi.org/10.1016/j.radi.2021.09.011>