

## EPIDEMIOLOGY OF MALARIA IN CHILDREN LIVING AT IGBO-ORA, SOUTH WESTERN NIGERIA\*

R.I. NWUBA<sup>1</sup>, Y.O. OMOSUN<sup>2,+</sup>, C.I. ANUMUDU<sup>1</sup>, S. ADORO<sup>1,§</sup>,  
A.B. ODAIBO<sup>1</sup>, and M. NWAGWU<sup>1</sup>

1. Cellular Parasitology Programme, Department of Zoology, University of Ibadan, Ibadan, Nigeria.

2. Department of Biological Sciences, Bells University of Technology, Ota, Ogun State, Nigeria.

§ Present address - Immunology Graduate Group, University of Pennsylvania.

(Received: August, 2007; Accepted: May, 2008)

### Abstract

Malaria transmission is seasonal with higher transmission occurring in the rainy season. The burden of malaria falls mainly on children and causes anaemia and fever. Children of school going age are affected and this leads to absence from school. Blood samples were collected from children aged 10 days to 15 years in dry and rainy seasons. Parasite densities were determined by microscopy. Malaria prevalence was higher in the rainy season than in the dry season. In the dry season, 42.4% of the children studied were positive for *P. falciparum*. While at the end of the rainy season 48.4% of the children were malaria positive. The parasite prevalence was not significantly different between males and females. Parasite densities varied from 18 to 52174 parasites per  $\mu$ l of blood. The most abundant group ranged from 1-100 (59%). There was a significant correlation between parasite density and age with the mean parasite density decreasing with age group. The study shows that malaria is more prevalent in the rainy season, and children in rural areas have high prevalence of asymptomatic parasitemia which might lead to symptomatic malaria. The results show that malaria immunity increases with age in both seasons.

**Key words:** Malaria, children, parasitemia, *Plasmodium falciparum*, immunity.

### 1. Introduction

The incidence rate of malaria worldwide is estimated to be 300 to 500 million clinical cases each year, with about 90% of these occurring in Africa, south of the Sahara. Most of these cases are caused by *Plasmodium falciparum* (Meek *et al.*, 2001). It has been stated that most of the children who come to the health centres in Nigeria are treated for malaria (Sule, 2003; Olarewaju and Johnson, 2001; Fawole and Onadoko, 2001, Gellert *et al.*, 1998, Angyo *et al.*, 1996). Also a study by Adediran *et al.*, (2003) showed that 50% of the children admitted to the Obafemi Awolowo teaching hospital children emergency ward that required transfusion had malaria parasitemia. Afolabi, *et al.*, (2001) related the presence of malaria parasites with anaemia in children in the first six months of life, while May *et al.*, (2000) and Lasingu, *et al.*, (2004) showed that the prevalence of anaemia increased with the extent of *P. falciparum* parasitemia. Transplacental transfer of *P. falciparum* to newborn babies was shown by the presence of parasitemia in children within seven days of birth (Egwunyenga *et al.*, 1995). The

presence of parasites in cord blood was related to decrease in weight of new born children (Egwunyenga *et al.*, 1996).

Malarial infection was found to be the most common infection among children of school age admitted for fever in an emergency room in Benin City (Akpede and Sykes, 1993). Studies have shown that school children in rural areas have a much higher prevalence of malaria parasite infection than those in the urban areas (Ademowo *et al.*, 1995; Kazadi *et al.*, 2004). Its being suggested that malaria adversely affects the performance of children at school (Fernando *et al.*, 2003). Children with asymptomatic parasitemia have an increased risk of having symptomatic malaria than children who were negative for malaria parasites (Njama-Meya *et al.*, 2004). The aim of this study was to determine the prevalence of asymptomatic malaria in children aged 10 days to 15 years. This involved looking at the influence of season and age on the prevalence of the parasites in the children.

+ corresponding author (email: yusufomosun@yahoo.com)

\* Presented in part at the First Faculty of Science Conference, Obafemi Awolowo University, Ile-Ife, July 3-5, 2007.

## 2. Materials and Methods

### Study Area:

Igbo Ora and Idere towns in Ibarapa local government area of Oyo state in south-western Nigeria were chosen as the study site. Three rivers, Opeki, Ofiki and Ayin pass through the area. *Anopheles gambiae* and *A. funestus* are the mosquito species found in this area (Lawrence, 1965). Igbo Ora is about 100 km from Ibadan, the state capital; the majority of the population is Yoruba. The climate consists of a warm dry season (November to March) and a cooler rainy season (April to October). The main occupation of the men is farming and hunting while the women are peasant farmers and retail traders (Achidi *et al.*, 1996).

### Study Design:

The cross sectional survey was carried out during the dry season, Jan - Mar 1999 and at the end of the rainy season Oct - Nov 1999. The study protocol was reviewed and approved by the Joint Ethical Committee of the College of Medicine and the University College Hospital, Ibadan. The subjects of the study included infants and children from 10 days to 15 years. 369 and 351 individuals were enrolled in the dry and rainy seasons respectively. Criteria for inclusion into the survey included the age, sex, length of time spent in the study site and informed consent.

### Blood Collection:

Blood (1-2 ml) was collected from the subjects by qualified medical doctors. The blood was stored in sample tubes with 0.12 M trisodium citrate and properly labelled. The blood samples were stored in ice, and transported to the Cellular Parasitology Laboratory in Ibadan. The blood samples were centrifuged at 8000 rpm for two minutes in a biofuge

table top centrifuge. The plasma obtained from it was stored at -80 °C.

### Parasitology:

Thick films were prepared and stained in 10% Giemsa solution in a Wheaton staining jar for 20 minutes and then allowed to dry. The parasites were counted under a microscope using the thick film on the basis of number of parasites per 200 white Blood Cells; this was converted to the number of parasites per  $\mu$ l of blood (WHO, 1985).

### Statistical Analysis:

The results were analyzed using correlation, ANOVA and students t test. The levels of significance were estimated at  $P < 0.05$  for ANOVA and students t test, while  $P < 0.01$  for correlation. The software packages used were Microsoft EXCEL and SPSS.

## 3. Results

In the dry season the most frequent group sampled were children aged 61-120 months and above (163 of 352), followed by 121-180 months (86 of 351), then 0-6 months (42) and 7-12 months (22), and 13-24 months (13), 49-60 months (9), finally 25-48 (8). The age of the individuals sampled ranged from one month to 180 months. 45.4% of the samples were male while 54.6% were female. At the end of the rainy season the most frequent group sampled were children aged 61-120 months and above (126 of 370), followed by 25-48 (76), 121-180 months (71), then 0-6 months (38), 49-60 months (26) and 7-12 months (12), and 13-24 months (8). The age of the samples ranged from one month to 168 months. 50.5% of the samples were male while 49.5% were female. *P. falciparum* parasitemia was prevalent both in the dry season and at the end of the rainy season (Fig. 1). The incidence rate in the two seasons the samples






Fig.1: Incidence rates of Malaria in the dry season and at the end of rainy season. The incidence rates of malaria parasitemia among 351 children, 10 days to 15 years old, during the dry season (January to March 1999) and among 369 children with the same age distribution at the end of the rainy season (October to November 1999), at Idere and Igbo Ora rural towns in south-western Nigeria. The incidence rate was calculated as the percentage of malarial parasite-positive individuals. The actual number of samples screened for parasites in each age group is shown at the top of each bar.



**Fig.2:** Parasite density classes of malaria positive smears. The number of parasite/200 leukocytes was counted and expressed as the number of parasites in a microlitre of blood. The different density classes were identified and the number of subjects in each class was determined and expressed as percentage of the total number of parasitemic subjects. The number of samples screened for parasites in each density class is shown at the top of each bar. The children were all asymptomatic for malaria.



**Figure 3:** Mean parasite density and age. The relationship between age and parasite density was ined. Age is expressed in years while parasite density is expressed as number of parasites in a microlitre of

were collected showed that there were more individuals positive for *P. falciparum* at the end of the rainy season within the different age groups (except 7-12 and 25-48 months age groups) than in the dry season (Fig. 1). In the dry season malaria prevalence rose from 55% to 83% and fell to 28%. At the end of the rainy season the malaria prevalence rose from 71% to 88% and fell to 38%. The trend of the incidence rate shows that malaria in the dry season peaked at 7-12 months and fell as the children got older, while at the end of the rainy season the malaria incidence peaked at 13-24 months and fell as the age of the children increased (Fig. 1). In the dry season, 42.4% of the individuals studied were positive for *P. falciparum*. At the end of the rainy season 48.4% of the children were malaria positive. In the dry season 42.3% (66) of the males and 42.1% (80) of the females were positive for malaria, while at the end of the rainy season 49.7% (92) of the males and 46.7% (85) of the females were positive for malaria. The children who were recruited for the study were all asymptomatic for malaria; there was no record of any one of them being clinically ill. Parasite densities varied from 18 to 52174 parasites per  $\mu\text{l}$  of blood. The most abundant group ranged from 1 - 100 (59%) (Fig. 2). There was a significant correlation ( $P > 0.01$ ,  $r_s = -2.48$ ) between parasite density and age with the mean parasite density decreasing with age group (Fig. 3).

#### 4. Discussion

Epidemiological studies have demonstrated that the burden of morbidity and mortality is concentrated among the youngest age groups under conditions of intense, perennial, stable transmission. The precise relationship between frequencies of parasite exposure, functional immunity and disease risk remains ill defined (Snow *et al.*, 1999). Children living in Igbo - Ora, south western Nigeria, were used in the study. Results from other studies have indicated a high level of exposure in this study area to malaria infection early in life (Achidi *et al.*, 1995). Umar *et al.*, (2002) stated that individuals living in these communities had a malaria morbidity of about 24.4% by looking at hospital data. What we did was to evaluate the number of children with parasitemia, since the children were malaria asymptomatic. The malaria incidence was higher at the end of the rainy season than the dry season (Fig. 1). The results showed that there were more children with parasites in their peripheral blood at the end of the rainy season than in the dry season. This agrees with previous studies carried out by Nwagwu *et al.*, (1998). This was attributed to the increased activity of the disease vector *Anopheles gambiae*, Bradley, (1991) reported that malaria transmission is proportional to mosquito density. Van der Kolk *et al.*, (2003) in a study carried

out in Cameroun showed that only *Anopheles gambiae* and *A. funestus* contributed to malaria transmission in their study area and their distribution was seasonal. The main vectors in Igbo - Ora have been reported to be *Anopheles gambiae* and *A. funestus* (Lawrence, 1965).

Results from this study showed that in the dry season the incidence rates increased from about six months of age 7-12 months old before it began to fall. The incidence rates at the end of the rainy season increased from about 6 months of age to 13-24 months before they began to fall. Children aged 121 - 180 months had the least malaria incidence rate in both seasons (Fig.1). A similar age - dependent trend was observed in a study previously carried out in Igbo - Ora (Achidi *et al.*, 1996), who showed that parasite densities in children in their first year of life increased from two to eight months then started to fall. Oyediran *et al.*, (1993) showed that in Igbo - Ora parasitemia decreased after the one to five years age group. The results of this study showed that children less than one year old were infected, and had higher incidence rates than children two to five years old. This could be due to transplacental transfer of the parasites from mother to child. It could also be suggested that as the children grew older they had more frequent exposure to malaria with their immunity increasing, while those that were younger and had less frequent exposure to the parasite had higher parasitemias due to their lower immunity. Parasite density classes of malaria - positive smears showed that 1-100 (59%) parasites class was most frequent (Fig. 2). This correlated with the previous work by Nwagwu *et al.*, (1998), and it implies that most individuals studied had few parasites in their peripheral blood.

This study showed that parasite density decreased significantly with age (Fig. 3) in agreement with the study carried out by Aribot *et al.*, (1996) and Sharma *et al.*, (2004) who showed that malaria prevalence and parasitemia were inversely correlated with age. Al-yaman *et al.*, 1996, showed that the geometric mean of *P. falciparum* density was highest in children less than two years of age. The results from this study show that there was higher malaria incidence in the rainy season than in the dry season. The parasites prevalence in the children is high and this could affect the children's attendance and performance in school. Children living in rural areas, should be screened routinely for parasites and treatment administered as this could help bring down the rate of absenteeism and increase the performance of the children at school. However this study shows that the ability of the children to resist malaria increases as they grow older.

## ACKNOWLEDGEMENT

We acknowledge the co-operation and endurance of the children who were used in this study. Appreciation also goes to the parents who gave permission for their children to participate in the study. I am grateful to O.O Ososanya, A.A. Adeniran, and A.E. Echoga for their clinical assistance, to E.O. Koyejo for enrolling the children, to A.O Olagoke for her co-operation in the use of the Idere Clinic and Nurses Fayemi and Toriola and the entire field staff at Igbo - Ora for helping with the welfare of the subjects.

This investigation received financial support under the Multilateral Initiative on Malaria (MIM) projects 980050 and A10581 through the UNICEF/UNDP/World Bank/WHO Special Programme for Research and Training in Tropical Diseases (TDR).

## REFERENCES

- Achidi, E.A., Perlmann, H., Salimonu, L.S., Perlmann, P., Walker, O. and Asuzu, M.C.A., 1995. Longitudinal Study of Seroreactivities to *Plasmodium falciparum* Antigens in Nigerian Infants During their First Year Of Life. *Acta Trop.*, 59(2), 173-83.
- Achidi, E.A., Salimonu, L.S., Asuzu, M.C., Berzins, K. and Walker, O., 1996. Studies on *Plasmodium falciparum* parasitemia and Development of Anemia in Nigerian Infants During their First Year of Life. *Am. J. Trop. Med. Hyg.*, 55(2), 138-43.
- Adediran, I.A., Adejuyigbe, E.A. and Oninla, S.O., 2003. Haematological Profile and Malarial Parasitaemia in Nigerian Children Requiring Emergency Blood Transfusion. *Niger. J. Med.*, 12(3), 130-3.
- Ademowo, O.G., Falusi, A.G. and Mewoyeka, O.O., 1995. Prevalence of Asymptomatic Parasitaemia in an Urban and Rural Community in South Western Nigeria. *Cent. Afr. J. Med.*, 41(1), 18-21.
- Afolabi, B.M., Salako, L.A., Mafe, A.G., Ovwigho, U.B., Rabi, K.A., Sanyaolu, N.O. and Ibrahim, M.M., 2001. Malaria in the First 6 Months of Life in Urban African Infants with Anemia. *Am. J. Trop. Med. Hyg.* 65(6), 822-7.
- Akpede, G.O. and Sykes, R.M., 1993. Convulsions with Fever of Acute Onset in School-Age Children in Benin City. *Nigeria. J. Trop. Pediatr.* 39(5), 309-11.
- Al-Yaman, F., Genton, B., Kramer, K.J., Chang, S.P., Hui, G., Baisor, M. and Alpers, M.P., 1996. Assessment of The Role of Naturally Acquired Antibody Levels to *Plasmodium falciparum* Merozoite Surface Protein-1 in Protecting Papua New Guinean Children from Malaria Morbidity. *Am. J. Trop. Med. Hyg.* 54 (5), 443-448.
- Angyo, I.A., Pam, S.D. and Szlachetka, R., 1996. Clinical Pattern and Outcome in Children with Acute Severe Falciparum Malaria at Jos University Teaching Hospital, Nigeria. *East Afr. Med. J.* 73(12), 823-6.
- Aribot, G., Rogier, C., Sarthou, J., Trape, J., Balde, A.T., Druilhe, P. and Roussillon, C., 1996. Pattern of Immunoglobulin Isotype Response to *Plasmodium falciparum* Blood-Stage Antigens in Individuals Living in a Holoendemic Area of Senegal [Dielmo, West Africa]. *Am. J. Trop. Med. Hyg.* 54 (5), 449-457.
- Bradley, D.J., 1991. Malaria. in: R.G Feachem and D.T. Jamison Eds., Disease and Mortality in Sub-Saharan Africa. Oxford University press. International bank for reconstruction and development/The World Bank. pp. 190-202.
- Egwunyenga, O.A., Ajayi, J.A. and Duhlińska-Popova, D.D., 1995. Transplacental Passage of *Plasmodium falciparum* and Serorevaluation of Newborns in Northern Nigeria. *J. Commun. Dis.* 27(2), 77-83.
- Egwunyenga, O.A., Ajayi, J.A., Popova-Duhlińska, D.D. and Nmorsi, O.P., 1996. Malaria Infection of the Cord and Birthweights in Nigerians. *Cent. Afr. J. Med.* 42(9), 265-8.
- Fawole, O.I. and Onadeko, M.O., 2001. Knowledge and Management of Malaria in Under Five Children by Primary Health Care Workers in Ibadan South-East Local Government Area. *Niger Postgrad. Med. J.* 8(1), 1-6
- Fernando, S.D., Gunawardena, D.M., Bandara, M.R., De Silva, D., Carter, R., Mendis, K.N. and Wickremasinghe, A.R., 2003. The Impact of Repeated Malaria Attacks on the School Performance of Children. *Am. J. Trop. Med. Hyg.* 69(6), 582-8.
- Gellert, S., Hassan, B.Y., Meleh, S. and Hiesgen, G., 1998. Malaria Prevalence and Outcome in the In-Patients of the Paediatric Department of the State Specialists Hospital (SSH), Maiduguri, Nigeria. *J. Trop. Pediatr.* 44(2), 109-13.
- Kazadi, W., Sexton, J.D., Bigonsa, M., W'Okanga, B. and Way, M., 2004. Malaria in Primary School Children and Infants in Kinshasa, Democratic Republic Of The Congo: Surveys from the 1980s and 2000. *Am. J. Trop. Med. Hyg.* 71(2 Suppl), 97-102.
- Lawrence, B.R., 1965. Medical Entomology at Igbo-Ora, Western Nigeria. *J. Nig. Med. Ass.* 2, 198-205.
- Lusingu, J.P.A., Vestergaard, L.S., Mmbando, B.P., Drakeley, C.J., Jones, C., Akida, J., Savaeli, Z.X., Kitua, A.Y., Lemnge, M.M. and Theander, T.G., 2004. Malaria Morbidity and Immunity among Residents of Villages with Different *Plasmodium falciparum* Transmission Intensity in North-Eastern Tanzania. *Malar. J.* 3 (1), 26.
- May, J., Falusi, A.G., Mockenhaupt, F.P., Ademowo, O.G., Olumese, P.E., Bienzle, U. and Meyer, C.G., 2000. Impact of Subpatent Multi-Species and Multi-Clonal Plasmodial Infections on Anaemia in Children from Nigeria. *Trans R. Soc. Trop. Med. Hyg.* 94 (4), 399-403.
- Meek, S., Hill, J. and Webster, J., 2001. The Evidence Base for Interventions to Reduce Malaria Mortality in Low and Middle-Income Countries. *CMH Working Paper Series Paper.* WG5, 6
- Njama-Meya, D., Kamya, M.R., Dorsey, G., 2004. Asymptomatic Parasitaemia as a Risk Factor for Symptomatic Malaria in a Cohort of Ugandan Children. *Trop. Med. Int. Health.* 9(8), 862-8.
- Nwagwu, M., Anumudu, C.A., Sodeinde, O., Ologunde, C.A., Obi, T.U., Wirtz, R.A., Gordon, D. M. and Lyon J.A., 1998. Identification of a Subpopulation of Immune Nigeria Adult Volunteers By Antibodies to the Circumsporozoite Protein of *Plasmodium falciparum*. *Am. J. Trop. Med. Hyg.* 58(5), 684-692.
- Olanrewaju, W.I. and Johnson, A.W., 2001. Malaria In Children In Ilorin, Nigeria. *East Afr. Med. J.* 78(3), 131-4.

- Oyediran, A.B.O.O., Topley, E., Osunkoya, B.O., Bamgboye, A., Williams, A.I.O., Ogunba, E.O. and Asuzu M.C., 1993. Severe Morbidity among Children in a Trial Malaria Chemo Prophylaxis with Pyrimethamine or Chloroquine in Ibarapa, Nigeria. *Afr. J. Med. Sci.* 22, 55-63.
- Sharma, S.K., Chattopadhyay, R., Chakrabarti, K., Pati, S.S., Srivastava, V.K., Tyagi, P.K., Mahanty, S., Misra, S.K., Adak, T., Das, B.S. and Chitnis, C.E., 2004. Epidemiology of Malaria Transmission and Development of Natural Immunity in a Malaria-Endemic Village, San Dulakudar, in Orissa State, India. *Am. J. Trop. Med. Hyg.* 71(4), 457-65.
- Snow, R.W., Craig, M., Deichmann, U. and Marsh, K., 1999. Estimating Mortality, Morbidity and Disability due to Malaria among Africa's Non-Pregnant Population. *Bull. W.H.O.* 77(8), 624-639.
- Sule, S.S., 2003. Childhood Morbidity and Treatment Pattern at the Multipurpose Health Centre, Ilesa, Nigeria. *Niger. J. Med.* 12(3), 145-9.
- Umar, U.S., Olumide, E.A. and Briegger W.B., 2002. Morbidity in Rural South-Western Nigeria: A One Year Follow Up of Voluntary Health Worker Consultations in Idere, Oyo State, Nigeria. *Afr. J. Med. Sci.* 31(4), 297-300.
- Van der Kolk, M., Tebo, A.E., Nimpaye, H., Ndongbol, D.N., Sauerwein, R.W. and Eling, W.M., 2003. Transmission of *Plasmodium falciparum* in Urban Yaoundé, Cameroon, is Seasonal and Age-Dependent. *Trans R. Soc. Trop. Med. Hyg.* 97(4), 375-9.
- W.H.O., 1985. Bench Aids for the Diagnosis of Malaria. Plate Numbers 1-8.