

## Ethnicity and gender variability in the diversity, recognition and exploitation of Wild Useful Fungi in Pobè region (Benin, West Africa)

Codjia JEI<sup>1\*</sup> and Yorou NS<sup>2</sup>

1. Ecole Nationale Supérieure des Sciences et Techniques Agronomiques de Kétou (ENSTA/ Ketou), BP 95 Kétou, Benin.

2. Faculté d'Agronomie, Université de Parakou, BP 123, Parakou, Benin.\* Corresponding author: [evans\\_codjia@netcourrier.com](mailto:evans_codjia@netcourrier.com)

Original submitted in on 12<sup>th</sup> February 2014. Published online at [www.m.elewa.org](http://www.m.elewa.org) on 30<sup>th</sup> June 2014. <http://dx.doi.org/10.4314/jab.v78i1.14>

### ABSTRACT

*Objective:* The ultimate goal of this study is to contribute to the enhancement of the livelihood of rural communities through a sustainable exploitation of Non Timber Forest Products, especially the Wild Useful Fungi (WUF).

*Methodology and results:* The study was undertaken in Pobè area located in Southeastern part of Benin (West Africa). Ethnomycological surveys were accomplished within three ethnic groups (Nago, Holli, Fon) from five (05) villages notably Ahoyéyé, Akouho, Igana, Issaba and Towe. We applied proportional sampling techniques to select 84 respondents. The semi-structured and unstructured interviews were conducted within the sampled population, using standardized ethnomycological questionnaires. Classical index such as the Reported Value Use of Gomez Beloz (RVU), Ethnobotanical Use Value (EUV) of a species for a use category, Total Ethnobotanical Use Value (TUV), Diversity Index (DI) and Pielou evenness (EI), as well as the Sorensen index (K) were used to assess the diversity, the level of knowledge, use of wild mushrooms and variability among all three ethnic groups. A total of 19 species including 12 edible ones were recorded, with *Collybia* sp. and *Volaviella volvacea* as the commonly exploited mushrooms in the study area (TUV between 2,14 and 2,40). All recorded species are used in the study area either as food, drugs or source of cash incomes. The study revealed a great variability of knowledge, expertise and of the level of exploitation among ethnic groups ( $K < 50\%$  for all three ethnic groups). However, there exists a certain consistency of local know-how among respondents of the same ethnic group ( $DI = 0,027 < DI \text{ max} = 37/2$ ). At the opposite, the study suggests that mycological know-how was held by a small group of men and a large group of women ( $EI < 0,5 : 0,0007 < 0,5 ; 0,004 < 0,5$ ) suggesting a differential use of fungal resources by rural communities.

*Conclusion and application of results:* This study provides key basic data for a sustainable exploitation of Wild Useful Fungi in Pobè region.

**Keywords:** Diversity, Know-how, uses / use value, value chain, ethnic groups, Wild fungi, Pobè, Benin.

### RÉSUMÉ

*Objectif :* Cette étude dont l'objectif global est de promouvoir la diversité et les usages des champignons sauvages s'est déroulée dans la commune de Pobè située dans le Département du Plateau.

**Méthodologie et résultats :** Cinq (05) villages à raison d'un village par arrondissement de la commune de Pobè ont été choisis pour conduire les enquêtes ethnomycologiques. Il s'agit des villages Ahoyéyé, Akouho, Igana, Issaba et Towé. Un échantillon proportionnel à la taille de chacun des 3 groupes ethniques (Nagot, Holli, Fon) a été appliqué et a permis le choix de 84 personnes sur lesquelles les enquêtes ont été conduites. Les entretiens semi-structurés et non structurés ont été conduits au sein de la population échantillonnée pendant la période d'étude (27 août - 29 octobre). De mêmes, les valeurs d'usages ethnobotaniques, l'indice de diversité et d'équitabilité, de même que l'indice K de Sorensen ont été utilisés pour apprécier la diversité, le niveau de connaissance et d'exploitation des champignons sauvages de même que leur variabilité entre les différents groupes ethniques du milieu. Les différents entretiens ont permis de recenser un total de 19 espèces de champignons dont 12 comestibles et 7 non comestibles. Les différentes espèces inventoriées sont utilisées à des fins alimentaires, commerciales, médicinales et magiques. Du fait de leurs valeurs d'usage plus élevées (comprises entre 2,14 et 2,40) ; les taxa fongiques *Collybia* et *Volvariella volvacea* sont les plus exploitées dans le milieu d'étude. L'étude a révélé une certaine homogénéité de savoir et connaissance ethnomycologique entre les membres d'un même groupe ethnique (valeur d'indice des enquêtés  $ID = 0,027 < ID \text{ max} = 37/2$ ). Par contre, il existe une grande variabilité de savoir, savoir-faire et du niveau d'exploitation d'un groupe ethnique à un autre ( $K < 50 \%$  pour tous les trois groupes ethniques). Par ailleurs, les différents indices d'équitabilité obtenus ( $IE < 0,5 : 0,0007 < 0,5 ; 0,004 < 0,5$ .) suggèrent que la plupart des connaissances sont détenues par un petit groupe au sein des hommes et par un grand nombre de femmes.

**Conclusion et application des résultats :** Cette étude fournit des données nécessaires pour une exploitation durable des champignons sauvages utiles dans la région de Pobè.

**Mots clés :** Diversité, Savoir-faire, usages/Valeur d'usage, variabilité, groupes ethniques, champignons sauvages, Pobè, Bénin.

## INTRODUCTION

Non Timber Forest Products (NTFPs) including inter alia edible wild fruits, nuts, vegetables, beverage, bushmeat, edible and medicinal mushrooms are used by tropical African rural populations for their survival and trade (Malaisse, 2010). Forest Food Resources become a major international concern because not only they display a high diversity, of the important role they play in the livelihood of rural communities but mostly because their exploitation raises up important questions relevant to ecosystem conservation. In Benin, though NTFPs are valuable to the rural communities, they gain very little interest from native scientists and decision makers (van der Zon and Grubben, 1976). Until recently ethnobotanical was been assembled on NTFPs by Assogbadjo *et al.* (2008, 2010) ; Fandohan *et al.* (2010) and Ekue *et al.* (2010). Socio-economic importance of NTFPs for local people in Benin was

addressed by Vodouhè *et al.* (2009). Some NTFPs have a large contribution in African local people's life, where mushrooms constitute a good example. Not only local people of Africa have a large interest in mushroom (Heim, 1977 ; Härkönen *et al.*, 1995 ; Yorou *et al.*, 2014), but local know-how is deep and full of tuition including patterns of edibility, therapeutic medicine, witchcraft (Rammeloo and Walley 1993, Walley and Rammeloo, 1994 ; Yorou and De Kesel, 2002 ; Guissou *et al.*, 2008). The overall objective of this study is to assess the diversity and promote the use of wild mushrooms in the town of Pobè (South Benin). Specifically, this study aims at : Making an inventory of wild useful mushrooms (edible, medicinal) in Pobè area, and assessing the various uses of wild mushrooms by the local people of Pobè.

## MATERIAL AND METHOD

**Study environnement:** The study was undertaken in Pobè region located in South-Eastern part of Benin (Fig 1). Pobè extends on an area of 400 km<sup>2</sup>, making up 11%

of the Plateau Province and 0.46% of Benin flash (Gassi, 2006). The study area is characterized by a bi-modal climate regime with two raining seasons (from April to

July and September to November) which contrast with two dry seasons (from October to March, and July-August). Pobè is the most humid zone of Benin totalizing about 1300 mm rainfall annually and 119 raining days, which largely surpasses the national average (CRAPP,2012). The mean temperature is about 28°C, with March-April being the hottest period. There are two types of soil in the area, including inter alia : ferrallitic, vertisol on sedimentary clay of Lama depression and hydromorphic soil (Dansou, 2011). Vegetation is constituted today by dense forest islands, Guinean

savannahs and plantation of oil palm tree (*Elaeis guineensis* Jacq.). The dense forests are composed of numerous species, but the most common are among others *Terminalia superba* (Engl.) Diels, *Triplochiton scleroxylon* K. Schum., *Dialium guineense* Wild., *Ceiba pentandra* (L.) Gaertn., *Azelia africana* (Sm.) Pers., *Piptadeniastrum africana* (Hook.f.) Brenan, *Cleistopholis patens* (Benth.) Engl. and Diels, *Albizia zygia* (DC.) J.F.Macbr., *Irvingia gabonensis* (Aubry-Lecomte ex O'Rorke) Baill. and mosaics of crops such as : peanut, cotton, maize (Dansou, 2011).




Figure 1 : Map of Pobè Province showing the study area in red. Source : Dansou, 2011.

**Sampling techniques:** Similarly to any region in Benin, the study area hosts a multi-ethnic population among

them the tribes of Holli, Nagot, Fon, Aizo, Bariba, Dendi, Ibo, Yoruba (Gassi, 2006) distributed within 32 villages. A

preliminary prospection was carried out in order to sample representative villages from the 32 villages and define adequate sampling size taking into account the size of each ethnic group. A total of 100 people was randomly sampled within the population in 5 District (Ahoyéyé, Akouho, Igana, Issaba and Towé) regardless the ethnic group. All 100 sampled persons were grouped

into three categories, notably the young (0 - 18 years old), Adults (18 to 35 years old) and Old (over 35 years old). To each of the 100 randomly sampled person, the question «Do you use wild mushrooms ?» was asked. The results of this preliminary surveys are presented in table 1 below.

**Table 1 :** Random sampling of 100 persons and preliminary record of mushrooms recognition and uses.

	Male			Female			Total
	Young (0 to 18 years)	Adults (18 to 35 years)	old (over 35 years)	Young (0 to 18 years)	Adults (18 to 35 years)	old (over 35 years)	
<b>Positive</b>	5	18	25	8	22	16	94
<b>Negative</b>	3	2	-	1	-	-	6
<b>Total</b>	8	20	25	9	22	16	100

As 94% of respondents know and commonly use mushroom, the sampling population size was calculated following sampling strategy of Dagnelie (1988) :

$$N = \frac{U_{1-\alpha/2}^2 \times P \times (1 - P)}{d^2}$$

with : **N** = the size of the respondents to be considered for detailed investigations, **P** = proportion of people knowing and using mushrooms from the preliminary surveys;  $U_{1-\alpha/2}$  is the value of the Normal random variable at

probability value  $1-\alpha / 2$ . For a probability value of 0,975 (or  $\alpha = 0,05$ ),  $U_{1-\alpha/2} \approx 1,96$  ; **d** ( $1-\alpha / 2$ ) margin error of any parameter to be computed from the survey and a value of 5 % was considered.

$$N = \frac{(1.96)^2 \times 0.94 \times 0.06}{(0.05)^2}$$

$$N = 86,66 \approx 87$$

This sampling size was proportionally distributed within all ethnic groups (Table 2).

**Table 2 :** Sampling size of the population within all ethnic groups

Ethnic group	Percentage (%)	Respondents per ethnic group		Male		Female	
<b>Nagot and Holli</b>	84,3	37 Nagot of 73	36 Holli of 73	18 Nagot of 37	18 Holli of 36	19 Nagot of 37	18 Holli of 36
<b>Fons</b>	12,5	11		5		6	
<b>Others</b>	2,8	3		1		2	
<b>Total</b>	100	87					

Representative ethnic groups are composed of Holli, Nagot and Fons. All three ethnic groups were considered for further interviews.

**Specimens sampling:** Forests visits were undertaken in dense forests, fallows and farmlands together with field guides from each ethnic group. Edible fungi were collected directly in the field, and dried after preliminary morphological features were recorded. The protocol of De Kesel *et al.* (2002) was followed for specimens sampling and conservation. For some specimens, professional

pictures were taken directly in the field. The sampled specimens were used to make semi-structured and structured interviews. Whenever needed, a photo guide of edible mushrooms compiled from previous works (De Kesel *et al.*, 2002 ; Yorou *et al.*, 2002) was used to complete the diversity we collected in the field. For

identification of fungi species we used identification books (De Kesel *et al.*, 2002 ; Eyi-Ndong *et al.*, 2011).

**Data analyses:** To evaluate the indigenous knowledge of ethnic groups about fungi species, the use frequencies of each species for each use category were given by each

respondent. With the help of these frequencies, scores were attributed as follow :

- The species is not used at all : 0
- The species is rarely used : 0.5
- The species is frequently used : 1
- The species is very frequently used : 1.5

These scores attributed allow calculating several classical index such as :

➤ 
$$RUV_i = \sum_i^n Species_i$$
 Reported Value Use of Gomez Beloz (RVU):

The total number of use for a species « e » per respondent « i ». However, we determine RUV per ethnic group (and no per respondent) for each species.

➤ Ethnobotanical Use Value (EUV) of a species for a use category: 
$$EUV_k = \frac{\sum_e EUV_{ek}}{N}$$

➤ Total Ethnobotanical Use Value (TUV): 
$$TUV_T = \sum_k EUV_k$$

Diversity Index 
$$DI = \frac{1}{\sum_e P_e^2}$$

**Pe** is the number of use cited per respondent « e » for a species divided by the total of use cited for the species. DI measures how many respondents use the species and how this knowledge is distribute among the respondents.

➤ Pielou evenness (Equitability Index)

It is Diversity index (DI) value divided by the maximal value of Diversity Index (**DI<sub>max</sub>**).

$EI = DI / DI_{max}$ .

This index measures the homogeneity degree of knowledge among respondents. EI varies between 0 and

1. In the practice, this index assesses the homogeneity degree of knowledge among all three ethnic group.

➤ The K Sorensen index was used to assess the diversity, the level of knowledge and use of wild mushrooms and variability among all three ethnic groups. This index is calculated via pair-wise comparison of ethnic groups :

**K = 100 x 2a / (2a + b + c)** with **a** the number of species used by both ethnic groups, **c** is the number of species used by ethnic group 1 and **b** that one used by ethnic group 2.

## RESULTS

**Diversity of Wild Useful Fungi in Pobè region:** A total of 12 edible species and 7 non edible species were collected during the study. The table 3 presents the list of edible species used by each ethnic group along with their

local name and etymologic significance. The following table 4 presents the list of non edible species per ethnic group. The photo 1 presents some edible species recorded in the study area.

**Table 3 :** Edible species, local name and preference level

N°	Scientific names	local name + appreciation			Signification		
		Nagot	Holli	Fon	Nagot	Holli	Fon
1	<i>Volvariella volvacea</i> (Bull.) Sing.	Ohunto èkpè (1.5)	Ohunto èkpè (1)	Dékpohunto (1)	mushroom of palm tree		
2	<i>Collybia</i> sp.	Okiki (1.5)	Okiki (1)	Okiki (1.5)	mushroom that comes out		
3	<i>Marasmius</i> sp.	Idjôdou (1)	Idjôdou (1)	Idjôdou (0.5)	-		
4	<i>Lentinus squarrosulus</i> Mont.	Oluawô (0.5)	Oluawô (0.5)	Ahô (0.5)	-		
5	mushroom sp1	Oludèrè (0.5)	Oludèrè (0.5)	-	-		
6	<i>Termitomyces letestui</i> (Pat.) Heim	OKo-adja (0.5)	-	-	-		
7	<i>Termitomyces schimperi</i> (Pat.) Heim	Oluérin (0.5)	Oluérin (0.5)	Oluérin or Lisso (0.5)	elephant mushroom		
8	<i>Marasmiellus inoderma</i> (Berk.) Singer	Eti-ologbo (0.5)	-	-	-		
9	mushroom sp2	Olubédjé (0.5)	Olubédjé (0.5)	-	-		
10	mushroom sp3	Oluébé (0.5)	-	-	-		
11	<i>Psathyrella tuberculata</i> (Pat.) A. H. Sm.	Oluchichi or ohunto égui (1)	ohunto égui (1.5)	Atinhunto (1)	Mushroom of tree		
12	mushroom sp4	Akantakpa (0.5)	Akantakpa (0.5)	Akantakpa (0.5)	-		

**Table 4 :** List of non edible species

N°	Scientific names	Non edible species		
		Local names		
		Nagot	Holli	Fon
1	mushroom sp5	-	Ohunto Dassèboun	-
2	<i>Leucocoprinus cretatus</i> Locquin ex Lanzoni.	Oluôchôchô	Ohunto édjo	-
3	<i>Gymnopilus cf. pupuratus</i> (Cooke and Masee) Singer	Ohunto édjo	Ohunto édjo	-
4	<i>Ganoderma lucidum</i> (Curtis) Kastern	Ohunto édjo	Ohunto édjo	-
5	<i>Pleurotus</i> sp.	Ohunto édjo	Ohunto édjo	-
6	<i>Gymnopilus</i> sp.	Ohunto édjo	Ohunto édjo	-
7	Mushroom sp6	Ohunto édjo	Ohunto édjo	-






Photo 1: (a – e) : a : *Volvariella volvacea*. b : *Collybia* sp. c : *Psathyrella tuberculata*. d : *Lentinus squarrosulus*. e : *Termitomyces schimperi*.

**Ethnomycological use value** : The uses and use index of each recorded species is presented in table 5 below.

**Table 5** : Use value of recorded species calculated per ethnic group.

	Nagot					Holli					Fon				
	EUV				TUV	EUV				TUV	EUV				TUV
	F	T	M	Ma		F	T	M	Ma		F	T	M	Ma	
<i>Volvariella volvacea</i>	1,41	0,94	0	0	2,36	1,4	1	0	0	2,40	1,31	0,82	0	0	2,14
<i>Collybia sp.</i>	1,43	0,89	0	0,027	2,35	1,23	0,83	0,055	0,028	2,15	1,27	0,72	0	0	2
<i>Marasmius sp.</i>	0,82	0,89	0	0	1,72	0,81	0,86	0	0,014	1,69	0,45	0,14	0	0	0,59
<i>L. squarrosulus</i>	0,25	0,24	0	0	0,5	0,055	0,042	0	0	0,097	0,32	0,09	0	0	0,40
mushroom sp1	0,23	0,20	0	0	0,43	0,18	0,15	0,014	0,014	0,36	0	0	0	0	0
<i>Termitomyces shimperi</i>	0,27	0,23	0	0	0,51	0,18	0,16	0	0	0,35	0,32	0,045	0	0	0,36
mushroom sp2	0,23	0,18	0	0	0,42	0,14	0,083	0	0	0,22	0	0	0	0	0
mushroom sp4	0,094	0,08	0	0	0,17	0,14	0,11	0	0	0,25	0,13	0,09	0	0	0,23
<i>Gerronema inoderma</i>	0,13	0,10	0	0	0,24	0	0	0	0	0	0	0	0	0	0
mushroom sp3	0,13	0,094	0	0	0,23	0	0	0	0	0	0	0	0	0	0
<i>Termitomyces letestui</i>	0,013	0,013	0	0	0,027	0	0	0	0	0	0	0	0	0	0
<i>Psathyrella tuberculata</i>	0,93	0,92	0	0	1,85	0,95	0,90	0,014	0,028	1,88	0,77	0,5	0	0	1,27

**Legend :**

F : Food ; T : Trade ; M : medicine ; Ma : Magico-religious, TUV : Total Ethnobotanical use value, EUV : Ethnobotanical Use Value of a species for a use category

Use categories made of wild fungi in Pobè area include: feeding ; trade ; medicine and magico-religious. The use of fungi varies from one ethnic group to another what is checked by the use value calculated (for instance : EUV Holli : 0.028 ; EUV Nagot : 0 ; EUV Fon : 0). *Volvariella volvacea* and *collybia sp* (TUV between 2,14 and 2,40) are the most used by all three ethnic group.

**Variability of indigenous mycological knowledge between ethnic groups:** The Diversity and Pielou Evenness values were calculated for each species and per ethnic group in order to measure how the knowledge

of the species is distributed within each ethnic group. Table 6 presents results of Diversity Index and Pielou Evenness.



**Table 6 :** Diversity Index and Pielou Evenness for each species within each ethnic group

		<b>Nagot</b>	<b>Holli</b>	<b>Fon</b>
<b><i>Volvariella volvacea</i></b>	DI	0,02	0,02	0,09
	DI max	37	37	11
	EI	0,0007	0,0007	0,008
<b><i>Collybia sp.</i></b>	DI	0,10	0,39	0,09
	DI max	37	31	11
	EI	0,002	0,01	0,008
<b><i>Marasmius sp.</i></b>	DI	0,02	0,06	0,36
	DI max	32	31	5
	EI	0,0009	0,002	0,07
<b><i>Lentinus squarrosulus</i></b>	DI	0,04	0,30	0,28
	DI max	18	4	8
	EI	0,002	0,07	0,03
<b>mushroom sp1</b>	DI	0,06	0,27	0
	DI max	17	13	0
	EI	0,003	0,02	0
<b><i>Termitomyces schimperi</i></b>	DI	0,07	0,08	0,4
	DI max	20	13	7
	EI	0,003	0,006	0,05
<b>mushroom sp2</b>	DI	0,06	0,16	0
	DI max	17	10	0
	EI	0,003	0,01	0
<b>mushroom sp4</b>	DI	0,16	0,11	0,44
	DI max	7	10	3
	EI	0,02	0,01	0,14
<b><i>Marasmiellus inoderma</i></b>	DI	0,111	0	0
	DI max	7	0	0
	EI	0,015	0	0
<b>mushroom sp3</b>	DI	0,12	0,12	0,12
	DI max	10	10	10
	EI	0,01	0,01	0,01
<b><i>Termitomyces letestui</i></b>	DI	0,25	0	0
	DI max	1	0	0
	EI	0,25	0	0
<b><i>Psathyrella tuberculata</i></b>	DI	0,02	0,15	0,10
	DI max	37	36	10
	EI	0,0007	0,004	0,01

Data from table 6 showed that there is a relative homogeneity of knowledge within people of same ethnic groups ( $DI < DI_{max} / 2$ ). The ethnomycological knowledge was more homogenous within Nagot people than in the other ethnic groups since the values of DI for this ethnic group was lower in all cases. The Pielou Evenness measures how the knowledge of a species is distributed between the respondents. EI values calculated were relatively low for all investigated species. In all case,

EI was lower than 0,5. This means that the ethnomycological knowledge is held by a small group of men and a large group of women.

**Variability of edible species between ethnic groups:**

The Sorensen similarity test was conducted between ethnic groups taken in pairs. The values calculated for Sorensen similarity test (K) are summarized in Table 7 below.

Table 7 : Sorensen similarity test between ethnic groups taken in pairs

		Between Holli and Nagot	Between Holli and Fon	Between Nagot and Fon
Values of K (%)		K= 46,1538 %	K= 46,6666 %	K= 42,4242 %
Number of species cited by ethnic group	Holli	12	12	-
	Nagot	9	-	9
	Fon	-	7	7

The value of Sorensen similarity Index (K) is lower than 50 in all cases. It means that there is differences in the use of wild mushrooms by the ethnic groups Nagot, Holli and Fon.

## DISCUSSION

### Diversity and variability of edible mushroom between ethnic groups :

The diversity of edible mushrooms in the study area was high and a total of 12 edible species was recorded. But these species are not fully exploited. In addition to the 12 edible species, 7 non edible species were recorded but still considered by local people as toxic. About fifty macromycete species are consumed by Nagot people of central Benin (Yorou and De Kesel, 2002). For the population Gbaya Bodoé of Centrafrique, Roulon-Doko (1998) had established a list of 60 edible mycotaxa confirming a good mycophagy and knowledge in various areas of this country. Over 60 edible mushroom species have been identified in Tanzania (Buyck *et al.*, 2000; Tibuhwa, 2001; Härkönen *et al.*, 2003). Of all species recorded in the study area, *Volvariella volvacea* is the most common. The study reveals that all three ethnic groups do not use the same fungal resources and the ethnomycologic knowledge varies from one ethnic groups to another ( $K < 50\%$  in all cases). Similar patterns have been reported in Niger (Hama *et al.*, 2012). Indeed, the populations who live at proximity of forest zones are familiar with the mushrooms (Ducousso *et al.*, 2003) and the knowledge accumulated by the populations is diversified, gathering inter alia : edibility, traditional medicine and other uses (De Kesel *et al.*, 2002). It's the

case of Holli in Pobè who hold a large ethnomycologic knowledge because probably they established in the area long time before the other ethnic groups, and thus they are more familiar with the surrounding forests than any other ethnic group.

### Culinary significance of wild edible fungi in Pobè region :

The species *Volvariella volvacea* and *Collybia* sp. are the most exploited species and present same nutritive/food importance as bush meat, fowl and fish (TUV between 2,14 and 2,40). According to respondents, *Volvariella volvacea* present the same taste as meat. This is in accordance with results from the rural communities in Cameroon where mushrooms are considered as meat for the poor (Van Dijk *et al.*, 2003). The importance of mushrooms, and their uses to supplement/substitute bushmeat is common in many West African countries (Yorou *et al.*, 2014). In Burkina Faso, the mushrooms *Phlebopus sudanicus* (Har. and Pat.) Heinem. is designated as the meat for the poor by the Bobo ethnic group (Guissou *et al.*, 2005). *Collybia* sp is also much esteemed because of its sweet taste. On the other side Nagot people from Wari-Marou, prefer larger mushrooms like *Termitomyces* (Yorou and De kesel, 2002). In general, *Termitomyces* species range among the most appreciated species in the whole tropical Africa (Yorou *et*

al., 2014 ; Yorou and De Kesel, 2002 ; Koné *et al.*, 2013). Because of their sporadic and highly spatially variable occurrence that rely upon the presence of intact termites mounds, *Termitomyces* are unfortunately not always available for consumption. In Central and Eastern Africa, chanterelles are the most consumed wild mushrooms (Yorou *et al.*, 2002 ; Eyi-Ndong *et al.*, 2011 ; Buyck, 1994). The population consume regularly the wild edible mushroom in several countries and they constitute an important contribution to improve nutritive value of daily diets, as showed in a study in Malawi (Abbott, 1999). In Jammu and Kashmir state (India), the majority of people questioned showed deep affection for the taste and preparation of wild mushrooms while only a few respondents showed aversion towards them. It was observed that the larger quantities of mushrooms are being consumed in the area and are regarded by many as an equivalent of meat (Kumar and Sharma, 2011). Majority of consumers in Ghana choose edible mushrooms for their taste, which enhance appetite and in the urban centres, mushrooms are included in soups and stews as delicacies (Apetorgbor *et al.*, 2005).

**Medicinal mushroom and use:** The species *collybia* sp., *Psathyrella tuberculata*, mushroom sp1 are used by Nagot and Holli to cure several diseases. They are used in the treatment of epilepsy, bilharzia, hair lice and scabies. The species are used as well to cure a disease called « Itankpa » that is expressed by a loss of skin of feet palm. Though *P. tuberculata* is commonly used in Pobe region for therapeutic purpose, the species is unknown by other ethnic groups of Benin (Nagot, Lokpa, Anii, Kotokoli, Bétamaribè and Peulh) who use this species exclusively as food. Many edible mushrooms have also pharmaceutical property. Like in China, the mushrooms are used abundantly in traditional medicine in tropical Africa and in the world. One hundred and eighty two (182) species of medicinal mushroom are recorded worldwide (Wasser and Weis, 1999). In Occidental Niger, the species such as *Ganoderma colossus* (Fr.) C.F. Baker, *Phellinus allardii* (Bres.) S. Ahmad and *Podaxis pistillaris* (L.) Fr. are used for the health treatment of people (Hama *et al.*, 2012). Similar observations are made in Nigeria where mushrooms are used for the treatment of malnutrition in infants, diabetes, obesity or hyperlipidemia, sterility, anaemia, mumps, fever and protein deficiency (Akpaja *et al.*, 2005; Okhuoya and Akpaja, 2005; Idu *et al.*, 2007). In Ghana, it is reported that *Volvariella volvacea* and *Termitomyces robustus* (Beeli) R. Heim, when eaten served as blood tonic. Of more, *Termitomyces globulus* R. Heim and Gooss.-Font. and *Termitomyces clypeatus* R. Heim, preferred by the

Asantes, Akyems and Akwapims, were also believed to lower blood pressure in hypertensive patients (Apetorgbor *et al.*, 2005).

**Use of wild mushroom on magico-religious plan:** Nagot and Holli hold an important knowledge about magico-religious use of wild mushroom. It is confirmed by ethnobotanical use value calculated (EUV : 0.028, 0.014 etc.). The species who are used here include *Collybia* sp., *Marasmius* sp., « oludère » and *Psathyrella tuberculata*. This species are supposed to bring the luck in the human life. Traders use this species to 'magically' get many customers. Thus, the powder of the cap of these species is used for both cases. The powder is mixed with soap and the trader or human use the mixture during showering. These species are used as love portions. In this case, the powder of any species cited above is mixed with the seed of *Aframomum melegueta* K. Shum. Similar mythological believes are commonly known by the Nigeria people. Here, the Yorubas used selected mushrooms with psychoactive and hallucinogenic effects for idol worshipping and spiritualism (Oso, 1975; Akpaja *et al.*, 2005; Okhuoya and Akpaja, 2005).

**Knowledge and sex influence in the recognition and exploitation of WEF:** The study reveals that of all respondents, the women knew mushrooms better than any other groups. Women search for, and pick mushrooms during field works, though the picking is free and none regulated. The women devote more time on mushroom activities above all food and trade. In most societies in Malawi, picking and trading edible fungi are predominantly a female occupation (Morris, 1987, 1994). Generally, there are no standard guidelines to detect edible species from toxic ones. However, ethnic groups use many local criteria to distinguish edible species from non edible species. In general, edibility criteria are linked to the form, color, fatness, hat. All species who present strange colour (yellow) are considered as toxic. These observations have been reported within Nagot people from central Benin, species with strange deeply colored (blue, red) are regarded as bad and non edible (Yorou and De Kesel, 2002). In Subsaharienne Africa, a change of colour of flesh and of their hymenophore, and thus, the majority of *Boletus* are considered as toxic by the population (Pierce, 1981 ; Thoen *et al.*, 1973). For some population, the change of colour, the disagreeable taste and odour, the tough flesh of mushrooms constitute the proof of toxicity (Eyi-Ndong *et al.*, 2011).

**Ethnicity variability in the use of WEF:** Use of edible mushroom varies between all ethnic groups interviewed. The value of Sorensen similarity Index (K) calculated

show that not all three ethnic groups Nagot, Holli and Fon use the same fungal resources. The Holli are often in contact with mushrooms, because they are true farmer and the hunter. Therefore, they meet often the species of mushrooms in their field and more at the time of the hunter in the forests where many species are encountered. The Holli are permanent forest residents in the study area. At the opposite, the Fon and Nagot are less farmer than Holli folk. Another fact they may explain differential uses of fungal resources is undoubtedly the differences in instalment history of the ethnic groups. More likely the diversity, and the overall uses made of food resources is positively correlated with the instalment background of each folk. In the study area, the Holli are known as the land owners, meaning that their instalment is more ancient than that ones of Fon and Nagot. As such, they are more familiar with the forest, along with forest food resources and the overall forest service they can get is increased. With a more recent instalment history, the Nagot and Fon will need more time to acquaint with the forests. Such differences have been demonstrated by Yorou and De Kesel (2002), Yorou *et al.* (2002a,b,c) in central Benin, where the Nagot people (who are land owners in this area) recognise and exploit more than 30 edible species, whereas Lokpa and Yom people of the same village (with relatively recent establishment) use fewer, but almost different species. Hama *et al.* (2012) reported similar patterns in Niger. Here, the Gourmantché and Peulh as the principal residents of the bush and the forest zones hold more ethnomycologic knowledge than neighbouring ethnic groups Djerma, Touareg and Haoussa. More investigations, addressing establishment history of local folks around forests will help elucidation such differences.

**Composition and origin of local names of wild mushrooms:** In general, local people of Pobè recognize two major groups of fungi : edible and non-edible mushrooms. Edible mushrooms are well known and distinguished. Nevertheless, non edible species are grouped under the same name. The Nagot and Holli call all non- edible mushrooms « ohunto éjo ». The

mushrooms are called Ohunto or Olu in Nagot as well as Holli. They are called « ohunto » in Fon. On the other hand, the Nagot people from central Benin call the mushrooms « Ossoussou » (Yorou and De Kesel, 2002 ). Sometimes, the generic name of mushrooms is completed by one epithet that often recalls of the habitat, morphology, taste or the ecology. As an example, in Nagot and Holli, palm tree mushroom (*Volvariella volvacea*) is called « ohunto èkpè » (èkpè means palm tree). Those of tree are generally called « ohunto égui or oluchichi » (égui means tree). Those produced by the termitary are called « ohunto gbodi or oluérin ». In addition, *Pleurotus* sp. are ranked in the group of non edible mushrooms and are called « Ohunto éjo » (éjo means snake) so all that is related at snake is considered as bad by ethnic groups. It's the case of people in Zambezi Miombo forest who pay no attention to non edible mushroom and hence they have no local name and are systematically considered as poisonous (ukavo or ikoko in Ndendeule and Yao, respectively) (Bloesch and Mbago, 2008). The vernacular names and their etymological significance vary from one ethnic groups to another. It's the case in Niger where the species *Ganoderma colossus* and *Phellinus allardii* are locally called « Maarliali » in Gourmantché and means forbid for pregnant women, « Baggy bodjel » in Fulfudé and means hare tomtom (Hama *et al.*, 2012). On the other side, in Jammu and Kashmir state (India) as many as 31 names of 71 wild mushrooms are recorded and transcribed. This vernacular names could be based on gross morphology or based on varied attributes. For instance, *Peziza vesiculosa* Bull. is recognized as 'Kann Kutch', which means ear like fungus in Kishtwari and Kashmiri. *Boletus* spp. are called 'Dailoo' (the fungus that breaks easily into pieces) in Bhadarwahi or 'Bhuto' and 'Bhutoo' (edible after roasting on fire) in Bhadarwahi and Gaddaishi dialects respectively. *Pleurotus* spp. are named as 'Saroori' (meaning growing on different host plants) in Kishtwari; 'Chur Siner' (growing on *Juglans regia*) in Kishtwari and Kashmiri languages (Kumar and Sharma, 2011).

#### ACKNOWLEDGEMENT

We want to express our sincere thanks to all the population of Pobè for sharing their mycological know-how. Our sincere thanks also go to various guides who help us during this study.

#### REFERENCES

- Abbott P., 1999. Non-timber forest products harvesting: lessons for seasonally sensitive management in miombo. In M.R. Ngulube, L. Mwabumba & P. Chirwa (eds). *Community-based management of Miombo woodlands in Malawi*: 70-89.
- Akpaja E.O., Okhuoya J.A., Eliwer Heferere B.A., 2005. Ethnomycology and indigenous uses of

- mushrooms among the Bini-speaking people of Nigeria: A case study of Aihuobabekun community near Benin City, Nigeria. *International Journal of Medicinal Mushroom* 7(3): 373-374.
- Apetorgbor M.M., Apetorgbor A.K., Nutakor E., 2005. Utilization and cultivation of edible mushrooms for rural livelihood in Southern Ghana. *17th Commonwealth Forestry Conference 2005-Colombo, Sri Lanka*. 22 p.
- Assogbadjo A.E., Glèlè Kakaï R., Adjallala F.H., Akomian F.A., Azihou F., Vodouhè G.F., Kyndt T., Codjia J.T.C., 2010. Ethnic differences in use value and use patterns of the threatened multipurpose scrambling shrub (*Caesalpinia bonduc* L.) in Bénin. *Journal of Medicinal Plants Research* 5(9): 1549-1557.
- Assogbadjo A.E., Glèlè Kakaï R., Chadare F.J., Thomson L., Kyndt T., Sinsin B., Van Damme P., 2008. Folk classification, perception and preferences of baobab products in West Africa: consequences for species conservation and improvement. *Economic Botany* 62: 74-84.
- Bloesch U. and Mbago F., 2008. The potential of wild edible mushrooms in the miombo woodlands of the Selous Niassa Wildlife Corridor for the livelihood improvement of the local population. *Mushroom study Selous Niassa Wildlife Corridor*. 36 p.
- Buyck B., 1994. Ubwoba. Les champignons comestibles de l'Ouest du Burundi. Bruxelles, AGCD. 123 pp.
- Buyck B., Eyssartier G., Kivaisi A., 2000. Addition to the inventory of the genus *Cantharellus* (*Basidiomycota, Cantharellaceae*) in Tanzania. *Nova Hedwigia* 71: 491-502.
- CRAPP, 2012. Rapport final sur les données climatiques. Pobe.
- Dansou S.B., 2011. Pluvial erosion and adaptation of people in township of Pobe. *Master's degree in the category of art, philosophy and sociology*. 75p.
- De Kesel A., Codjia J.T.C., Yorou S.N., 2002. *Guide des champignons comestibles du Bénin*. National Botanic Garden of Belgium, Brussels. 135 p + 35 pl.
- Ducouso M., Ba A.M., Thoen D., 2003. Les champignons ectomycorhiziens des forêts naturelles et des plantations d'Afrique de l'Ouest: une source de champignons comestibles. *Bois et forêts des tropiques* 275(1) : 51-63.
- Ekué M., Sinsin B., Eyog-Matig O., Finkeldey R., 2010. Uses, traditional management, perception of variation and preferences in ackee (*Blighia sapida* K.D. Koenig) fruit traits in Benin: implications for domestication and conservation. *Journal of Ethnobiology and Ethnomedicine* 6 : 12.
- Eyi-Ndong H., Degreef J., De Kesel A., 2011. *Champignons comestibles des forêts denses d'Afrique centrale, Taxonomie et identification*. ABC Taxa Vol 10. 262 p.
- Fandohan B., Assogbadjo A.E., Glèlè Kakaï R., Kyndt T., De Caluwé E., Codjia J.T.C., Sinsin B., 2010. Women's traditional Knowledge, use value, and the contribution of tamarind (*Tamarindus indica* L.) to rural households' cash income in Bénin. *Economic Botany* 64: 248-259.
- Gassi B., 2006. Monographie de la Commune de Pobe. *Programme d'appui au démarrage des Communes*. 46p.
- Gómez-Beloz A., Rucinski J.C., Balick M. J., Tipton C., 2003. Double incision wound healing bioassay using *Hamelia patens* from El Salvador. *Journal of Ethnopharmacology* 88 (2003) : 169 – 173.
- Guissou K.M.L., Sankara P., Guinko S. 2005. *Phlebopus sudanicus* ou "la viande des Bobos", un champignon comestible dans le département de Satin au Burkina Faso. *Crypto. Myco.* 26(3): 195-204.
- Guissou K.M.L., Lykke A.M., Sankara P., Guinko S. 2008. Declining wild mushroom recognition and usage in Burkina Faso. *Econ. Bot.* 62(3): 530-539.
- Hama O., Ibrahim D., Baragé M., Alhou B., Daniëls P.P., Infante F., 2012. Use of some species of Macromycetes in traditional pharmacopoeia in occidental Niger (West Africa). *Journal of Applied Biosciences* 57: 4159– 4167.
- Härkönen M., Saarimäki T., Mwasumbi L. 1995. Edible mushrooms of Tanzania. *Karstenia* 35: 1-92.
- Härkönen M., Niemelä T., Mwasumbi L., 2003. Tanzanian mushrooms. Edible, harmful and other fungi. *Norrinia* 10: 1-200.
- Heim R., 1977. *Termites and mushrooms. The termitophile mushroom of balck Africa and meridional Asia*. Paris, Boubée. 207 p.
- Idu M., Osemwegie O., Timothy O., Onyibe H.I., 2007. A survey of plants used in traditional healthcare by



- Waja tribe Bauchi State, Nigeria. *Plant Archives* 7(2): 535-538.
- Kumar S. and Sharma Y.P., 2011. Diversity of wild mushroom from Jammu and Kashmir (India). *Proceedings of the 7th International Conference on Mushroom Biology and Mushroom Products (ICMBMP7)*. Section: Economical and societal features. 568-577 p.
- Malaisse F., 2010. How to live and survive in Zambezian open Forest (Miombo Ecoregion). *Gembloux, Presses*. 422 pp.
- Morris B., 1987. Common mushrooms of Malawi. Oslo, *Fungiflora*: 108 p.
- Morris B., 1994. Bowa: Ethnomycological notes on the macrofungi of Malawi. In J.H. Seyani & A.C. Chikuni (eds). *Proceedings of the XIIIth Plenary meeting of AETFAT*, Vol. 1: 635-647.
- Okhuoya J.A. and Akpaja E.O., 2005. Mycomedicine and Ethnomycology: The Nigerian Experience. *International Journal of Medicinal Mushrooms* 7(3): 439-440.
- Oso B.A., 1975. Mushrooms and the Yoruba people of Nigeria. *Mycologia* 67(2):311-319.
- Pearce G.D., 1981. *An introduction to Zambia's wild edible mushrooms and how to use them*. Zambia, Forest Dept. 28 pp.
- Rammeloo J. and Walley R., 1993. The edible fungi of Africa south of the Sahara: a literature survey. *Scripta Bot. Belg.* 5: 1-62.
- Roulon-Doko P., 1998. Chasse, cueillette et cultures chez les Gbaya de Centrafrique. *L'Harmattan*. Paris. 540 pp.
- Thoen D., Parent G., lukungu T., 1973. L'usage des champignons dans le Haut-Shaba (République du Zaïre). *Bull. Trim. Centr. Etudes Probl. Soc. Econ.* 100-101: 69-85.
- Tibuhwa D., 2001. *Taxonomy of Termitomyces from Tanzania coastal areas*. MSc. thesis, University of Dar es Salaam.
- van der Zon A.P.M. and Grubben G.J.H., 1976. Les légumes-feuilles spontanés et cultivés du Sud-Dahomey. Communication 65. *Koninklijk Instituut voor de Tropen*. Amsterdam. 111p.
- Van Dijk H., Awana-Onguene N., Kuyper T.W., 2003. Knowledge and utilization of edible mushrooms by local populations of the rainforest of south Cameroon. *Ambio*. 32:19-23.
- Vodouhê F.G., Coulibaly O., Greene C., Sinsin B., 2009. Estimating the Local Value of Non Timber Forest Products to Pendjari Biosphere Reserve Dwellers in Benin. *Economic Botany* 63(4): 397-412.
- Walley R. and Rammeloo J., 1994. The poisonous and useful fungi of Africa south of the Sahara. *Scripta Botanica Belgica* 10: 1-56.
- Wasser S.P. and Weis A.L., 1999. General description of the most important medicinal higher basidiomycetes mushrooms. *International journal of Medicinal Mushrooms* 1: 351-370.
- Yorou S.N. and De Kesel A., 2002. Ethnomycologic knowledge of Nagot in Center of Benin (West Africa). *Proceeding of XVI the AETFAT congress, Brussels 2000. Systematic and Geographic of Plants* 71: 627-637.
- Yorou S.N., De Kesel A, Sinsin B., Codjia J.T.C., 2002a. Diversity and productivity of edible mushroom of Wari-Marô forest. (Benin, West Africa). *Proceedings of XVIth AETFAT Congress, Brussels 2000. Systematic and Geographic of Plants* 71: 613-625.
- Yorou S.N., De Kesel A., Codjia J.T.C., Sinsin B., 2002b. Biodiversity of edible mushroom of Benin. *Proceedings of the Symposium-Workshop on Biodiversity in Benin*. Abomey-Calavi (Benin) October 30<sup>th</sup> to November 18<sup>th</sup> 2002. 231-240 pp.
- Yorou S.N., De Kesel A., Sinsin B., 2002c. Preliminary studies of edible larger fungi in Soudanian Woodlands of Benin. 7<sup>th</sup> congress of the International Mycological Association, Oslo August 2<sup>nd</sup> to 7<sup>th</sup>, 2002. *Abstract book*. 127 p.
- Yorou S.N., Kone A.N., De Kesel A., Guissou M.L., Ekue M.R., 2014. Biodiversity and sustainable use of Wild Edible Fungi in the Soudanian Centre of Endemism : a plea for their valorisation. In Bâ et al (eds). *Ectomycorrhizae in the Tropics and Neotropics*. CRC Press, Science Publisher. London. 241-269 pp.