

## Mycorrhizal status of *Olea europaea* spp. *oleaster* in Morocco

Fadoua Sghir, Mohamed Chliyah, Warda Kachkouch, Mohamed Khouader, Amina Ouazzani Touhami, Rachid Benkirane and Allal Douira

Laboratoire de Botanique et de Protection des Plantes, Université Ibn Tofaïl, Faculté des Sciences, B.P. 133, Kenitra, Maroc.

Corresponding author email: [douiraallal@hotmail.com](mailto:douiraallal@hotmail.com)

Original submitted in on 10<sup>th</sup> November 2012. Published online at [www.m.elewa.org](http://www.m.elewa.org) on 31<sup>st</sup> January 2013.

### ABSTRACT:

**Objective:** This study describes the mycorrhizal oleaster status (*Olea europaea* ssp. *oleaster*.) in the Moroccan ecosystems.

**Methodology and results:** Soil samples were extracted from the rhizosphere of the oleaster tree groves in several regions of Morocco. The frequency and the levels of the arbuscular mycorrhizal fungi (AMF) inside the root bark were measured by assigning an index of mycorrhization from 0 to 5 (Derkowska et al., 2008). The results showed that the AM fungal colonization structures were hyphae, coils and vesicles. The mycorrhizal frequency and intensity reached respectively 70 and 6% in the Bnifougass site. The highest spore density was in the order of 364 g soil spores/100 g and the genus *Glomus* was the dominant one.

The tentative identification test of VAM (Vesicular-Arbuscular Mycorrhizae) species, isolated from the rhizosphere of the oleaster trees, revealed the presence of five fungus species: *Glomus intraradices*, *Glomus clarum*, *Glomus versiforme*, *Acaulospora colossica*, *Scutellospora heterogama*.

**Conclusion:** In all the studied sites the oleaster roots were Mycorrhized. These results open up many opportunities for the application of the controlled mycorrhization in the oleaster plants nurseries production

**Keywords:** Morocco, oleaster, rhizosphere, endomycorrhizae.

### INTRODUCTION

The semi-arid Mediterranean ecosystems are increasingly experiencing critical situations as a result of anthropogenic pressure and different weather conditions like higher temperatures, decreased rainfall and prolonged periods of drought. These pressures have caused the decline of forests, accelerated soil degradation and the developed symbioses transformation, especially mycorrhizal symbioses that can be depleted (Sieverding, 1991; Duponnois et al., 2007). It is well established that the functioning and the stability of terrestrial ecosystems are

mainly dependent on the composition and on the specific vegetation diversity (Hooper and Vitousek, 1997, Tilman et al., 1996). Mycorrhizal fungi play a role in the vegetative cover durability, especially in terrestrial ecosystems of the arid areas (Brundett, 1991, van der Heijden et al., 1998). Mycorrhizal fungi are an essential link between soil and plants. They play an important role in the dynamic of the ecosystem (Bever et al., 1997, van der Heijden et al., 1998) and in the development and the adaptation of vascular plants to their environments (Brundett, 2009), in particular an


adaptation to the drought stress (Strullu, 1991; Varma, 1998). They form a close relationships with plants by providing a particular mineral nutrition needed for their development (van der Heijden et al., 1998) facilitating principally the mobilization and the assimilation of the essential minerals such as phosphorus (Boulard, 1990; Sieverding, 1991, Jakobsen et al., 1992; Smith and Read, 1997) and water (Smith and Read, 1997) by the plants. Symbiosis also plays an important role in the restoration of the degraded soils and in the plants prophylaxis against pathogens (Strullu et al., 1991; Diop, 1996). The wild olive tree (oleaster) is a woody species that occupies a major place in the Moroccan forest ecosystems. This species of the Mediterranean regions occupies the plains and the low mountains in Morocco and does not seem to

demand a lot the type of soil (Benrahmoune and Dubruille, 2003). The occupied areas by this species are very small following the prolonged drought periods that existed in the Mediterranean area and the clearing by indigenous people for the cereal culture installation (Benabid, 2000). These disturbances in the range of the oleaster distribution in Morocco influence probably the microbial community living in close association with the oleaster rhizosphere. The main objective of this work is to know the diversity of the endomycorrhizal fungi naturally present in the oleaster rhizosphere (*Olea oleaster* (Hoofg. and Link.)) growing in some regions of Morocco and to assess the spore diversity of these fungi. In Morocco, the mycorrhizal associations in the oleaster rhizosphere are not known yet.

## MATERIAL AND METHODS

**Presentation of sampling sites:** Soil samples were collected from five sites (Kenitra: Mehdiya, Ouezzane,

Tirghiouine, Tamzazit, Bnifougass) located in three different regions. Fig. 1)



**Fig. 1. Location of the sampling sites.** Sidi Kacem region: Khnicbat (1); Meknès region (2); Taza region: Guercif (3); Sefrou region: Bni fougas (4), Tamzazit (5), Tirghiouine (6); Missour region (7).

The samples were taken in April 2010 from the oleaster rhizosphere (five trees per site at a rate of one kilogram of soil per tree) at a depth of 25 cm and a composite sample of soil was achieved by site. Very fine roots, more likely to be mycorrhized and easily observed under the microscope were taken at the same time with the soil.

**Physico-chemical analyzes of soil:** The main physicochemical soils characteristics (pH, electrical conductivity, organic matter, carbon, nitrate, Nitrogen, ammoniacal nitrogen, mineral nitrogen, phosphore and potassium) were determined by conventional analyzes performed by the soil analysis laboratory of 'Office Régional de Mise en Valeur Agricole du Gharb' (ORMVAG) in Kenitra.

**Roots Extraction and measuring of the roots mycorrhization rate:** The mycorrhization parameters were evaluated by the overall assessment of 30 fragments, as described by Kormanik and McGraw (1982) and Trouvelot et al. (1986) and (Kormanik and McGraw, 1982). The roots were prepared using the method of Koske and Gemma (1989). They were first washed with water and the finest ones were cut into a length of 1 cm and then immersed in a solution of 10% potassium hydroxide (KOH) and placed in a water bath at 90 °C for one hour. The root fragments were then whitened by adding a few drops of hydrogen peroxide H<sub>2</sub>O<sub>2</sub> to the KOH solution. After 5 min, fragments were rinsed with distilled water and stained with a solution of cresyl blue for 15 min at 90 °C in water bath. They were finally rinsed with distilled water and observed under a microscope, each fragment being carefully checked along its entire length, at magnifications of 100 and 400 to record mycorrhizal structures: arbuscules, hyphal walls, vesicles, hyphae intra- and intercellular hyphae extra matrix. The AMF arbuscules and vesicles frequency and levels inside the root bark were measured by assigning an index of mycorrhization from 0 to 5 (Derkowska et al., 2008): 0: no, 1: trace, 2: less than 10%, 3: 11 to 50%, 4: 51 to 90%, 5: more than 91%. The mycorrhizal frequency (F %), reflects the importance of the host plant root system infection by mycorrhizal fungi:

$$F\% = 100 \times (N - N_0) / N$$

With, N: number of the observed fragments and N<sub>0</sub>: number of non-mycorrhizal fragments.

**Mycorrhizal Intensity (M %):** The mycorrhizal Intensity (M %) is defined as the proportion of the root invaded by endomycorrhizal:

$$M\% = (95n_5 + 70n_4 + 30n_3 + 5n_2 + n_1) / N$$

n<sub>5</sub>, n<sub>4</sub>, n<sub>3</sub>, n<sub>2</sub> and n<sub>1</sub> denote the number of fragments scored 5, 4, 3, 2 and 1.

**Content of arbuscules (A %) of the mycorrhized part:**

$$A\% = (100m_3 + 50m_2 + 10m_1) / 100$$

Where MA<sub>3</sub>, MA<sub>2</sub>, MA<sub>1</sub> are the percentages (%) respectively assigned to the notes A<sub>3</sub>, A<sub>2</sub>, A<sub>1</sub>, with, MA<sub>3</sub> = (95 + 70 n<sub>5</sub> n<sub>4</sub> A<sub>3</sub> A<sub>3</sub> + 30 + 5 n<sub>2</sub> n<sub>3</sub> A<sub>3</sub> A<sub>3</sub> n<sub>1</sub> + A<sub>3</sub>) / N. The same for A<sub>1</sub> and A<sub>2</sub>.

In this formula, n<sub>5</sub>A<sub>3</sub> represents the number of fragments marked 5 with A<sub>3</sub>; n<sub>4</sub>A<sub>3</sub> marked the number of fragments 4 with A<sub>3</sub>;

A<sub>0</sub>: no arbuscules, A<sub>1</sub>: some arbuscules 10%, A<sub>2</sub>: moderately abundant arbuscular 50%, A<sub>3</sub>: very abundant arbuscular: 100%.

**Content of vesicles (V %):** It is calculated in the same manner as that of the arbuscular content:

$$V\% = (100 + 50 MV_3 MV_2 MV_1 + 10) / 100$$

Where MV<sub>3</sub>, MV<sub>2</sub>, MV<sub>1</sub> are the percentages (%) respectively assigned notes V<sub>3</sub>, V<sub>2</sub>, V<sub>1</sub>, with, MV<sub>3</sub> = (95 + 70 n<sub>5</sub> V<sub>3</sub> V<sub>3</sub> n<sub>4</sub> + 30 + 5 n<sub>2</sub> n<sub>3</sub> V<sub>3</sub> V<sub>3</sub> n<sub>1</sub> + V<sub>3</sub>) / N. The same for V<sub>1</sub> and V<sub>2</sub>.

In this formula, n<sub>5</sub>V<sub>3</sub> represents the number of fragments marked with 5 with V<sub>3</sub>; n<sub>4</sub>V<sub>3</sub> the number of fragments 4 with V<sub>3</sub>;

V<sub>0</sub>: no vesicles; V<sub>1</sub>: some vesicles 10% V<sub>2</sub>: 50% moderately abundant vesicles; V<sub>3</sub> abundant vesicles: 100%.

**Spores extraction:** Spores were extracted following the wet sieving method described by Gerdemann and Nicolson (1963). In a 1 L beaker, 100g of each composite sample of soil is submerged in 0.5 L of tap water and stirred for 1 minute with a spatula. After 10 to 30 seconds of settling, the supernatant is passed through a sieve of four bunks with decreasing mesh size (500, 200, 80 and 50 microns). This operation was repeated twice. Content retained by the sieves of 200, 80 and 50 microns was divided into two tubes and centrifuged for 5 min at 2000 rev / min. The supernatant was discarded and a viscosity gradient is created by adding 20 ml of sucrose solution at 40% in each centrifuge tube (Walker et al., 1982). The mixture is rapidly stirred and the tube provided in the centrifuge again for 1 min at 3000 rpm / min. Unlike the first centrifuging, the supernatant is poured onto the sieve of 50 μm, the resulting substrate was rinsed with distilled water to remove sucrose and then disinfected with an antibiotic solution (streptomycin). The spores were then recovered with a little distilled water in an Erlenmeyer.

**Species richness and frequency of occurrence of spores:** Species richness is the total number of species observed in every sampling site and the frequency of occurrence of species corresponds to the percentage of sites where each species is detected.

## RESULTS

**Physical and chemical properties of soil:** An analysis of physico-chemical characteristics of soils (Table 1) collected has showed an alkaline pH for almost all the studied sites (pH from 7.97 in Kenitra and above 8 for the other sites). The contents of mineral nitrogen substrates vary from one site to another, they range from 9.8 ppm in the site of Tamzazit and 101.72 ppm in the site of Tirghiouine. The level of available

**Statistical Analysis:** The statistical treatment of results focused on the analysis of variance with a single classification criterion (ANOVA1).

phosphorus and organic matter vary respectively between 3 and 34 ppm and 2.45 and 8.26%. The available potassium content reaches 1028 ppm in the soil of Bnifougass and is of the order of 1000 ppm in the soil of Kenitra. The rate of carbon fluctuates between 1.42% (site of Tamzazit) and 8.29% (site of Tirghiouine).


**Table 1:** Physical and chemical properties of soil samples.

Locality	pH	Total limestone (%)	Electrical Conductivity (mmhos/cm) (1/5)	Organic Matter (%)	Carbon (%)	Nitrate nitrogen (ppm)	Ammoniacal nitrogen (ppm)	Mineral Nitrogen (ppm)	Assimilable phosphore (ppm)	Assimilable potassium (ppm)
Bnifougass	8.04	16.5	0.15	6.3	3.66	28.08	52.08	80.16	5	1028
Ouezzane	8.01	24	0.13	3.62	2.10	64.48	14.76	79.24	6	341
Tamzazit	8.17	14.8	0.15	2.45	1.42	26.64	73.16	9.8	34	940
Tirghiouine	8.01	8.2	0.21	8.29	4.81	36.00	65.72	101.72	21	911
Kénitra	7.97	17.3	0.11	4.31	2.5	62.00	8.64	70.64	3	100


### Mycorrhizal load Characterization of the oleaster:

Microscopic examinations of root fragments treated by the method of Phillips and Hayman (1970) and stained with cresyl blue revealed the presence of different structures characterizing vesicular arbuscular: inter- and intracellular hyphae vesicles of varying shapes and arbuscules. The mycorrhizal frequency of the roots

measured in the different studied sites has been variable, reaching 70% in the site of Bnifougass and 35% in the site of Ouezzane (Fig. 2). The mycorrhization intensity that corresponds to the percentage of mycorrhizal root cortex remains low, around 6%. It's the roots of oleaster of Bnifougass which presented this intensity of mycorrhization (Fig. 3).




**Fig. 2:** Mycorrhizal frequency of the oleaster trees in the studied sites.



**Fig. 3 :** Mycorrhizal intensity of the oleaster trees in the studied sites.

Moreover, the arbuscular contents vary in the studied sites. The highest arbuscular content was recorded in the site of Bnifougass (28%). Meanwhile, the recorded one in the site of Kenitra reached 4% (Fig. 4). The vesicle contents also present variations from one site to another (Fig. 5). They were null in the site of Kenitra.

Meanwhile, they vary between 1% (Ouezzane) and 2% (Tamzazit). Concerning the estimation of the density of spores in the rhizosphere of oleaster growing in the studied sites, the average recorded varies between 364 (Bnifougass) and 168 spores/100g of soil (Ouezzane) (Fig.6).



**Fig. 4 :** Roots arbuscular content of the oleaster trees in the studied sites.




Fig. 5 : Roots vesicular content of the oleaster trees in the studied sites.





Fig. 6 : Spores number of the AM fungal in the oleaster trees rhizosphere in the studied sites .

**Spores diversity of the arbuscular Mycorrhizae (AM):** Preliminary identifications have allowed noting that the spores encountered belong to five species in the order of *Glomales* (Fig. 7): *Scutellospora heterogama*, *Acaulospora colaussica*, *Glomus versiforme*, *G. intraradices* and *G. clarum*. *Glomus clarum* was the most dominant species; its occurrence frequency (Fig. 8) was around 80%. Those of *Glomus*

*sp2* and *Glomus versiforme* reach 33% and 30% respectively. Meanwhile, the occurrence frequencies of *Acaulospora sp1* and *Entrophospora sp.* Species richness (Fig. 9) varied according to the soil sampling sites. It is 7 to 8 species in the site of Bnifougass and Ouazzane and varied between 3 species in the site of Tirghiouine and 6 species in the site of Tamzazit.



**Fig. 7:** Spore of *Glomus versiforme* (A) ; *Acaulospora colossica* (B); *Glomus clarum* (C) , *Glomus intraradices*(D). (G: × 400)




Fig. 8: IS fungi species frequency for eight AM fungi detected in the oleaster trees rhizosphere in the studied sites.




Fig. 9: AM fungi species richness detected in the oleaster trees rhizosphere in the studied sites.

## DISCUSSION AND CONCLUSION

In the all studied sites the oleaster roots were Mycorrhized. Characteristic structures of AM fungi were observed: intra-and extra radical mycelium, vesicles and arbuscules. The oleaster was regarded as a mycotrophic species (Maremmani et al., 2003; Beddiar et al., 2008). The mycorrhizal frequency and intensity reached respectively 70 and 6% in the Bnifougass site, and the average rates of arbuscules and vesicles reached 28 and 2% respectively in the site of Bnifougass and that of Tamzazit. Bouamri et al. (2006)

reported a negative correlation between the mycorrhizal intensity of root cortex and the concentration of available phosphorus in the soil. Indeed, the highest intensity of root mycorrhization of the palm trees was observed at the sites where the soil is deficient on phosphorus, organic matter and nitrogen. For the oleaster, the highest intensity of mycorrhization was recorded in the roots of the oleaster growing in the soil at the site of Bnifougass which are low in phosphorus (5 ppm), but contains mineral nitrogen content in the



order of 25ppm. On the other hand, in the soil of Tamzait site have been shown the highest phosphorus content (34 ppm) and low nitrogen content (9.8 ppm), the intensity of mycorrhizal roots is about 2%. The mycorrhizal roots rate of the oleaster observed in this study appeared low compared with those recorded in Morocco by Abbas *et al.* (2006) and Manaut *et al.*, (2009) respectively *Tetraclinis articulata* (between 27 and 57%) and *Ceratonia siliqua* (40%). According to Diagne and Ingleby (2003), the weakness of the infection may due to the soil poverty on N, P and K. The same authors have highlighted the weakness of the endomycorrhization (<45%) using as substrate for the breeding of plants in disturbed or eroded soil from the north (Tunisia) and the south (Senegal) of the Sahara, compared to weakly disturbed or undisturbed soil (Oasis of Gabes) (up to 100% infection).

The low mycorrhizal infection observed could also be attributed to the method used in estimating the mycorrhization rate. Indeed, the method Gridline would be faster but less accurate than that used in some works, including those of Kormanik and McGraw (1982). Analysis of spore communities CMA found at the rhizosphere of oleaster showed that on average their number was small not exceeding 364 spores / 100 g dry soil. In Morocco, this number is about 170 spores/100 g dry soil at the association *Quercus ilex-Tetraclinis articulata* (Bakkali Yakhlef *et al.*, 2009), from 63 to 98 spores / 100 g soil in coastal dunes of the Souss-Massa (Hatimi and Tahrouch, 2007) and from 2 to 22 spores/100 g soil in the rhizosphere of *Casuarina* sp. (Tellal *et al.*, 2008). The spores density is less than those observed in the rhizosphere of other plants in arid mycotrophic habitats and semi-arid areas such as palm tree (295 to 1900 g of soil spores/100) Tafilalt (Bouamri *et al.*, 2006), the argan trees (900-2080 spores/100 g soil) in southwestern Morocco (Nouaim, 1994). In

general, the fluctuation in the number of MVA spores observed would be allocated to the process of spore formation, their germination and degradation (Smith, 1980), the sampling season (Gemma *et al.*, 1999) and soil and climatic variations (Koske, 1987; Johnson *et al.*, 1991) and microbiological soil (Anderson *et al.*, 1984; Dalpe, 1989).

In the all collected soils, eight species of AM fungi were isolated. Diversification of host plants at trapping could allow the identification of other species. Species isolated belong to three genera: *Glomus*, and *Acaulospora Entrophospora*. The *Glomus* genus is largely dominant. This dominance has been reported by several authors in coastal dunes (Nicolson *et al.*, 1979, Giovannetti *et al.*, 1983, Bergen *et al.*, 1984, Schenck *et al.*, 1980; Ragupathy, 1998; Hatimi and Tahrouch, 2007) and in tetraclinaies (Abbas *et al.*, 2006). It was also found in Burkina Faso by Ba *et al.*, (1996) and in other tropical regions such as Senegal (Diop *et al.*, 1994). The test for identification of some MVA species revealed the presence of five species belonging to two different genera: *Scutellospora heterogama*, *Acaulospora colaussica*, *Glomus versiforme*, *Glomus intraradices* and *Glomus clarum*. Mycorrhizal richness was recorded at all study sites, according to Tacon (1978), AM fungi were not specific and there was no natural site where there was absence of endophytes, but the CMA differ in their infective powers and numbers that appeared more or less important depending on the host and the competition between different endomycorrhizal species.

The diversity of arbuscular mycorrhizal fungi naturally present in the soils of oleaster can be selected and used in reforestation and restoration of degraded ecosystems and even in improving the production of vigorous olive or oleaster plants.

## REFERENCES

- Abbas, Y., Ducouso, M., Abourough, M., Azcon, R. and Duponnois, R., 2006. Diversity of arbuscular mycorrhizal fungi in *Tetraclinis articulata* (Vahl) Masters woodlands in Morocco. *Ann For Sci.*, 63, 285–291.
- Baataoui F.Z., Boumezzough A., Ouahmane L., Hafidi M., & Duponnois R. 2009. Les communautés animales du sol (macro et microarthropodes) et les champignons mycorrhiziens : influence de l'horizon, altitude et saison. Actes du congrès international "Biotechnologie microbienne au service du développement" (MICROBIOD) Marrakech, Maroc, 02-05 Novembre 2009. [www.ucam.ac.ma/microbiona](http://www.ucam.ac.ma/microbiona).
- Bakkali Yakhlef S., Abbas Y., Abourouh M., Hafidi M. & Duponnois R., 2009. Diversité phénotypique et moléculaire des champignons mycorrhiziens de l'association *Quercus ilex-Tetraclinis articulata*. Actes du congrès international "Biotechnologie microbienne au service du développement" (MICROBIOD) Marrakech,

- MAROC 02-05 Novembre 2009.  
[www.ucam.ac.ma/microbiona](http://www.ucam.ac.ma/microbiona)
- Beddiar A., Mekahlia M.N. et Meddad-Hamza A., 2008. Réponse de l'oléastre (*Olea oleaster* (Hoofg. et Link.) et du Sorgho (*Sorghum vulgare* L.) à l'inoculation par quelques morphotypes du genre *Glomus* extraits de sols algériens. Journée francophones de mycorhizes, 4 et 5 septembre 2008. UMR INRA 1088/CNRS 5184/ Université de Bourgogne Plante-Microbe-Environnement BP 86510, 21065 DIJON Cedex France, p. 29. <http://www2.dijon.inra.fr/pme/>
- Benabid A., 2000. Flore et écosystème du Maroc. Evaluation et préservation de la biodiversité. Ibis Press, Paris, p
- Benrahmoune Z. et Dubruille Ch., 2003. Invitation à l'amour des plantes. Guide floristique illustré de la Réserve Biologique de Sidi Boughaba. Edition Acriptura, 319 p.
- Bergen M., Koske R.E., 1984. Vesicular-arbuscular mycorrhizal fungi from sand dunes of Cape Cod, Massachusetts. Transactions of the British Mycological Society, 83: 157-158.
- Bever J.D., Westover K.M., Antonovics J., 1997. Incorporating the soil community into plant population dynamics: the utility of the feedback approach. J. Ecol., 85: 561-763.
- Bouamri B., Dalpé Y., Serrhini M.N. & Bennani A., 2006. Arbuscular Mycorrhizal fungi species associated with rhizosphere of *Phoenix dactylifera* L. in Morocco. African Journal of Biotechnology, 5(6) : 510-516.
- Boullard B., 1990. Guerre et paix dans le règne végétal, Edition Marketing, 181-291p.
- Brundrett, M.C., 1991. Mycorrhizas in natural ecosystems. In: Macfayden A., Begon M., Fitter, A.H. (Eds.), Advances in Ecological Research, vol. 21. Academic Press Ltd., London, pp. 171-213.
- Brundrett, M.C., 2009. Mycorrhizal associations and other means of nutrition of vascular plants: understanding the global diversity of host plants by resolving conflicting information and developing reliable means of diagnosis. Plant and Soil, 320(1-2):37-77.
- Bever J.D., Morton J.B., Antonovics J., Schultz P.A., 1996. Host-dependent sporulation and species diversity of arbuscular mycorrhizal fungi in a mown grassland. J. Ecol., 84:71-82.
- Dalpé Y., 1989. Inventaire et répartition de la flore endomycorhizienne de dunes et de rivages maritimes du Québec du Nouveau Brunswick et de la nouvelle. Ecosse Rev. Ecol. Syst., 116: 219 – 236.
- Derkowska E., Sas Paszt L., Sumorok B., Szwonek E., Gluszek S., 2008. The influence of mycorrhization and organic mulches on mycorrhizal frequency in apple and strawberry roots. J. Fruit Ornament. Plant, 16: 227-242.
- Diagne O. & Ingleby K. 2003. Ecologie des champignons mycorrhiziens arbusculaires infectant *Acacia raddiana*. In : Un arbre au désert. Paris, IRD Editions, pp.205-228.
- Diop T.A., Plenchette C., Strullu D.G., 1994. *In vitro* culture of sheared mycorrhizal roots. Symbiosis, 17:217-227.
- Diop T.A., 1996. Les mycorhizes à vésicules et arbuscules. J. Fac. Sci. Dakar, 2: 49-64.
- Duponnois R., Garbaye J., Bouchard D. et Churin J.L., 1993. The fungus-specificity of mycorrhization helper bacteria (MHBs) used as an alternative to soil fumigation for ectomycorrhizal inoculation of bare-root Douglasfir planting stocks with *Laccaria laccata*. Plants & Soil, 157, 257-262.
- Duponnois R., Plenchette C., Prin Y., Ducosso M., Kisa M., Ba A.M., Galiana A., 2007. Use of mycorrhizal inoculation to improve reforestation process with Australian *Acacia* in Sahelian ecozones. Ecological engineering, 29: 105-112.
- Gemma J.N., Koske R.E. & Carreiro M., 1989. Seasonal variation in spore abundance and dormancy of *Gigaspora gigantea* and in Mycorrhizal inoculum potential of a dune soil. Mycologia, 80 : 211-216.
- Gerdemann J.W., et Nicolson T.H., 1999. Direct interaction between the arbuscular mycorrhizal fungus *Glomus intraradices* and different rhizosphere micro-organisms. New Phytol., 141:525-533.
- Gerdemann, J.W. et Nicolson, T.H., 1963. Spores of mycorrhizal *Endogone* species extracted from soil by wet sieving and decanting. Trans. Br. Mycol. Soc., 46, 235.
- Giovannetti M. et Nicolson T.H., 1983. Vesicular arbuscular mycorrhiza in Italian sand dunes. Trans. Br. Mycol. Soc., 80(3): 552-556.

- Giovannetti M., Sbrana C., Avio L., Citernes A.S., Logi C., 1993. Differential hyphal morphogenesis in arbuscular mycorrhizal fungi during pre-infection stages. *New Phytol.*, 125: 587–593.
- Hatimi et S. Tahrouch. (2007) Caractérisations chimique, botanique et microbiologique du sol des dunes littorales du Souss- Massa.
- Hatim A. & Tahrouch S., 2007. Caractérisations chimique, botanique et microbiologique du sol des dunes littorales du Souss-Massa. *Biomatec Echo.*, 2(5) : 85-97.
- Hooper D.U. et Vitousek P.M., 1997. The effects of plant composition and diversity on ecosystem processes. *Science*, 277: 1302-1305.
- Jakobsen, I., Abbott, L.K. et Robson, A.D., 1992. External hyphae of vesicular-arbuscular mycorrhizal fungi associated with *Trifolium subterraneum*. I. Spread of hyphae and phosphorus inflow into roots. *New Phytol.*, 120: 371–380.
- Jasper D.A., Abbot L.K., Robson D. (1991). The effect of soil disturbance on vesicular-arbuscular mycorrhizal fungi in soils from different vegetation types. *New Phytologist*, 118: 471-476.
- Jasper, D.A., 1994. Management of mycorrhizas in revegetation. In *Management of Mycorrhizas in Agriculture, Horticulture and Forestry* ed. Robson A.D., Abbott L.K. and Malajczuk N., pp. 211–219. Dordrecht: Kluwer.
- Johnson N.C., Zak D.R., Tilman D., Pflieger F.L., 1991. Dynamics of vesicular-arbuscular mycorrhizae during old-field succession. *Oecologia*, 86 : 349-358.
- Kabsy EL H., 2000. Corrélation entre l'incidence du Bayoud, la charge mycorrhizienne et les pratiques culturales dans la palmeraie de Tafilalet. Mémoire de 3<sup>ème</sup> cycle en agronomie. E.N.A.
- Kormanik P.P. and McGraw A.C., 1982. Quantification of Vesicular-arbuscular Mycorrhizae in Plant Roots. In: *Methods and Principles of Mycorrhizal Research*, Sheed, N.C. (Ed.). American Phytopathological Society, St. Paul, pp: 37-45.
- Koske R. E., 1987. Distribution of VA mycorrhizal fungi along a latitudinal temperature gradient. *Mycologia*, 79: 55-68.
- Köske R.E. and Gemma J.N., 1989. A Modified Procedure for Staining Root to Detect VAM. *Mycological Research*, 92 : 486-505.
- Le Tacon F., 1978. La Présence de calcaire dans le sol. Influence sur le comportement de l'Épicéa commun (*Picea excelsa* Link.) et du Pin noir d'Autriche (*Pinus nigra nigricans* Host). *Annales des Sciences forestières*, 35: 165-174.
- Maremmani A., Bedini S., Matosevic I. et Manuela Giovannetti P.E.T., 2003. Type of mycorrhizal associations in two coastal nature reserves of the mediterranean basin. *Mycorrhiza*, 13: 33-40.
- Mukerji K. G., Kapoor A.(1986). Occurrence and Importance of Vesicular- Arbuscular Mycorrhizal Fungi in Semi-arid Regions of India. *Forest Ecology and Management*, 16: 117-126.
- Manaut N., Ouahmane L., Hafidi M. et Duponnois R., 2009. Rôle de la symbiose mycorrhizienne dans l'amélioration de la domestication du caroubier en pépinière. Actes du congrès international "Biotechnologie microbienne au service du développement" (MICROBIOD) Marrakech, MAROC 02-05 Novembre 2009. [www.ucam.ac.ma/microbiona](http://www.ucam.ac.ma/microbiona)
- Nicolson T.H., Johnston C., 1979. Mycorrhiza in Gramineae III. *Glomus fasciculatum* as the endophyte of pioneer grasses in maritime sand dunes. *Trans. Br. Mycol. Soc.*, 72: 261–268.
- Nicolson T.H. et Schenck N.C., 1979. Endogonaceous mycorrhizal endophytes in Florida. *Mycologia*, 71, 1: 178-198.
- Nouaim R., 1994. Ecologie microbienne des sols d'arganeraies: Activités microbiologiques des sols et rôle des endomycorhizes dans la croissance et la nutrition de l'arganier (*Argania spinosa* (L.) Skeels). Thèse de Doctorat, Faculté des Sciences Ibn Zohr, Agadir, 193p + Annexes.
- Philips J.M. et Hayman D.S., 1970. Improved procedures for clearing roots and staining parasitic and vesicular arbuscular mycorrhizal fungi for rapid assessment of infection. *Trans. Brit. Mycol. Soc.*, 55:158-161.
- Ragupathy S.; Nagarjan G.; Mahadevan A., 1998. Mycorrhizae in coastal sand dunes of Tuticorin, Tamil Nadu. *Jour. Envir. Biol.*, 19(3): 281-284.

- Schenck N.C. et Kinloch R.A., 1980. Incidence of mycorrhizal fungi on six field crops in monoculture on a newly cleared woodland site. *Mycologia*, 72: 445-456.
- Schenck N. C., Spain J. L., Sieverding E. & Howeler R. H., 1984. Several new and unreported vesicular-arbuscular mycorrhizal fungi (Endogonaceae) from Colombia. *Mycologia*, 76, 685-699.
- Sieverding, E., 1991. Vesicular–Arbuscular Mycorrhiza Management in Tropical Agrosystems. GTZ, Germany, pp. 79–85.
- Smith T.F., 1980. The effect of season and crop relation with abundance of spores of vesicular arbuscular(VA) mycorrhizal endophytes. *Plant and soil*, 57: 475-479.
- Smith S.E. et Read D.J., 1997. *Mycorrhizal Symbiosis*. 2<sup>nd</sup> edn. UK: Academic Press.
- Strullu D.G., Romand C., Plenchette C., 1991. Axenic culture and encapsulation of the intraradical forms of *Glomus spp.* *World J. Microbiol. Biotechnol.* 7: 292-297.
- Stutz, J.C. et Morton, J.B., 1996. Successive pot cultures reveal high species richness of arbuscular mycorrhizal fungi in arid ecosystems. *Can. J. Bot.* 74 : 1883–1889.
- Strullu D.G., 1991. *Les mycorhizes des arbres et plantes cultivées*. Technique et documentation. Lavoisier, Paris, France.
- Tellal M., 2008. Contribution à l'étude de la symbiose Casuarina-microorganismes et son importance sur la production de plants en pépinière et la fertilité du sol. Thèse de Doctorat. Université Ibn Tofail, Fac. des Sciences, Kénitra, Maroc, 135p.
- Tellal M., Qarro M., Arahou M., Abourouh M., Tellal R., Ghazi L. & Douira A., 2008. Effet de l'Actinomycète *Frankia* sur la croissance et la fixation de l'azote de *Cusuarina glauca* et *C. cunnunghamiana*. *Revue sécheresse*, 19(3): 211-216.
- Tilman D., Wedin D., Knops J., 1996. Productivity and sustainability influenced by biodiversity in grassland ecosystems. *Nature*, 379, 718–720.
- Trouvelot A., Kough J. L. & Gianinazzi V., 1986. Mesure de taux de mycorhization VA d'un système racinaire. Recherche de méthodes d'estimation ayant une signification fonctionnelle. In *physiological and genetic aspects of mycorrhizal*, V. Gianinazzi-Pearson et S. Gianinazzi. (éd.), INRA, Paris, pp 217-221.
- Walker C., 1982. Systematics and taxonomy of arbuscular endomycorrhizal fungi (Glomales) a possible way forward. *Agronomie* 12: 887-897.