

Choice of place for childbirth: prevalence and correlates of utilization of health facilities in Chongwe district, Zambia

*Hazemba AN, Siziya S

Department of Community Medicine, School of Medicine, University of Zambia.

ABSTRACT

Objectives: To determine the prevalence and correlates for utilization of health facilities for childbirth in a rural Chongwe district, Zambia.

Design: A cross sectional study was carried among 250 mothers who had delivered babies within one year prior to the survey. A Backward multiple logistic regression method was used to determine independent predictors for utilization of health services for childbirth.

Main Outcome Measures: Percent utilization of health facilities.

Results: A total of 250 mothers were recruited into the study, of whom the majority were below the age of 25 year (41.2%) and were married (84.0%). A third (32.8%) of the mothers were not able to read and write. The rate of health facility utilisation for childbirth was 42.8%. Independent predictors for utilisation of a health facility were place of last childbirth, and knowledge that traditional birth attendants (TBAs) are given none food items (excluding money) after assisting delivery. Compared to mothers who delivered their last pregnancy at a health facility, mothers who had their last childbirth at home were 85% (AOR=0.15, 95%CI [0.10, 0.22]) less likely to deliver their current pregnancy at a health facility. Knowledge that a TBA was given none food items (excluding money) after assisting delivery was negatively associated with delivery at a health facility (AOR=0.55, 95%CI [0.37, 0.83]).

Conclusion: The importance of giving birth at a health facility should be communicated to mothers who give birth at home, during postnatal visits or clinic outreach sessions.

* Corresponding author:
Alice Ngoma Hazemba
Department of Community Medicine
School of Medicine
University of Zambia
P O Box 50110
Lusaka, Zambia

INTRODUCTION

Zambia adopted the concept of Primary Health Care (PHC) to bring health services including maternity services as close to the family as possible. It is estimated that 50% of households in rural areas are within a radius of 5 km from a health facility¹. In spite of the closeness to a health facility, only 27.9% of live births were delivered at such health facilities², suggesting that 22.1% of the expectant women did not deliver at a health facility despite being within reach of a health facility. There is overwhelming evidence that distance to a health facility is a strong determinant of the choice for maternal health services^{3,4}. However, factors other than distance to health facility have been reported to be associated with health service utilization for childbirth such as education⁵⁻⁸, maternal age^{3,6}, parity^{6,8}, economic status⁷, cultural factors and beliefs, lack of skilled staff at primary health care level, and health worker negative attitudes towards expectant mothers⁹.

Provision of safe motherhood is of utmost importance in the reduction of maternal mortality. Zambia has one of the highest maternal mortality rate in the world at 729/100000 live births², and can be as high as 1300/100000 live births in some remote areas¹⁰. Safe deliveries can only be guaranteed if deliveries are conducted at a health facility by skilled staff. However, staffing of qualified staff for safe deliveries in health facilities in Zambia remains a challenge. Overall, 33% of rural health centres are staffed by unqualified health workers¹¹, and the shortage of midwives is even more critical.

While the ZDHS 2001-2002² provides rates of deliveries conducted at health facilities by background characteristics (maternal age, birth order, residence, province, mother's education, and antenatal care visits), there is scanty information on the factors associated with utilization of health facilities for childbirth. Furthermore, parity, place of delivery of last child, and

Key words: Utilization, health facility, childbirth, rural area, Zambia.

traditional birth attendants-related factors were not examined in this report. We, therefore, conducted this study to determine factors associated with the utilization of health facilities for childbirth in a rural setting of Chongwe district in Zambia. Due to varying rates of utilization of health facilities for child birth between rural areas, we also aimed to estimate this prevalence in a rural district closest to the capital city, Lusaka, of Zambia.

METHODS

Study Design

A cross-sectional study was conducted to collect information on the study factors.

Study site

The description of the study site is given elsewhere¹² but in brief Chongwe had a ratio of 1 midwife to 1,063 women in a population of 145,000 people of whom 31,900 were women in the reproductive age range 15-49 years.

Study population, Sample size, and sampling

Mothers in the reproductive age group of 15-49 years, having delivered six months prior to data collection and attending children's clinic were requested to take part in the study.

Out of the total population of 145,000 (Chongwe District Health Information System report, 2000) people in the district, 4% were children aged less than one year, giving a total of 5800 children. Therefore in a month it was estimated that there would be 483 children under the age of one year. Using a Statcalc programme in Epi Info version 6, with this population size, assuming expected frequency of 50% ± 5% and a 95% confidence level, the required minimum sample size was 217. Furthermore, with a 95% response rate the adjusted minimum sample size was 225.

The first 13 mothers attending children's clinic on the day of data collection from each of the 24 health facilities were selected. However, 4 health facilities were not reached, and in some health facilities only a few mothers attended the children's clinic, thus reducing the sample size.

Data collection techniques

Structured questionnaires were used for interviews. Both closed and open-ended questions were utilised for collecting data. The questionnaire included items on socioeconomic characteristics; obstetric factors, decision making, place delivery of last child, and traditional birth attendants-related factors

Data processing and analysis

Filled-in questionnaires were checked for completeness and consistency of the responses. Open ended questions were post-coded and entered on the questionnaire. Data entry was done using Epi Info version 6. Further, editing of the data occurred after data entry by running frequencies and checking for out of range responses. Preliminary analysis was done in Epi Info version 6 as well. The Yates' corrected Chi-squared test was used to determine associations between predictor variables and the outcome, and the Yates corrected Chi-squared test was not valid, the Fisher's exact test was used. Meanwhile the Backward logistic regression was used to determine independent predictors for utilization of health services for childbirth. Adjusted Odds ratios (AOR) are reported together with their 95% confidence intervals (CI).

Ethical considerations

The protocol was reviewed and approved by the by University of Zambia, Research Ethics Committee. Permission to conduct the study was sought from the Chongwe district Director of Health. All respondents consented before participating in the study.

RESULTS

Sample description

Overall, 250 mothers participated in the survey. The majority, 103 (41.2%) were of age less than 25 years, 89 (35.6%) were aged 25-34 years, and 58 (23.2%) were of age 35 years or more. In terms of the marital status of the participants, 40 (16.0%) were single, and 210 (84.0%) were married. Most, 227 (90.8%), of the participants were house wives, and only 23 (9.2%) were employed. About a third, 80 (32.8%), of the participants were not able to read and write, while 170 (68.0%) were able to do so. The rate of utilization of a health facility for childbirth was 42.8%.

Bivariate analyses

Table 1 shows socio-demographic correlates for choice of place of childbirth. Ability to read and write was positively associated with utilization of health facility for child birth (AOR=2.23, 95%CI [1.23, 4.11]). Parity was also significantly associated with choice of health facility for childbirth (Table 2). Compared with mothers who had 3 or more children, mothers who had one child were 92% (AOR=1.92, 95%CI [1.01, 3.67]) more likely to deliver at a health facility. Mothers who had their last pregnancy delivered at home were 97% (AOR=0.03, 95%CI [0.01, 0.06]) less likely to utilize health facilities for their current childbirth compared to mothers who had delivered their last pregnancy at a health facility. Mothers who thought that there were more women delivered by traditional birth attendants were 42% (AOR=0.58, 95%CI [0.34, 0.99]) less likely to deliver at a health facility compared to mothers who thought otherwise (Table 3).

Table 1. Socio-demographic correlates for choice of place of childbirth

Factor	Health Facility Delivery n (%)	Home delivery n (%)	p value
Age			
<25	51 (47.7)	52 (36.4)	0.199
25-34	34 (31.8)	55 (38.5)	
35+	22 (20.6)	36 (25.2)	
Marital Status			
Single	21 (19.6)	19 (13.3)	0.239
Married	86 (80.4)	124 (86.7)	
Occupation			
House Wife	95 (88.8)	132 (92.3)	0.464
Employed	12 (11.2)	11 (7.7)	
Able to read and write			
Yes	83 (77.6)	87 (60.8)	0.008
No	24 (22.4)	56 (39.2)	

Table 2. Obstetric correlates for choice of place for childbirth

Factor	Health Facility Delivery n (%)	Home delivery n (%)	p value
Number of children			
1	52 (48.6)	46 (32.2)	0.030
2	25 (23.4)	46 (32.2)	
3+	30 (28.0)	51 (35.7)	
Decided on choice of place for delivery			
Self	67 (62.6)	97 (67.8)	0.358
Husband	29 (27.1)	28 (19.6)	
Other	11 (10.3)	18 (12.6)	
Place of delivery of last child			
Home	18 (16.8)	125 (87.4)	<0.001
Health facility	89 (83.2)	18 (12.6)	

Table 3. Traditional birth attendants knowledge-related correlates for choice of place for childbirth

Factor	Health Facility Delivery n (%)	Home delivery n (%)	p value
Thinks more women were delivered by TBAs than in health facility			
Yes	53 (49.5)	90 (62.9)	0.047
No	54 (50.5)	53 (37.1)	
No payment in any form given to TBAs after assisting delivery			
Yes	16 (15.0)	22 (15.4)	0.933
No	91 (85.0)	121 (84.6)	
Money given to TBA after assisting delivery			
Yes	48 (44.9)	79 (55.2)	0.134
No	59 (55.1)	64 (44.8)	
Food given to TBA after assisting delivery			
Yes	42 (39.3)	42(29.4)	0.133
No	101 (70.6)	65(60.7)	
None food items (excluding money) given to TBAs after assisting delivery			
Yes	45 (42.1)	76 (53.1)	0.108
No	62 (57.9)	67 (46.9)	
TBAs offered delivery services			
Yes	100 (93.5)	139 (97.2)	0.213
No	7 (6.5)	4 (2.8)	
TBAs offered PNC			
Yes	12 (11.2)	19 (13.3)	0.766
No	95 (88.8)	124 (86.7)	

Multivariate analysis

Of the factors considered in bivariate analysis, only place of delivery of last child, and giving traditional birth attendants non food items (excluding money) after assisting a delivery were significantly associated with choice of place for childbirth. Compared to mothers who delivered their last pregnancy at a health facility, mothers who had their last childbirth at home were 85% (OR=0.15, 95%CI [0.10, 0.22]) less likely to deliver their current pregnancy at a health facility. Mothers who stated that a traditional birth attendant was given none food items (excluding money) after assisting a delivery were 45% (OR=0.55, 95%CI [0.37, 0.83]) less likely to deliver at a health facility compared to mothers who stated otherwise.

DISCUSSION

We found a rate of utilization of health facility for childbirth in rural Chongwe district in Zambia of 42.8%. This rate is higher than the national figure of 27.9% for rural areas in Zambia. This indicates the large magnitude of variation of health care utilization rate for childbirth between rural districts in Zambia. Compared to other rural areas in the region, our rate is higher than what has been reported in rural Kenya of 5.4%¹³. However, our finding is comparable with that reported in another study in rural Kenya of 47.5%¹⁴, and 37% reported in rural areas of the Orange Free State in South Africa¹⁵.

In the current study, place of last childbirth, and knowledge that traditional birth attendants were given none food items (excluding money) were significantly associated with delivering at a health facility.

We found that mothers who reported that TBAs were given none food items (excluding money) after assisting delivery were 45% less likely to utilize a health facility for childbirth. This finding supports the assertion that it is cheaper for women to be assisted at delivery by TBA than delivery at a health facility in rural areas. However, none food items may be more expensive compared to the cost of delivery at a health facility. In order to convince women in Chongwe district to deliver at a health facility, there is need to compare costs of none food items given to TBAs for delivery assistance and those of delivery at a health facility. A similar study done in rural Kenya¹⁴ Hodgkin found that the average cost of delivery in health facilities was much higher than for TBAs. Almost half of the TBAs in this rural area in Kenya did not charge their clients, and this may or may not be the case in Chongwe. Even if delivery by TBAs

was cheaper in Chongwe, it is important that the quality of delivery services offered by TBAs is evaluated.

The availability of delivery assistance by TBAs has been reported to be associated with non-utilization of a health facility⁴. In communities such as Chongwe that are poorly serviced by health facilities, TBAs will always be used, and there is need to train TBAs in safe delivery, and referral of expectant mothers to health facilities in time to get to the centres. In Chongwe district, Hazemba and Siziya¹² found that 31.4% of TBAs were not trained and recommended that more TBAs be trained.

We found that having had last childbirth at home was negatively associated with current delivery at a health facility. This may have been the case when the last delivery at home was complication-free, and mothers would have had a follow-up delivery at home as well. It is important that these mothers are reached when they seek postnatal care services or during clinic outreach sessions and talked to about the importance of delivering children at a health facility despite that they had a previous safe delivery at home. Having had a safe previous delivery is not a guarantee that the follow-up delivery would also be complication-free. To the best of our knowledge, this is the first time a significant association between place of last childbirth and choice of place of current childbirth is being reported.

Limitations of the study

A number of factors that may confound our findings were not considered in our study such as distance to a health facility, and cultural factors (beliefs and practices). However, we believe these factors may have little influence on our findings that are home-based factors. Our design was a cross sectional study and as with all such studies we can not ascribe causations to the relationships found in the current study, and based on self-report we are not sure whether the respondents may or may not have misreported. However, because the questionnaire was anonymous we have no reason to believe that the respondents may have knowingly misreported in our survey. Our sample was non-random, and as such may not be representative of the target population but may be representative of women attending Under 5 clinics that was our sampled population.

CONCLUSIONS

The importance of giving birth at a health facility should be communicated to mothers who give birth at home. These can be reached when they attend postnatal services or at mobile clinics.

ACKNOWLEDGEMENT

We are grateful to the mothers for their participation in the study. We wish also to express our appreciation to the United States Agency for Development (USAID) for funding the study, and to management for Chongwe District Health Management Team for the permission granted to carry out the study. Lastly but not the least, we are also indebted to Dainess Chinyama for the support she rendered during the period of data collection.

REFERENCES

1. WHO. Health action in crises. <http://www.who.int/hac/crises/zmb/Zambia.aug05.pdf>. Accessed 9 August 2008.
2. Central Statistical Office [Zambia], Central Board of Health [Zambia], ORC Macro. Zambia Demographic and Health Survey 2001-2002. Calverton, Maryland, USA: Central Statistical Office, Central Board of Health, and ORC Macro, 2003.
3. Al-Nahedh NNA. Factors affecting the choice of maternal and child health services in a rural area of Saudi Arabia. *Eastern Mediterranean Health Journal* 1995;261-9.
4. Esimai OA, Ojo OS, Fasubaa OB. Utilization of approved health facilities for delivery in Ile-Ife, Osun State, Nigeria. *Niger J Med* 2002;11:177-9.
5. Osubor KM, Fatusi AO, Chiwuzie JC. Maternal health-seeking behavior and associated factors in a rural Nigerian community. *Matern Child Health J* 2006;159-69.
6. Chowdhury RI, Islam MA, Gulshan J, Chakraborty N. Delivery complications and healthcare-seeking behaviour: the *Bangladesh Demographic Health Survey*, 1999-2000.
7. Tann CJ, Kizza M, Morison L, Mabey D, Muwanga M, Grosskurth H, Elliot AM. Use of antenatal services and delivery care in Entebbe, Uganda: a community survey. *BMC Pregnancy Childbirth* 2007 Oct 11;7:23.
8. al-Nasser AN, Bamgbove EA, Abdullah FA. Providing antenatal services in a primary health care system. *J Community Health* 1994;19:115-23.
9. Kyomuhendo GB. Low use of rural maternity services in Uganda: impact of women's status, traditional beliefs and limited resources. *Reprod Health Matters* 2003;11:16-26.
10. WHO. Health action in crises. <http://www.who.int/hac/crises/zmb/Zambia.aug05.pdf>. Accessed 9 August 2008.
11. Miti SK. The human resources crisis in the health sector in Zambia and efforts by the Ministry of Health to address the crisis. *Med J Zambia* 2007;34:2-4.
12. Hazemba A, Siziya S. Utilisation of maternal care services offered by traditional birth attendants in Chongwe rural district. *Med J Zambia* 2007;34:125-9.
13. Cotter K, Hawken M, Temmerman M. Low use of skilled attendants' delivery services in rural Kenya. *J Health Popul Nutri* 2006;24:467-71.
14. Hodgkin D. Household characteristics affecting where mothers deliver in rural Kenya. *Health Econ* 1996;5:333-40.
15. Cronje HS, Joubert G, Chapman RD, de Winnaar B, Bam RH. Utilisation of maternity services by black women in rural and urban areas of the Orange Free State. *S Afr Med J* 1995;85:1190-1.