

Microbiology of chronic suppurative otitis media at Queen Elizabeth Central Hospital, Blantyre, Malawi: A cross-sectional descriptive study

M Chirwa¹, W Mulwafu², J M Aswani³,
PW Masinde³, R Mkakosya⁴, D Soko⁴

1. ENT Department, Kamuzu Central Hospital, Lilongwe, Malawi

2. College of Medicine, University of Malawi and Queen Elizabeth Central Hospital, Blantyre, Malawi

3. Department of ENT/Head and Neck Surgery, University of Nairobi, Kenya

4. Department of Pathology and Laboratory Sciences, College of Medicine, Blantyre Malawi

Abstract

Background

Chronic suppurative otitis media (CSOM) is still a significant health problem in developing countries. Therefore, it was pertinent to determine the local Malawian microbiology in order to guide adequate treatment, avoid complications, and provide records for future reference.

Aim

The study sought to determine the CSOM-causing microorganisms at Queen Elizabeth Central Hospital in Blantyre, Malawi, and establish their relationship signs and symptoms, and with the demographic pattern of the study.

Methods

This was a hospital-based cross-sectional descriptive study carried out at the ENT outpatient clinic and the Microbiology Department of Queen Elizabeth Central Hospital. The sample comprised 104 patients with unilateral or bilateral active CSOM, who met the inclusion criteria. All patients were evaluated through a detailed history and clinical examination. Pus samples from draining ears were collected by aspiration with a sterile pipette. The specimens were immediately sent for microbiological analysis. Data were analyzed using SPSS version 20.

Results

The study found that *Proteus mirabilis*, *Pseudomonas aeruginosa*, and *Staphylococcus aureus* were the most prevalent aerobic bacteria, while *Bacteroides* spp. and *Peptostreptococcus* spp. were the commonest anaerobic bacteria causing CSOM. These CSOM-causing microorganisms were predominant among males aged 18 years and below. Some CSOM-causing microorganisms were—significantly more so than the others—characteristically associated with each of the following clinical features: quantity of pus drainage, mode of onset, otalgia, hearing loss, location of tympanic membrane perforation, and mucosal appearance.

Introduction

Chronic suppurative otitis media (CSOM) is defined as a perforation of the tympanic membrane, with persistent drainage of pus from the middle ear, lasting at least two weeks¹. The global burden of illness from CSOM is estimated to involve about 65 to 330 million individuals with draining ears, 60% (39 to 200 million) of whom suffer from significant hearing impairment.² Over 90% of the burden is borne by developing countries in Southeast Asia, the Western Pacific Region, and Africa².

Typical pathogens reach the middle ear following insufflations of respiratory pathogens through the eustachian tubes from the nasopharynx and spread from the external ear canal inwards through a non-intact tympanic membrane^{3,4}. Studies on microbiologic diagnoses of CSOM differ in regard to patient age, geography, and the presence of complications such as cholesteatomas, and these inconsistencies likely

impact some of the variation in reported pathogens. A portion of the variability observed may be related to differences in sampling and processing methods^{3,5}. Knowledge of the true frequency of polymicrobial infection, particularly the extent of anaerobic involvement, is limited by differences in collection and culture techniques^{6,7}. Traditional swab specimen collection has been associated with contamination with normal skin flora like *Staphylococcus epidermidis*, diphtheroids and anaerobic organisms, such as *Propionibacterium acnes*⁸.

Aerobes, anaerobes, and fungi are all potential pathogens in CSOM. Understanding of the microbiology of chronic otitis media is important for efficient and effective treatment, and prevention of complications and antibiotic resistance. The objective of this study was to determine the CSOM-causing microorganisms and associated factors among CSOM patients attending the ENT clinic at Queen Elizabeth Central Hospital (QECH) in Blantyre, Malawi.

Methods

This was a hospital-based cross-sectional descriptive study.

This study was carried out in the ENT outpatient clinic at QECH. Queen Elizabeth Central Hospital has bed capacity of 1,200 beds and serves the population of Blantyre District and the Southern Region of Malawi, which has a population of about 5.5 million people. Microbiological analyses were carried out at the QECH microbiology laboratory and University of Malawi College of Medicine's laboratory.

The study was done between July and September 2013. The study population comprised CSOM patients attending the ENT outpatient clinic at QECH during the study period. Inclusion criteria were all patients with actively draining CSOM who consented or whose guardians consented to participate in the study and did not meet any of the exclusion criteria. Exclusion criteria were patients on antibiotic or antifungal treatment (ear drops or systemic) within the previous two weeks, patients with draining ears of less than two weeks duration, patients with draining ears but intact tympanic membrane (otitis externa), and patients who refused to consent to participate in the study.

Sample size determination

The sample size was determined using the Cochran (1963:75) formula to yield a representative sample for proportions. Thus, a sample size of 104 CSOM patients was taken to increase the representativeness of the sample, minimize sampling errors, increase generalizability of the results, and cater for attrition (10%).

Sampling method

The study utilized simple random sampling to select CSOM patients. Subjects were recruited into the study as they came to the clinic until the required number was obtained with strict application of the inclusion and exclusion criteria.

Social demographic and medical history

A detailed clinical history that captured age, gender, duration of discharge and antibiotic therapy was taken for all included

patients. Patients of any age, both genders, unilateral or bilateral draining ears, resulting from CSOM of more than two weeks, were included in the study. Demographic data were taken and, together with the patient's medical history and physical examination findings, were entered in the patient's study file.

Bacterial isolation

Pus specimens from draining ears were taken on the first day of contact with the patient. The ears were inspected first; pus from the outer part of the ear canal was then cleaned by suction. A sterile pipette was then introduced through a sterile aural speculum placed in the external auditory canal, and each specimen was aspirated from the bony part of the ear canal (inner two-thirds) or the middle ear cavity. Pus specimens were collected from both ears for patients with bilateral draining. Each collected specimen was immediately placed in an anaerobic jar, under aseptic conditions, and transported, within one hour of collection, to the microbiology laboratory for routine microbiological culture and identification.

For bacterial isolation, the specimens were inoculated on blood agar, MacConkey's agar, chocolate agar and Robertson's cooked meat medium for aerobic and anaerobic cultures. For anaerobic bacteria, anaerobic blood agars were incubated in an anaerobic jar to permit recovery of anaerobic pathogens. For fungi isolation, a part of the pus specimen was cultured on Sabouraud's dextrose agar. The culture plates were incubated at 37 °C for 24 to 48 hours. Anaerobic culture plates were incubated for seven days to allow growth of anaerobes, which grow slowly compared to aerobes. The isolates from the culture plates were identified using Gram staining, colony morphology, catalase, coagulase, oxidase, and biochemical strips. For fungal growth, lactophenol cotton blue was used for final identification, and culture was for no less than seven days.

Data management

All data collected in the study were sorted, coded, and entered in a computer using SPSS version 20 software. Data were cross-checked against the data files for any inconsistencies or obvious data entry errors. The laboratory request form was also checked for the desired test. The data entry and editing was done throughout the study process. The demographic details, characteristics, and particulars of the subjects, in terms of predictability and determination of risk of CSOM, were analyzed using the Chi-square test. Cross tabulations were done to establish relationships between variables and Chi-square tests were used to test association. Data from bacterial isolation were analyzed using qualitative methods.

Ethical consideration

Ethical approval was obtained from the University of Malawi's College of Medicine Research and Ethics Committee.

Results


There were 64 males (61.5%) and 40 females (38.5%). The mean age was 17.8 years where 64 (61.5%) were aged below 18 years and 40 (38.5%) were aged 18 years and above. The range of the ages was 2 to 64 years, and the median age was 14 years.

Clinical presentation

There were 90 patients (86.5%) with unilateral disease and 14 (6.7%) with bilateral disease. The commonest mode of onset was ear pain in 87 study patients (73.8%), followed by upper

respiratory tract infection (URTI) in 19 (16.1%), trauma in 7 (5.9%), and foreign body in 5 patients (4.2%). Hearing loss was reported in 100 ears (84.7%); it was persistent in 69 ears (68.6%) and fluctuating in 31 (31.4%). Otorrhoea was observed in the left ear in 61 patients (51.7%) and in the right ear in 57 (48.3%). Pus drainage was foul smelling in 90 ears (76.2 %) and odourless in 28 (23.8 %). The pattern of drainage was continuous in 65 ears (55.1%) and recurrent in 53 (44.9%) (Figure 1).

Figure 1: Type and pattern of pus drainage



Examination findings showed that the quantity of pus in the canal was scanty in 63 ears (53.4%) and copious in the other 55 (46.6%). The location of tympanic membrane perforation was central in 54 ears (45.8%), subtotal in 27 (22.9%), marginal in 26 (22.0%), attic in 9 (7.6%), and total perforation in 2 ears (1.7%). Mucosal appearance in the middle ear was injected in 57 (48.4%), hyperplastic in 41 (34.7%), and sclerotic in 20 ears (16.9%). Granulation tissue was present in 20 ears (16.9%).

Laboratory findings

There was an equal number (n = 58; 49.2%) of mixed and pure cultures, while 2 (1.7%) specimens produced no growth. On Gram staining, gram-negative bacteria accounted for 84 specimens (72.4%) and gram-positive bacteria were found in 32 specimens (27.6%). Bacterial findings were categorized into aerobes and anaerobes. The most common aerobes identified from the specimens were gram-negative bacteria, which included *Proteus mirabilis* (n = 44; 28.6%), *Pseudomonas aeruginosa* (n = 32; 20.8%), and *Escherichia coli* (n = 13; 8.4%). The gram-positive aerobes identified included *Staphylococcus aureus* (n = 31; 20.1%) and coagulase-negative staphylococci (n = 6; 4.0%). Other organisms were identified in small numbers (Table 1). Anaerobes were isolated from 39 (33.6%) of the specimens. The most common anaerobes identified were *Bacteroides* spp. in 18 specimens (15.5%), *Peptostreptococcus* spp. in 12 (10.3%), and *Clostridium* spp. in 7 (6.0%) (Table 2). Half of the positive cultures were mixed cultures. Aerobes were identified in the mixed cultures, and mixtures of *P. mirabilis* and *P. aeruginosa* were the most common finding (n = 19; 16.4%). Mixtures of *P. mirabilis* and *E. coli* were found in 8 specimens (6.9%) and *S. aureus* with *P. mirabilis* was found in 7 specimens (6.0%). Mixtures of aerobes and anaerobes were isolated from 34 (29.3%) of the cultures. These comprised mainly *Bacteroides* spp. (n = 15; 12.9%), *Peptostreptococcus* spp. (n = 9; 7.8%) and *Clostridium* spp. (n = 6; 5.2%). Others were observed in small numbers.

Fungal isolates were identified in 21 specimens (18.1%). The fungal isolates consisted of *Aspergillus* spp. (n = 12; 10.3%) and *Candida* spp. (n = 9; 7.8%).

Table 1: Aerobes isolated

Species	Frequency	Percent
Gram-positive aerobes		
<i>Staphylococcus aureus</i>	31	20.1%
Coagulase-negative staphylococci	6	4.0%
<i>Streptococcus pyogenes</i>	3	1.9%
<i>Streptococcus pneumoniae</i>	2	1.3%
Total	42	27.3%
Gram-negative aerobes		
<i>Proteus mirabilis</i>	44	28.6%
<i>Pseudomonas aeruginosa</i>	32	20.8%
<i>Escherichia coli</i>	13	8.4%
<i>Klebsiella pneumoniae</i>	6	4.0%
<i>Proteus vulgaris</i>	6	4.0%
<i>Diphtheroides</i> spp.	4	2.6%
<i>Enterobacter cloacae</i>	3	1.9%
<i>Morganella morganii</i>	1	0.6%
<i>Raultella ornithinolytica</i>	1	0.6%
Coliforms	1	0.6%
<i>Enterobacter cloacae</i>	1	0.6%
Total	112	72.7%

Table 2: Anaerobes isolated

Species	Frequency	Percent
<i>Bacteroides</i> spp.	18	15.6%
<i>Peptostreptococcus</i> spp.	12	10.3%
<i>Clostridium</i> spp.	7	6.0%
<i>Prevotella melaninogenica</i>	2	1.7%
Total	39	33.6%

Table 3: Associations between CSOM-causing microorganisms and clinical features

Clinical feature	Wald chi-square	df	p-value
Otorrhoea	0.046	2	0.831
Type of discharge	1.027	2	0.598
Quantity of discharge	12.208	2	0.002
Type of odour	0.682	1	0.409
Mode of onset	13.565	4	0.009
Presence of otalgia	5.054	1	0.025
Presence of hearing loss	10.628	1	0.001
Character of pus discharged	8.518	4	0.074
Location of TM perforation	38.279	3	0.000
Mucosal appearance	35.445	2	0.000
Granulation tissue	0.478	1	0.489

TM = tympanic membrane

Table 4: Most common organisms associated with specific clinical features of CSOM

Clinical feature	Most common organism
Continuous pus drainage	<i>P. mirabilis</i>
Recurrent pus discharge	<i>S. aureus</i>
Otalgia	<i>P. mirabilis</i>
Persistent hearing loss	<i>S. aureus</i>
Central perforation	<i>P. mirabilis</i>
Marginal perforation	<i>Aspergillus</i> spp.

Association between CSOM-causing microorganisms and symptoms and signs

The chi-square test was used to determine the relationship between the different CSOM-causing microorganisms and symptoms and signs. Some CSOM-causing microorganisms were—significantly more so than the others ($p < 0.05$)—characteristically associated with each of the following clinical features: quantity of pus drainage, mode of onset, otalgia, hearing loss, location of tympanic membrane perforation and mucosal appearance (Table 3, Table 4).

Continuous pus drainage was associated with 20 *P. mirabilis* cultures (17.2%), 11 *P. aeruginosa* cultures (9.5%), 8 *Bacteroides* cultures, and 8 *Aspergillus* cultures (6.9%). *S. aureus* was mostly associated with recurrent discharge (6 specimens; 5.2%). The mode of onset was mainly acute ear pain in specimens with *P. mirabilis* ($n = 16$; 13.7%), *P. aeruginosa* ($n = 15$; 12.8%), and *Aspergillus* spp. ($n = 9$; 7.7%). Otalgia was most commonly associated with *P. mirabilis* ($n = 26$; 22.0%), *P. aeruginosa* ($n = 19$; 16.1%), *S. aureus* ($n = 13$; 11.0%), and *Aspergillus* spp. ($n = 14$; $p = 11.9\%$). Persistent hearing loss was mainly present in ears whose specimens grew *S. aureus* ($n = 16$; 16.0%), *P. aeruginosa* ($n = 13$; 13.0%), *Bacteroides* spp. ($n = 7$; 7.0%), and *Aspergillus* spp. ($n = 7$; 7.0%). The location of tympanic membrane perforation was mainly central in ears whose specimens grew *P. mirabilis* ($n = 15$; 12.7%), *P. aeruginosa* ($n = 9$; 7.6%) and *Aspergillus* spp. ($n = 6$; 5.1%). Subtotal tympanic membrane perforations were mainly associated with specimens that grew *S. aureus* ($n = 5$; 4.2%) and marginal perforations were associated mostly with specimens that grew *Aspergillus* spp. ($n = 12$; 12.0%) and *P. mirabilis* ($n = 7$; 5.9%). The mucosal appearance was mainly injected in ears whose specimens grew *S. aureus* ($n = 6$; 5.2%) and *Candida* spp. ($n = 5$; 4.2%). Hyperplastic mucosal appearance was mostly associated with *P. mirabilis* ($n = 11$; 9.3%), *P. aeruginosa* ($n = 6$; 5.1%), and *Candida* spp. ($n = 3$; 2.5%). Sclerotic mucosa was mostly associated with specimens that grew coliforms, *P. vulgaris* and *Peptostreptococcus* spp. (Table 4).

Discussion

CSOM was most prevalent in children and young adults than in older participants. This is similar to studies reported by others in India and Pakistan^{9,10}. There are several reasons to explain this observation. The eustachian tubes in children are shorter, narrower, and more horizontal than in adults, and frequent upper respiratory tract infections are more common in children¹¹. However, these findings differ from the findings of another study from Singapore, which showed that the disease was more prevalent in the age group of 31 to 40 years¹². In our study, males were more affected than females, and this is similar to findings reported from Ethiopia¹³.

Otorrhoea was observed in the left ear in 61 cases (51.7%), and in the right ear in 57 cases (48.3%). There were 7 bilateral cases. This is comparable with results from a local study and a study from Nigeria¹⁴. Patients in our study, who had bilateral disease, exhibited no differences in terms of CSOM-causing microorganisms between the right and left ears. However, in one study conducted in South Africa, 44.4% of patients with bilateral disease showed differences in the distribution of bacteria, indicating that separate pus specimens need to be taken in bilateral disease¹⁶. The commonest mode

of onset was acute ear pain, which was reported by nearly three-quarters of the patients. Otagia was present in 111 ears (94.1%), which was higher compared to a study in Iraq, where the prevalence of otalgia was 41.7%¹⁷.

The pus drainage was mainly purulent and foul smelling in 90 ears (76.2%) and mucopurulent and odourless in 28 ears (23.8%). This is similar to a study in Bangladesh, where aural drainage was mucoid or mucopurulent in 80 (80%) of patients with tubotympanic CSOM and foul smelling, scanty ear drainage was present in 88 (88%) of patients with atticointral CSOM¹⁸. Findings in a study done in Iraq indicated that pus drainage was foul smelling in 30 ears (25%), which was different from this study. The difference can be attributed to variation in population characteristics and geographical location.

Hearing loss was persistent in 69 ears (68.6%) and fluctuating in 31 ears (31.4%). This is in agreement with a study by Sheahan et al., where 80% of cases had hearing loss¹⁹.

Microbiology

In the present study, gram-negative rods were more than twice as common as gram-positive. It is usually seen that aerobic gram-negative rods outnumber gram-positive cocci in CSOM^{20,21}. This could be attributed to exposure to contaminated water.

The most common bacterial isolates causing CSOM in this study were *P. mirabilis*, *P. aeruginosa*, and *S. aureus*. A study conducted in rural Malawi in 1998 also showed that *P. mirabilis* was the commonest aerobic bacteria²². The findings from the rural setting were similar to those in this study from an urban setting of Malawi in terms of CSOM-causing microorganisms. *P. mirabilis* has been found to be the commonest isolate in rural Kenya,²³ as well as in urban areas of Congo²⁴ and Ethiopia¹³. Findings of *P. mirabilis* as a common isolate suggests that infections are from the community, where there is a constant circulation of *Proteus* and *Staphylococcus* species, as opposed to the health facilities. Although the recommended treatment for *Pseudomonas aeruginosa* is ciprofloxacin ear drops, this is not available in most health facilities in Malawi. As such, treatment with ciprofloxacin ear drops may not explain the relatively lower yield of *P. aeruginosa* ($n = 32$; 20.8%), compared to other studies^{25,26,27}. Various researchers, over the past few decades, have isolated *P. aeruginosa* from 48 to 98% of patients with CSOM. Other studies have shown that *S. aureus* is the most common, especially when cholesteatoma is present^{25,26,27}. The ability of these organisms to form biofilms may contribute to their frequency in CSOM. In this study, *E. coli* was isolated from 13 cultures (8.4%) and *Klebsiella* spp. from 6 (3.9%); these are comparable to findings reported elsewhere.^{22,10} More frequent isolation of faecal bacteria, like *E. coli* and *Klebsiella* spp., indicates that individuals are at high risk of infection from poor hygiene and sanitary conditions.

In this study anaerobes were isolated in 33.1% of the total sampled specimens, most commonly *Bacteroides* spp., followed by *Peptostreptococcus* spp. and *Clostridium* spp.. This is comparable to findings from other reported studies^{28,16}. Other research findings have showed *Peptococcus* spp., *Peptostreptococcus* spp., and *Prevotella melaninogenica* as the commonest anaerobes^{29,30}. Some studies have reported isolation rates of anaerobes of 20% to 50%^{13,24}. Improved isolation of anaerobes can be attributed to improved collection, transportation, and processing techniques of

the specimens. Mixed aerobic and anaerobic cultures also characterize chronic infection in CSOM, suggesting a potential synergy between aerobic and anaerobic bacteria. It has been reported that polymicrobial infections are more pathogenic than monomicrobial infections³¹.

In this study, fungal isolates were identified in 21 (17.8%) of the sampled specimens, and *Aspergillus* spp. were the commonest, followed by *Candida* spp.; this is comparable with results published elsewhere³². Fungal infections of the middle ear are common as fungi thrive well in moist ears³³. The most commonly isolated fungi in CSOM are *Candida* and *Aspergillus*³⁴.

This study showed statistically significant characteristic associations of some CSOM-causing microorganisms with specific aspects of the CSOM symptomatology, which is contrary to a study where bacteriological findings had no significant effect on symptoms and signs³⁵.

One limitation for this study was that antibiotic susceptibility testing was not done. We recommend that future studies evaluate the drug sensitivities of local microorganisms that cause CSOM and subsequently identify changes in bacteriological profile and, possibly, the emergence of multidrug-resistant strains resulting from indiscriminate use of antibacterial agents.

Conclusions

Proteus mirabilis, *Pseudomonas aeruginosa*, and *Staphylococcus aureus* were the most prevalent aerobic bacteria, while *Bacteroides* spp. and *Peptostreptococcus* spp. were the commonest CSOM-causing anaerobic bacteria. These CSOM-causing microorganisms were predominant among males aged 18 years and below. The disparity in findings from previous research can be attributed to the variation in climate, community environment, patient population, and indiscriminate use of antibiotics. There was a correlation between particular CSOM-causing microorganisms and specific clinical features of CSOM.

References

1. Smith AW, Macharia I, Mugwe P, Hatcher J. Randomized control trial of treatment of chronic otitis media in Kenyan school Children. *Lancet*, 1996, 348: 1128-1133
2. Jose A. Chronic otitis media: Burden of Illness and Management .Child and Adolescent Health and Development Prevention of Blindness and Deafness. World Health Organization (WHO). Geneva, Switzerland, 2004.
3. Roland PS. Chronic otitis media: a clinical overview. *Ear Nose Throat J* 2002; 81: 8
4. Verhoeff M, et al. Chronic otitis media: a review. *Int J Pediatr Otorhinol* 2006; 70-71
5. Vartiainen E, Vartiainen J. Effect of aerobic bacteriology on the clinical presentation and treatment results of CSOM. *J Laryngol Otol* 1996; 110: 315
6. Brook I, Burke P. The management of acute, serous and CSOM: the role of anaerobic bacteria. *J Hosp Infect* 1992; 22 Suppl A: 75
7. Brook I. The role of anaerobic bacteria in otitis media: microbiology, pathogens-is, and implications on therapy. *Am J Otolaryngol* 1987; 8: 109
8. Adoga AS, Malu D, Badung BP, Obiesie IV. Swab and Aspiration collection methods and antibiograms in chronic otitis media at Jos University Teaching Hospital. Which is superior? *Ann Afr Med* 2010 vol 9. Issue 4:230-234

9. Poorley VK and Lyer A. Study of bacterial flora in chronic otitis media and its clinical significance. *Indian J Otolaryngol Head and Neck Surg* 2002;54:91-5
10. Mansoor T, Mussani MA, Khalid G, Kumal M. *Pseudomonas aeruginosa* in Chronic Otitis media: sensitivity spectrum against various antibiotics in Karachi. *J Ayub Med Coll Abbottabad* 2009; 21(2):120-23
11. Mawson S, Pollack M. Special role of *Pseudomonas aeruginosa* in chronic otitis media. *Ann Otol Rhinol Laryngol Head and Neck Surg.*, 1988, 97 (Suppl 130): 10-13 46: 488-97
12. Loy AHC, Tan AL, Lu PKS. Microbiology of chronic otitis media in Singapore. *Singapore Med J* 2002;43:296-9
13. Melaku A, Lulseged S. Chronic otitis media in a childrens hospital in Addis Ababa, Ethiopia. *Ethiop Med J.* 1999 Oct; (4): 237-46
14. Van Hasselt P, van Kregten E. Treatment of chronic otitis media with ofloxacin ear drops: A clinical/bacteriological study in a rural area of Malawi. *Int J Paediatr Otorhino* 2002;63(1):49-56
15. Olatoke F, Ologe FE, Nwawolo CC, Saka MJ. The prevalence of hearing loss among school children with chronic otitis media in Nigeria and its effect on academic performance. *Ear Nose Throat J* 2008;87(12):E19
16. Tiedt NJ, Butler IR, Atkins MD, Elliot E et al. Paediatric chronic otitis media in the Free State Province: Clinical and audiological features. *The South African Med J*; vol103, no 7(2013)
17. Ahmed, M., Ihsan E., Jassim M., Prevalence and patterns of chronic suppurative otitis media and hearing impairment in Basrah city *Journal of Medicine and Medical Sciences* Vol. 1(4) pp. 129-133 May 2010
18. Chowdhury M. A., Alauddin M., comparative study between tubotympanic and atticofurcal types of chronic suppurative otitis media *Bangladesh Med Res Counc Bull* 2002 Apr;28(1):36-44
19. Sheahan P., Donnelly M., Kane R., Clinical features of newly presented cases of CSOM *Journal of Laryngology and Otology* 2001;115:962-6
20. Kumar H, Seth S. Bacterial and Fungal study of 100 cases of Chronic otitis media. *J Clin Diag Res.*2011;5:1224-7
21. Shyamla R, Reddy SP. The study of bacteriological agents of chronic otitis media-aerobic culture and evaluation. *J Microbiol Biotechnol Res.* 2012;(2):152-62
22. Van Hasselt et al. Bacteriology of Chronic Otitis Media amongst children in Nkhosha District of Malawi *ENT and Audiology News* Vol. 12 May/June 2013
23. Macharia I, Mugwe P et al. Bacteriology of chronic otitis media in Garissa district, Kenya: A point prevalence study. *International Journal of Paediatric Otorhinolaryngology* vol 77, issue 7, July 2013, p1107-1111
24. Nyembue DT, Tshiswaka JM, Sabue MJ. Bacteriology of Chronic otitis media among Congolese children. *Acta Otolaryngol Belg.* 2003; 57(3):205-8
25. David S. Haynes. Peril-operative antibiotics in chronic otitis media. *Ear, Nose and Throat Journal*, August, 2002
26. Constanble L, Butler I. Microbial flora in chronic otitis media. *J Infect* 1982; 5: 57-60
27. Iqbal SM, Hassan A, Shafiq M. Chronic otitis media: disease pattern and drug sensitivity. *J Surg Pak* 2006; 11: 17-19
28. Nikakhlagh S, Khosrani A, Mand Rahidi N et al. Microbiological findings in patients with Chronic otitis media. *J Med Science.* 2008; 8(5): 503-506
29. Sweeney G, Picozzi G, Browning GG. A quantitative study of aerobic and anaerobic bacteria in CSOM. *J Infect* 1982; 5: 49-55
30. Maji PK, Chatterjee S, Chatterjee J et al. 2007. The investigation of bacteriology of chronic otitis media in patients attending a tertiary care hospital with special emphasis in seasonal variation. *Indian J. Otol Head and Neck Surg*, 59: 128-131
31. Brook I. Microbiology of polymicrobial abscesses and implication for therapy. *J Antimicrobial Chemother* 2002;50:805-1
32. Olu Ibekwe A, Zain Al Shareef, Ashraf Benayum. Anaerobes and Fungi in Chronic otitis media. *Ann Otol Rhinol Laryngol* 106: 1997;106:694-52
33. Tiwari S et al. Chronic bilateral Suppurative Otitis media caused by *Aspergillus terreus*, *Mycoses* 38(7-8)(1995).
34. Rajat P, Deepak J, Vikran N et al. Microbiology of Chronic Otitis media in a Tertiary Care Setup of Uttarakhand State. *N Am J Med Sci.* 2013 April; 5(4):282-285
35. Vartiainen E., Vartiainen J., Effects of aerobic bacteriology on clinical presentation and treatment results of CSOM *Journal Laryngology and Otology* 1996 April, 110(4):315-8