

Case report

Giant gastric lipoma mimicking well-differentiated liposarcoma

Mohamed Moncef Hamdane^{1,&}, Ehsen Ben Brahim¹, Mériam belhaj Salah¹, Nooman Haouas², Ahmed Bouhafa², Achraf Chedly-Debbiche²

¹Department of pathology, Habib Thameur Hospital. 8 rue Aly Ben Ayed. 1008. Montfleury, Tunis, Tunisia, ²Department of visceral surgery, Habib Thameur Hospital. 8 rue Aly Ben Ayed. 1008, Montfleury, Tunis, Tunisia

[&]Corresponding author: Mohamed Moncef Hamdane, 60 Avenue Bahi Ladgham, Ennasr 2, 2037. Ariana, Tunisia

Key words: Stomach, lipoma, liposarcoma, pathology, cytogenetics

Received: 9/13/2011 - Accepted: 9/2/2012 - Published: 9/23/2012

Abstract

Authors report the case of a 51-year-old man, presenting with epigastralgia of recent onset. Physical exam was unremarkable. Endoscopy revealed a large, ulcerated, submucosal, antral tumor. CT scan reveals an antral mass with fat attenuation. The patient underwent a total gastrectomy. Macroscopic examination identified in the antral wall a 9-cm, well-circumscribed, nodular lesion, with a yellow, greasy cut surface. On histological examination, the tumor was composed of a mature adipocytes proliferation, showing significant variation in cell size, associated to some lipoblasts. Nuclei were sometimes large, slightly irregular, but without hyperchromasia nor mitosis. Diagnosis of a well-differentiated liposarcoma was suspected and molecular cytogenetic analyses showed no MDM2 nor CDK4 gene amplification on fluorescent in situ hybridization. The diagnosis of lipoma was made. Twelve months following surgery, the patient is doing well.

Pan African Medical Journal. 2012; 13:16

This article is available online at: <http://www.panafrican-med-journal.com/content/article/13/16/full/>

© Mohamed Moncef Hamdane et al. The Pan African Medical Journal - ISSN 1937-8688. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Introduction

Lipomas are relatively uncommon tumors in the gastrointestinal tract, often located in the right colon [1]. Gastric lipomas are even more unusual with approximately 220 cases reported in the literature [2]. The diagnosis is strongly suggested by abdominal CT scan findings and is confirmed by histology. Histopathological diagnosis is usually easy. However, the tumor may sometimes undergo significant inflammatory changes leading to misdiagnosis of this lesion with a well-differentiated liposarcoma (WDLS).

Patient and observation

A 51-year-old man, in otherwise excellent general health, presented with epigastralgia of recent onset. The patient denied any history of gastrointestinal hemorrhage, nausea, vomiting, and change in bowel habits, fevers, or melena. Physical examination was unremarkable. Endoscopy revealed a soft, large, ulcerated, submucosal mass in the gastric antrum. Multiple biopsies were obtained but were all superficial, showing unspecific inflammation of the gastric mucosa. The abdominal CT scan revealed a round, well circumscribed, low-attenuation, gastric antral mass, measuring approximately 9 cm in diameter (**Figure 1**). The patient underwent total gastrectomy.

Gross examination of the surgical specimen identified in the antral submucosa, a well-circumscribed, smooth, nodular lesion, measuring 9x7.5x5 cm, with a homogeneous, yellow, greasy cut surface. Overlying mucosa was partially ulcerated (**Figure 2**).

Histological examination revealed a submucosal tumor (**Figure 3**), composed of a mature adipocytes proliferation, showing significant variation in cell size (**Figure 4**), associated to some lipoblasts (**Figure 5**), in a fibromyxoid background. Nuclei were sometimes large, slightly irregular, but without hyperchromasia nor mitosis. Many branching capillaries were seen. Areas of lower cellularity, displaying chronic inflammatory changes and containing bland spindle cells were also noted. The overlying mucosa was ulcerated (**Figure 6**) and the muscularis propria was focally dissociated by the tumor. The diagnosis of a WDLS was suspected. An immunohistochemical study was performed. Tumor cells were reactive with anti-HMGA2 and didn't express S-100 protein, CD34, MDM2 and CDK4. A cytogenetic study was then performed, showing no MDM2 nor CDK4 gene amplification on fluorescent in situ hybridization (FISH). The diagnosis of a benign lipoma was so made. The patient underwent an uneventful recovery. One year following surgery, he is doing well.

Discussion

Gastric lipoma is a rare, benign lesion, accounting for only 5% of gastrointestinal tract lipomas and fewer than 1% of all gastric tumors [3,4]. They typically occur in the fifth or sixth decade of life with equal sex incidence [2,4,5]. 75% of gastric lipomas are located in the antrum and are usually submucosal in origin [2].

Clinical symptoms depend on the size of the lesion. When small (< 2cm), lipomas are usually asymptomatic and they are often discovered fortuitously [1,3,4]. When the tumors are large (> 3–4cm), patients often present with upper gastrointestinal hemorrhage, either chronic or acute, caused by ulceration of the neoplasm [1,4]. Abdominal pain and obstructive symptoms are also common, especially if there is endoluminal growth that could cause intussusception [1,4].

Endoscopically, gastric lipomas typically appear as a soft, sharply defined, submucosal mass, often yellow in color as opposed to the surrounding pink mucosa. Typically, 3 clues help to identify these lesions as lipomas on endoscopic examination: the “tenting sign”, in which the overlying mucosa is easily retracted with the biopsy forceps; the “cushion sign” which occurs when the forceps produces a soft, cushioning indentation when applied to the lipoma; and the “naked fat” sign, which refers to exposed adipose tissue on the surface of the lipoma that pokes through the normal overlying mucosa after multiple biopsies of the normal mucosa are performed [6]. Occasionally, the lesion may be associated with a centrally located, superficial ulceration, caused by pressure necrosis. At times, ulceration can be fairly extensive, leading to the false impression of a more aggressive lesion [4,6]. Abdominal CT scan is the imaging examination of choice. It strongly suggests the diagnosis by showing a well-circumscribed lesion with a uniform, fatty density and an attenuation ranging from –70 to –120 H [4]. Endoscopic ultrasound examinations (EUS) is also a very good method for the diagnosis of gastric lipomas [2]. The typical findings of EUS reveal the tumor as an hyperechoic neoplasm in the submucosal layer.

Diagnosis is confirmed only after histopathological examination of surgical specimen. Standard biopsies are often inadequate because of the submucosal location of the tumor. On gross examination, gastric lipomas appear as solitary, smooth, soft masses, most often more than 2 cm in diameter, with an average size of 6.5 cm. On cut surface, they appear bright yellow, round, greasy, and encapsulated unless they have become infarcted. Large lesions may be ulcerated, like in our patient [1,2,4]. Histologically, gastric lipomas are sharply circumscribed tumors, usually surrounded by a thick fibrous capsule. They are composed of mature adipocytes, relatively uniform in size and lacking cytologic atypia. The tumor is usually centered in the submucosa and often compresses the overlying muscularis mucosae [1,7].

When the tumor is large, there is a progressive tendency for the submucosal mass to extrude into the lumen, leading to traumatic and inflammatory changes and resulting in necrosis, ulceration, and hemorrhage. Secondary changes including nuclear hypertrophy, hyperchromasia, fat necrosis, fatty cysts, and foamy macrophages may be observed. Lipoblasts and variation in cells size could, in addition, be present. The tumor can, in this situation, mimic a WDLS [3,7,8]. In such situation, diagnosis is difficult and morphological features are insufficient to classify the tumor. Cytogenetics and molecular biology offer new powerful tools for differentiating benign and malignant lipomatous tumors. WDLS are characterized by giant marker and ring chromosomes, sometimes as a sole finding or occasionally in association with other numerical or structural alterations. The giant marker and ring chromosomes contain amplified sequences of 12q13-15, the site of several genes (e.g., MDM2, GLI, SAS, CDK4, and HMGIC). This structural abnormality results in the consistent amplification of MDM2 and the frequent amplification of the adjacent genes, SAS, CDK4, and HMG1C [8]. The high specificity and sensitivity of detection of MDM2 and CDK4 amplification in WDLS and dedifferentiated liposarcomas have been demonstrated and analysis of these abnormalities using FISH or polymerase chain reaction has recently been shown to be an interesting means of identifying and separating WDSL from various benign lipomatous lesions [8-10].

The treatment of choice for a symptomatic gastric lipoma is circumferential excision with a clear margin of normal tissue. Due to the benign nature of this lesion an extensive surgical procedure such as an extended gastrectomy is not necessary. Simple local enucleation or partial gastric

resection is sufficient to remove the lipoma without fear of relapses or malignant degeneration [2]. The prognosis of patients with gastric lipomas is good. Malignant degeneration has not been reported [2].

Conclusion

Fatty tumors are rare in the gastrointestinal tract. Differentiating benign from malignant neoplasms is sometimes difficult in morphologic features. Cytogenetic procedures are, in these cases, the only means for an accurate diagnosis.

Competing interests

The authors declare no competing interests.

Authors' contributions

All the authors have contributed to this manuscript in ways that are compliant with the ICMJE authorship criteria. All the authors have read and approved the final version of this manuscript.

Figures

Figure 1: Abdominal CT scan: A large, low-attenuation, gastric mass

Figure 2: Macroscopic features: A well-circumscribed nodular lesion with a yellow, greasy, cut-surface

Figure 3: An adipocytic proliferation located in the submucosa (HEx40)

Figure 4: The tumor is made of an adipocytic proliferation showing a significant variation in cells size, with many branched capillaries, in a fibromyxoid background (HEx100)

Figure 5: Lipoblasts are present within the adipocytic proliferation (HEx400)

Figure 6: The overlying mucosa is focally ulcerated (HEx40)

References

1. Odze RD, Goldblum JR. *Surgical Pathology of the GI Tract, Liver, Biliary Tract and Pancreas*. 2nd ed. Saunders Elsevier: Philadelphia, 2009
2. Saltzman JR, Carr-Locke DL, Fink SA. Lipoma Case Report. *MedGenMed*. 2005 Jan 18;7(1):16. **This article on PubMed**
3. Fernandez MJ, Davis RP, Nora PF. Gastrointestinal lipomas. *Arch Surg*. 1983 Sep;118(9):1081-3. **This article on PubMed**
4. Thompson WM, Kende AI, Levy AD. Imaging characteristics of gastric lipomas in 16 adult and pediatric patients. *AJR Am J Roentgenol*. 2003; 181(4):981-985. **This article on PubMed**
5. Chu AG, Clifton JA. Gastric lipoma presenting as peptic ulcer: case report and review of the literature. *Am J Gastroenterol*. 1983; 78(10):615-618. **This article on PubMed**
6. Taylor AJ, Stewart ET, Dodds WJ. Gastrointestinal lipomas: a radiologic and pathologic review. *AJR Am J Roentgenol*. 1990; 155(6):1205-1210. **This article on PubMed**
7. Fenoglio-Preiser CM, Noffsinger AE, Stemmermann GN, Lantz PE, Isaacson PG. *Gastrointestinal Pathology: An Atlas and Text*, 3rd Edition. Lippincott Williams & Wilkins: Philadelphia, 2008
8. Weiss SW, Goldblum JR. *Enzinger and Weiss's Soft Tissue Tumors*, 5th ed. Mosby Elsevier: Philadelphia, 2008
9. Boltze C, Schneider-Stock R, Jäger V, Roessner A. Distinction between lipoma and liposarcoma by MDM2 alterations: a case report of simultaneously occurring tumors and review of the literature. *Pathol Res Pract*. 2001; 197(8):563-568. **This article on PubMed**
10. Shimada S, Ishizawa T, Ishizawa K, Matsumura T, Hasegawa T, Hirose T. The value of MDM2 and CDK4 amplification levels using real-time polymerase chain reaction for the differential diagnosis of liposarcomas and their histologic mimickers. *Hum Pathol*. 2006; 37(9):1123-1129. **This article on PubMed**




Figure 1

Abdominal CT scan: A large, low-attenuation, gastric mass



Figure 2

Macroscopic features: A well-circumscribed nodular lesion with a yellow, greasy, cut-surface




Figure 3
An adipocytic proliferation located in the submucosa (HEx40)




Figure 4

The tumor is made of an adipocytic proliferation showing a significant variation in cells size, with many branched capillaries, in a fibromyxoid background (HEx100)




Figure 5
Lipoblasts are present within the adipocytic proliferation (HEx400)




Figure 6
The overlying mucosa is focally ulcerated (HEx40)