

Supplementary Information, *S. Afr. J. Chem.*, 2015, **68**, 53–56

A Facile and Practical *p*-Toluenesulfonic Acid Catalyzed Route to Dicoumarols Containing an Aroyl group

Saeed Khodabakhshi^{1,*}, Bahador Karami², Khalil Eskandari^{2,*} and Alimorad Rashidi¹

¹*Nanotechnology Research Center, Research Institute of Petroleum Industry, Tehran, Iran*

²*Department of Chemistry, Yasouj University, P. O. Box 353, Yasouj, 75918-74831, Iran*

Table of contents

¹H NMR and ¹³C NMR spectra of products **3b**, **3e**, **3f**, **3g**, **3h**




Figure 1 ^1H NMR spectrum of **3b**




Figure 2 ^{13}C NMR spectrum of **3b**




Figure 3 ¹H NMR spectrum of **3e**




Figure 4 ^{13}C NMR spectrum of **3e**




Figure 5 ^1H NMR spectrum of **3f**




Figure 6 ^{13}C NMR spectrum of **3f**




Figure 7 ^1H NMR spectrum of **3g**




Figure 8 ^{13}C NMR spectrum of **3g**




Figure 9 ¹H NMR spectrum of **3h**




Figure 10 ^{13}C NMR spectrum of **3h**