



A review of the factors that influence erythrocyte osmotic fragility

NA Igbokwe

¹ Department of Veterinary Physiology and Biochemistry, Faculty of Veterinary Medicine, University of Maiduguri, PMB 1069 Maiduguri, Borno State, Nigeria

*Correspondence: Tel.: +2348060175771; E-mail: naigbokwe@gmail.com

Copyright: © 2018 Igbokwe. This is an open-access article published under the terms of the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Publication History:
Received: 02-12- 2017
Accepted: 22-06-2018

Abstract

Erythrocyte osmotic fragility is a laboratory test which evaluates the stability of the erythrocyte membrane to osmotic stress. The aim of this review is to highlight research findings on the intrinsic and extrinsic factors that influence erythrocyte osmotic fragility. The extrinsic factors include the type, ionic strength and pH of incubation media, type of anticoagulant and storage time of the blood, ambient temperature, drugs, medicinal plant extracts, xenobiotics, chemical agents; whereas intrinsic factors are age, sex, breed, species, pregnancy, lactation and genetic factors. Membrane composition, ion transports, aquaporin action, lipid peroxidation, and eryptosis of erythrocytes are involved in the variability of osmotic fragility. Increased osmotic fragility and improved osmotic stability of erythrocytes are pathophysiological phenomena that require appropriate interpretation in research or clinical investigation and the understanding of the factors affecting osmotic fragility will aid in the laboratory assessment of conditions related to erythrocytes where erythrocyte osmotic fragility test is used.

Keywords: Erythrocyte, Extrinsic factors, Intrinsic factors, Osmotic fragility, Osmotic stability

Introduction

Osmotic stability and fragility of erythrocytes are related measurable quantities in haematology which estimate haemolysis under hypoosmotic stress. In the interpretation of estimated haemolysis, an increased haemolysis is defined as an increase in osmotic fragility or a decrease in osmotic stability in an inverse corollary (Igbokwe, 2016). Erythrocyte osmotic fragility is determined by conducting a fragility test which is used as aid in the diagnosis of hereditary spherocytosis and haemolytic diseases. It supplements a stained cell examination to detect morphologic erythrocyte abnormalities and alterations leading to destruction of the erythrocytes (Kolanjippan *et al.*, 2002). It is used for studies of membrane permeability and as a potential

biomarker of oxidative membrane damage in pathologic conditions as well as toxicant or xenobiotic or pesticide-induced oxidative membrane damage to erythrocytes (Sharma *et al.*, 2010). It is also a tool for screening for alpha-thalassemia (Sirichotiyakul *et al.*, 2004), beta-thalassemia trait (Thool *et al.*, 1998; Bobhate *et al.*, 2002; Manju *et al.*, 2006), for pigmenturia (Legrange *et al.*, 1995) and for hypertensive patients who will benefit from diuretic therapy (Fansamade, 1999). Transportation stress (Minka & Ayo, 2010) and vitamin E status (Pillai *et al.*, 1992) can be assessed in animals by erythrocyte osmotic fragility test.

Some extrinsic factors which include temperature, pH of isotonic solution (Oyewale *et al.*, 1991;

Oyewale, 1991a; Oyewale, 1991b; Oyewale, 1993; Oyewale, 1994a; Oyewale *et al.*, 2011; Islah *et al.*, 2016), osmolarity and type of media (Igbokwe & Igbokwe, 2016a; Igbokwe & Igbokwe, 2016b), oxygenation (Lewis & Ferguson, 1966; Dacie & Lewis, 1995; Mafudvaze & Erlwanger, 2007), season (Oladele *et al.*, 2003; Habibu *et al.*, 2016), ionic strength of media (Igbokwe & Igbokwe, 2015), medicinal plants (de Freitas *et al.*, 2008) and drugs (Boelsterli *et al.*, 1983; Chikezie, 2007) can affect erythrocyte osmotic fragility. Certain intrinsic factors such as age (Perk & Perk, 1964a; Oyewale & Ajibade, 1990; Oyewale 1991b; Igbokwe *et al.*, 2016), genes (Krogmeier *et al.*, 1993; Gueorguiev *et al.*, 1999), species (Soliman & Amrousi, 1966; Coldman *et al.*, 1969; Aldrich *et al.*, 2006), breed (Schalm *et al.*, 1975; Oyewale & Durotoye, 1988; Olayemi & Oyewale, 2002; Habibu *et al.*, 2013), phenotype (Igbokwe *et al.*, 2015a), gender (March *et al.*, 1966; Habibu *et al.*, 2014; Igbokwe *et al.*, 2016), pregnancy and lactation (Habibu *et al.*, 2014; Igbokwe *et al.*, 2015b), egg laying (Oyewale, 1990), size (Troiano *et al.*, 2000) and differences in erythrocyte membrane composition (Fairley *et al.*, 1988; Hagve *et al.*, 1993) can also affect the osmotic fragility of erythrocytes. The resistance of erythrocytes to haemolysis may be increased or decreased in haematological disorders (Jain, 1973), glycemia (Sekar & Selvam, 1994) and disease processes (Kobo *et al.*, 2014). Glycosylation of haemoglobin and erythrocyte membrane proteins can also cause changes in osmotic fragility of erythrocytes (Ramana *et al.*, 1997). Chikezie *et al.* (2010) reported that differences in erythrocyte osmotic fragility of humans are under the control of individual genotype of erythrocytes. Variations in some physicochemical properties and oxidant levels of red blood cell genotypes contribute to differences in mechanical stabilities and capacities of erythrocyte to withstand osmotic stress (Srouf *et al.*, 2000a, Srouf *et al.*, 2000b; Senturk *et al.*, 2001; Richards *et al.*, 2007). Extracts of some medicinal plants (Suboh *et al.*, 2004; Akinwande *et al.*, 2007; Gomes de Sou *et al.*, 2007, Chikezie & Uwakwe, 2011) and haemoparasites can also affect erythrocyte fragility (Silva *et al.*, 1989; Santoro *et al.*, 1994). The aim of the present review was to identify and understand the factors that could influence erythrocyte osmotic fragility and how they affect erythrocyte osmotic stability.

Intrinsic Factors

Genetic factors

Decreased osmotic fragility in C57BL/6J strain of mice, compared to another strain DBA/2J, was directly controlled by their genotype (Dewey *et al.*, 1982) and the osmotic resistance allele was autosomal and recessive to the susceptible one (Schaefer & Dewey, 1989). Gene effect on osmotic fragility seemed to influence erythrocyte membrane ion transport in mice (Norman & Dewey, 1985; Armsby *et al.*, 1996) and correlated with milk fat traits in dairy cattle (Krogmeier *et al.*, 1993). Erythrocytes from sheep with extra α -gene were larger (with increased MCV) and had increased erythrocyte osmotic fragility than those of sheep without the genes (Pieragostini *et al.*, 2003). Chikezie *et al.* (2010) reported that osmotic fragility differed in three erythrocyte genotypes (HbAA, HbAS and HbSS) of males in humans. Erythrocytes with HbSS genotype exhibited the least stability and HbAA showed the most stability. Phenotypic drift caused physiological variation in erythrocyte osmotic fragility in Sahel goats (Igbokwe *et al.*, 2015a).

Species

Elephant erythrocytes are highly resistant to osmotic lysis when compared to erythrocytes from other species of mammals due to the increased surface to volume ratio of their large erythrocytes (Nirmalan *et al.*, 1967; Silva & Kuruwita, 1994). Erythrocyte osmotic fragility was reported to be highest in West African dwarf goat (0.55-0.75%); lowest in White Fulani cattle (0.35%-0.55%) and that of sheep was in between the two (0.45%-0.65%) when the three species were compared (Olusanya & Adepoju, 1979). Buffenstein *et al.* (2001) reported that erythrocytes from kangaroos were more resistant to osmotic stress than erythrocytes of sheep. In a study involving adult animals (Perk *et al.*, 1964b), osmotic fragility of erythrocytes of the following animals was reported in an increasing order; camel, chicken, dog, pig, rabbit, guinea pig, mouse, rat, hamster, horse, donkey, ox, cat, sheep and goat. Birds with elliptical erythrocytes are less osmotically fragile than species with round erythrocytes (Viscor & Palomeque, 1982). Erythrocytes from the guinea fowl are more fragile than those of the domestic fowl (Durotoye & Oyewale, 1988). Erythrocytes from chicken were reported to be more fragile than those of ostrich (Olowookorun & Makinde, 1998).

Breed

Erythrocyte osmotic fragility was lower in Nigerian indigenous dogs than in German shepherd dogs (Olayemi *et al.*, 2009; Ogunyemi & Olayemi, 2016). However, Adebisi *et al.* (2014) reported no difference in erythrocyte osmotic fragility between Rottweiler dogs and Nigerian indigenous dogs. Oyewale & Durotoye (1988) reported that erythrocytes from local Nigerian breeds were more susceptible to osmotic haemolysis than those of exotic breeds of fowl. Among exotic breeds, breed differences exist between the osmotic fragility of erythrocytes from White Leghorn and New Hampshire breeds of fowl, and Marshal, Ross and Hubbard-Anak cross breeds of broiler chickens (March *et al.*, 1966; Habibu *et al.*, 2013), with the Marshal breed erythrocyte as the most stable.

Age

Oyewale (1991b) reported that adult guinea fowls (156 weeks of age) had more osmotically fragile erythrocytes than the younger ones (21 weeks of age). Erythrocytes were osmotically more fragile in the young (8-10 weeks old) than adult (52-80 weeks old) local Nigerian ducks (*Anas platyrhynchos*) (Oyewale *et al.*, 1998a) and turkeys (Oyewale & Ajibade, 1990; Azeez *et al.*, 2011a) due to variation in nutrition and cholesterol-phospholipid ratio in the membrane. Erythrocyte osmotic fragility decreased as Gezel and Makoei breeds of sheep (Asri-Rezaei *et al.*, 2006) and cattle (Basarab *et al.*, 1980) grew older. Erythrocyte membrane stability increased from age 20 to 94 years in women (Penha-Silva *et al.*, 2007), while erythrocytes of premature infants were less osmotically fragile than those of full-term infants (Bautista *et al.*, 2003). In normal males between the ages of 18 and 78 years, age-related effect was shown to increase both the mean fragility of erythrocytes and the variability of the fragilities of the erythrocytes within individual blood sample (Bowdler *et al.*, 1981). Mosior & Gomulkiewicz (1988) reported decreased osmotic fragility in older human and bovine erythrocytes as their erythrocytes aged in circulation, while Rifkind *et al.* (1983) and Mosior & Gomulkiewicz (1985) reported an increase in osmotic fragility as human erythrocytes aged in circulation.

Sexual dimorphism hormones and other hormones

Erythrocyte osmotic fragility was higher in males than in females in cattle (Olayemi 2004, Olayemi, 2007), birds (March *et al.*, 1966; Oyewale &

Durotoye, 1988; Oyewale, 2004) and sheep (Durotoye 1987) probably due to stabilizing effect of estrogen. Erythrocytes from females were osmotically more fragile than those of males in dogs (Ogunyemi & Olayemi, 2016), African giant rats (Oyewale *et al.*, 1998b), goats (Habibu *et al.*, 2014), turkeys (Azeez *et al.*, 2011b) and humans (Olorunshola *et al.*, 2012). However, sexual dimorphism was absent in swine (Makinde, 1986), camels and donkeys (Oyewale *et al.*, 2011), cattle (Basarab *et al.*, 1980) and Sahel goats (Igbokwe *et al.*, 2016).

Circulating estrogen in high levels protected erythrocyte membrane and decreased erythrocyte fragility in cows (Olayemi, 2004; 2007) and birds (March *et al.*, 1966; Oyewale, 1990). Testosterone had no effect on erythrocyte osmotic fragility in birds (March *et al.*, 1966). Parathyroid hormone (PTH) caused an increase in osmotic fragility of human erythrocytes (Bogin *et al.*, 1982; Malachi *et al.*, 1986). PTH caused a significant influx of calcium into erythrocyte, which was not associated with potassium leak. The increased calcium influx affected the spectrin-actin of the cytoskeletal network of the erythrocyte and altered the stability and integrity of the cell membrane. Osmotic fragility of erythrocytes was increased in birds with hypothyroidism when compared with normal birds (Dariyerli *et al.*, 2004). Leylek *et al.* (1998) reported significant increase in erythrocyte fragility in groups of women with labour induction or augmentation with oxytocin than in other groups. Progesterone interacted with the soluble protein component of erythrocyte membrane and improved the stability of the membrane (Devenuto *et al.*, 1969) but progesterone reduced erythrocyte osmotic fragility of human erythrocytes (Kaya & Saito, 1985; Yoong *et al.*, 2003).

Pregnancy and lactation

Erythrocyte osmotic fragility increased in pregnant women due to destabilizing effect of progesterone on the membrane (Nakamura, 1983; Arora *et al.*, 1994; Emembolu & Mba, 1994) than non-pregnant women mostly in the third trimester of pregnancy (Magid *et al.*, 1982). However, no difference was recorded between mean erythrocyte fragility of pregnant and non-pregnant women (Suhail *et al.*, 2010) and Red Sokoto does (Habibu *et al.*, 2014). Krogmeier *et al.* (1993) reported a decrease in erythrocyte osmotic fragility in Holstein cows that were in late lactation as compared to those in early lactation. Habibu *et al.* (2014) reported no significant

difference in osmotic fragility between erythrocytes of lactating and non-lactating Red Sokoto does. In Sahel does, erythrocyte osmotic fragility decreased in late pregnancy due to improved membrane stabilization by progesterone and increased during lactation because of decreased concentration of cholesterol and triglycerides in the membrane (Igbokwe *et al.*, 2015b).

Membrane composition

The composition of the erythrocyte membrane determines its elasticity and deformability and influences the ability of the erythrocyte to withstand osmotic stress. In humans, low deformability, increased level of surface phosphatidylcholine and low stomatin level in erythrocytes were associated with increased erythrocyte osmotic fragility (Orbach *et al.*, 2017). The osmotic stability of erythrocytes in control and heat exposed hamsters was increased with the addition of linolenoyl and stearoyl sorbitol which revealed that unsaturated acyl groups of membrane lipids contribute to the higher osmotic stability of erythrocyte (Livne *et al.*, 1972). Raz & Livne (1973) reported that unsaturated fatty acids (stearic, oleic, linoleic and linolenic acids, the methyl esters of these acids, as well as their hydroxy analogs) are able to protect erythrocytes from hypotonic haemolysis. In human erythrocytes, linoleic acid caused haemolysis and increased erythrocyte osmotic fragility due to its detergent-like action (Thomas-George *et al.*, 1979). Matured human erythrocytes were osmotically more stable than reticulocytes due to increased membrane cholesterol (Karai *et al.*, 1982a) which was suggested to be due to the inactivation of lecithin-cholesterol acyltransferase (Karai *et al.*, 1982b). Hagve *et al.* (1993) observed an increased level of n-3 fatty acids in the erythrocyte membrane of healthy females fed fish oil for twenty-eight days which led to a decrease in osmotic fragility. A similar report was made by Kiron *et al.* (1994) in rainbow trout fish, but Kogawa *et al.* (1998) reported an increase in osmotic fragility of rabbit erythrocytes treated with free fatty acids. In camels, decreased erythrocyte osmotic fragility was attributed to high concentration of total lipids, cholesterol, proteins, sphingomyelin and phosphatidylcholine in the membranes when compared with the concentrations of these parameters in the erythrocyte membranes of sheep and goats (Al-Qarawi & Mousa, 2004). The variations observed in the lipid composition and structure of erythrocytes of humans, goats and sheep affect their ability to resist detergent induced

lysis (Koumanov *et al.*, 2005). Increased erythrocyte osmotic fragility was observed in rabbits fed high fat diets due to distortion of ion fluxes and membrane properties (Abdelhalim & Moussa, 2010). Rats fed dietary oils had decreased erythrocyte osmotic fragility indicative that the fatty acid composition of rat erythrocyte membranes was affected by the fatty acid composition of the dietary fats (Kirchgessner *et al.*, 1994). Mineo & Hara (2005) reported same in rat erythrocytes incubated in saline medium containing short-chain fatty acids.

Esterases are enzymes in erythrocyte membrane that catalyse the hydrolysis of an ester into its alcohol and acid (Fukami and Yokoi, 2012) and some of these esterases could be responsible for the resistance of erythrocytes to osmotic stress. Al-Qarawi & Ali (2003) reported that the activities of three esterases; aspirin esterase, cholinesterase, and nitrophenylacetate esterases were lowest in camels, highest in goats while those of cattle and sheep fell in between.

Extrinsic Factors

Incubation temperature and time

Erythrocyte osmotic fragility increased when incubation temperature was decreased in sheep (Oyewale 1991a), rat, rabbit, cattle, and pig (Oyewale, 1992), domestic fowls and guinea fowls (Oyewale, 1991b), pigeon, lizard and toad (Oyewale, 1994a) and humans (Murphy, 1967; Seeman *et al.*, 1969; Aloni *et al.*, 1977). In camels (Aloni *et al.*, 1977), ducks (Oyewale *et al.*, 1998a) and peafowls (Oyewale, 1994a), erythrocyte osmotic stability decreased as the incubation temperature increased. The osmotic resistance of erythrocytes exposed to detergents (Triton X-100[®] and sodium dodecyl sulfate) decreased with increase in incubation temperature (Bielawski, 1990). However, increasing the incubation temperature to 46⁰C for 1hr did not affect erythrocyte fragility in humans (Van der Walt & Russell, 1978).

Potential of hydrogen (pH) of media

An inverse relationship between the pH of the media and osmotic fragility of erythrocytes has been reported in camels (Aloni *et al.*, 1977; Islah *et al.*, 2016), sheep and goats (Oyewale, 1991a; Oyewale *et al.*, 1991), pigeon and lizard (Oyewale, 1994a), domestic fowl (*Gallus domesticus*) (Oyewale, 1991b), rat, cattle, pig, rat (Oyewale, 1992), and humans (Jacob & Parpart, 1931) with an increase in the pH of the hypotonic saline medium causing a decrease in erythrocyte osmotic fragility attributed to oxidative

damage from elevation of endogenous cytotoxic metabolites and release of metals with variable valencies from metalloproteins after acidification.

Anticoagulant and storage of blood sample

When used as an anticoagulant, ethylene diaminetetra-acetic acid (EDTA) increased erythrocyte osmotic fragility when compared with heparin (Kafka & Yermiahu, 1998). Mafudvaze & Erlwanger (2007) reported that blood from ostriches collected in EDTA, had an increased osmotic fragility compared to blood collected in heparin after 12 hours of storage. Erythrocytes stored over a long period of time develop artifacts (Veale *et al.*, 2011; van de Watering, 2011) and become very fragile (Epps *et al.*, 1994). Erythrocyte osmotic fragility increased after storage for 40 days when blood samples from dogs (Price *et al.*, 1988) and humans (Ogunro *et al.*, 2010) were preserved in citrate-phosphate-dextrose-adenine (CPDA-1) medium. A marked increase in osmotic fragility was observed when erythrocytes were stored in CPD-A2 for 42 hrs (Beutler *et al.*, 1982). Erythrocyte osmotic fragility was not affected after storing non-detergent washed human erythrocytes for 24 hours (Okwusidi, 2004, 2011) and 48 hours (Okwusidi, 2002).

Erythrocytes stored in glass bottles at 0°C showed increased haemolysis and osmotic fragility when compared to erythrocytes stored in polyvinylchloride (PVC) bags plasticized with di (2-ethylhexyl) phthalate (DEHP) at the same temperature. The addition of DEHP to erythrocytes stored in glass bottles at 0°C decreased haemolysis and osmotic fragility to levels equal to those in PVC bags (Yamamura *et al.*, 1991). Kanas *et al.* (2013) reported reduced erythrocyte osmotic fragility in blood stored in bags made with butyryl trihexyl citrate (BTHC) than in bis (2-ethylhexyl) phthalate (DEHP). A comparison of the osmotic fragility of erythrocytes from stored blood samples from various species revealed that after 24hrs at 10°C, osmotic fragility increased in erythrocytes of domestic fowl, lizard and toad (Oyewale, 1994b), goat and pig (Oyewale, 1993); but decreased in erythrocytes of cattle, mouse, rabbit and rat (Oyewale, 1993), pigeon and peafowl (Oyewale, 1994b). After 24hrs of storage at 4°C, the osmotic fragility of camel erythrocytes increased. Osmotic fragility of erythrocytes of camel and donkey also increased after 78hrs of storage at 4°C, (Oyewale *et al.*, 2011). Erythrocytes stored in isotonic phosphate-adenine-guanosine-glucose-saline-mannitol (PAGGSM) or erythrosol-4 had improved osmotic

stability when compared with erythrocytes stored in saline-adenine-glucose-mannitol (SAG-M) media due to the reduction in size (Veale *et al.*, 2011). Oxidative injuries to erythrocytes during storage led to membrane damage and made the erythrocytes more fragile with increased susceptibility to osmotic lysis (Chaudhary & Katharia, 2012).

Incubating media other than saline

The haemolysis of human erythrocytes in 0-10 g/dl glucose solutions was described by a decreasing sigmoid curve with complete or no haemolysis occurring at concentrations of 0-2 g/dl (0-68 mosmol/l) or 4-10 g/dl (137-342 mosmol/l), respectively (Lemos *et al.*, 2011). Glucose caused haemolysis in erythrocytes by altering the membrane (Marar, 2011). No haemolysis of human, rabbit, cattle, hamster, guinea pig, pig and sheep erythrocytes occurred in solutions of $\geq 0.4\%$ glucose (Matsuzawa & Ikarashi, 1979). Haemolysis of sheep erythrocytes was prevented in glucose concentration of 0-6%. As the concentration of glucose increased, the degree of haemolysis decreased from 100% to about 2% haemolysis in 0-4% glucose solution, but then rose in much higher concentrations for dog, mouse and rat erythrocytes (Matsuzawa & Ikarashi, 1979). Igbokwe & Igbokwe (2016a) reported a decrease in erythrocyte osmotic fragility of Sahel goats in hyposmolar concentrations of glucose when compared with similar concentrations of saline speculated to be influenced by glucose transporter protein and ion fluxes. Incubation in sucrose media reduced erythrocyte osmotic fragility in Sahel goats media due to decreased permeability of sucrose (Igbokwe and Igbokwe, 2016b) and in humans by inducing shrinkage (Martins *et al.*, 2012).

Nutrition

Erythrocytes were osmotically more stable in hypotonic solutions amongst well-nourished humans (Penha-silva *et al.*, 2007) and kangaroos (Buffenstein *et al.*, 2001) than from the malnourished which later became normal after nutritional therapy (Kaplay, 1978). Hyperglycaemia increased osmotic fragility of human erythrocytes (Jain, 1989).

Decrease in osmotic fragility and change in the shape of erythrocytes were reported in vitamin B6 deficient rats (Kual *et al.*, 1995). Pre-treatment of rats with vitamin C reduced erythrocyte osmotic fragility which was increased due to exposure to chlorpyrifus (Ambali *et al.*, 2010; Uchendu *et al.*, 2011). Vitamin C (ascorbic acid) administration before transportation protected the erythrocyte

membrane in pigs (Adenkola *et al.*, 2010a, Adenkola *et al.*, 2010b) and goats (Minka & Ayo, 2010) transported by road for several hours. Vitamins C and E supplementation also reduced erythrocyte osmotic fragility and oxidative damage in rats (Etlik *et al.*, 1997; Kraus *et al.*, 1997) and chickens (Azeez *et al.*, 2011a). In Gezel and Makoei breeds of sheep, increased vitamin E (α -tocopherol) levels associated with the increase in age caused a decrease in erythrocyte osmotic fragility (Asri-Rezaei *et al.*, 2006). Erythrocytes from vitamin E deficient sheep were protected against detergent induced haemolysis after the cells were preincubated with α -tocopherol *in vitro* (Stevenson & Jones, 1989). Vitamin E was reported to inhibit membrane peroxidation and protein oxidation and restored activities of superoxide dismutase and catalase on membrane of erythrocytes treated with mefenamic acid (Ahmad & Suhail, 2002) and glucose (Marar, 2011) thereby improving erythrocyte osmotic stability. Marques *et al.* (1986) reported the possibility of a toxic effect of higher doses of vitamin E supplementation rather than a protective role on erythrocyte membrane stability after *in vitro* treatment of erythrocytes with alpha-tocopherol.

In vivo treatment with vitamin E decreased erythrocyte osmotic fragility in patients on haemodialysis and peritoneal dialysis (Uzum *et al.*, 2006). Rats treated with acetaminophen had decreased erythrocyte glutathione content and activity of Na^+ K^+ -ATPase enzyme with increased osmotic fragility but supplementation with vitamin E restored the glutathione content, Na^+ K^+ -ATPase activity and osmotic fragility to normal (Suhail & Ahmad, 1995). Treatment with vitamin C (Alhassan *et al.*, 2010), E or combination of vitamins E and C during hot-dry season increased packed cell volume (PCV) and haemoglobin concentration but decreased erythrocyte osmotic fragility in Wistar rats (Wahab *et al.*, 2010).

Extracts of leaves, plants and herbs

Leaf extract of *Carica papaya L* stabilized erythrocyte membrane and reduced osmotic fragility in dengue patients (Ranasinghe *et al.*, 2012). Aqueous extracts of *Anacardium occidentale*, *Psidium guajava*, and *Terminalia catappa* leaves (Chikezie & Uwakwe 2011) and methanol extract of *Solanum aethiopicum* (Anosike *et al.*, 2012) reduced the osmotic fragility of human erythrocytes by stabilizing the membrane. Crude extracts of *Artemisia absinthium*, *Lippia sp.*, *Cymbopogon citratus* and *Mentha villosa* decreased erythrocyte osmotic fragility while extracts of

Bryophyllum sp. and *Solidago microglossa* increased erythrocyte osmotic fragility in humans (de Freistas *et al.*, 2008). Biltto *et al.* (2012) reported that pre-incubation of erythrocytes with rutin and α -naphtho flavone did not have any effect on osmotic fragility but quercetin and 3, 5, 7-tri- hydroxy-4'-methoxy flavone-7-rutinoside reduced osmotic fragility of human erythrocytes. Oral administration of ethanol extracts of *Jatropha gossypifolia* (Pohl) (Oyedemi *et al.*, 2015a) and *Adenopus breviflorus* (Benth) (Oyedemi *et al.*, 2015b) did not affect erythrocyte osmotic fragility in male Wistar rats. Ferrali *et al.* (1997) also reported that quercetin reduced osmotic fragility of rat erythrocytes. A mixture of flavonoids decreased erythrocyte osmotic fragility in rats experimentally infected with *Trypanosoma brucei* (Kobo *et al.*, 2014). Increased osmotic fragility due to membrane damage was reported in rat erythrocytes treated *in vivo* and *in vitro* with onions (*Allium cepa*) and garlic (*Allium sativa*) (Salami *et al.*, 2012). Vanillin, a naturally occurring food-flavoring agent protected erythrocytes against carbon tetrachloride (CCl_4)-induced erythrocyte damage in Wistar albino rats by decreasing osmotic fragility, lipid peroxidation and degradation of membrane proteins (Makni *et al.*, 2012). Erythrocytes of rabbits fed diets containing *Hibiscus sabdariffa*, a plant reported to have antioxidant effect improved osmotic stability when compared with erythrocytes from rabbits that were sham treated (Adenkola & Oluremi, 2014). Aqueous extract of *Alstonia congolensis* increased erythrocyte osmotic fragility in guinea pigs (Akinwande *et al.*, 2007).

Sharma *et al.* (1981) reported increased erythrocyte osmotic fragility in calves poisoned by lantana. Erythrocyte osmotic fragility increased due to lipid peroxidation and oxidative stress in cattle intoxicated with *Senico sp* (Bondan *et al.*, 2005). Gossypol, a yellow polyphenolic compound, from whole cottonseed or cotton seed meal had been reported to increase the osmotic fragility of lactating cows (Mena *et al.*, 2004), growing heifers (Colin-Negrete *et al.*, 1996), cattle (Wyse *et al.*, 1991), sheep and goats (Menges 1991; Matondi *et al.*, 2007). Herman (1969) reported that extracts of *Plasmodium lophurae* increased erythrocyte osmotic fragility in ducks.

Transportation stress

Erythrocyte osmotic fragility increased in pigs (Adenkola and Ayo, 2009) and goats (Minka & Ayo, 2010) but decreased in Nera black chicken (Azeez *et al.*, 2011a) and humans (Olorunshola *et al.*, 2012)

after transportation stress. Olaifa *et al.* (2012) reported increased erythrocyte osmotic fragility in donkeys due to post packing stress.

Exercise

In horses, aerobic exercise increased the temperature and pH of blood, reduced the size of the erythrocytes due to changes in the lipid composition and protein structure of the cell membrane and then decreased the erythrocyte osmotic fragility (Hanzawa *et al.*, 1996). However, anaerobic exercise decreased the pH of blood, increased the size of the erythrocytes and increased erythrocyte osmotic fragility (Hanzawa *et al.*, 1996; Hanzawa, 2000). Hanzawa *et al.* (1999) observed that regardless of splenic contraction for the release of erythrocytes, osmotic fragility increased with heavy exercise in normal and splenectomized horses.

Increased erythrocyte osmotic fragility caused by high-intensity exercise was recorded in horses (Hanzawa *et al.*, 1998) and humans (Davis & Brewer, 1935; Anuradha *et al.*, 1995). Yusof *et al.* (2007) reported a decrease in erythrocyte osmotic fragility due to *in vivo* alterations in erythrocyte membrane proteins, especially in the α - and β -spectrins in six non-smoking experienced male ultra-marathon runners. Increased erythrocyte osmotic fragility was reported in humans after whole body vibration exercise (Monteiro *et al.*, 2013). Erythrocyte osmotic fragility decreased in the rainbow lizard (*Agama agama*) after swimming exercise (Azeez & Oyewale, 2010).

Season

Habibu *et al.* (2016) reported that erythrocytes from Red Sokoto and Sahel goat kids were osmotically more fragile in the hot-dry than in the cold-dry season. Adenkola *et al.* (2011) reported that the osmotic fragilities of goat and cattle erythrocytes were higher during the cold harmattan than in hot dry season. However, osmotic fragility of erythrocytes from rats kept at hypothermic temperature decreased when compared with those from rats kept at room temperature (Peinado *et al.*, 1993). Increased erythrocyte osmotic fragility was seen in heat-exposed hamsters due to changes in membrane properties than those not exposed to heat (Kuiper *et al.*, 1971; Livne *et al.*, 1972).

Water deprivation

Increased erythrocyte osmotic fragility was reported in chickens (Yagil *et al.*, 1976) and Peking ducks

(Baloyi *et al.*, 2006) deprived of water for more than 24 hours. However, Mafudvaze *et al.* (2008) did not observe any change in erythrocyte osmotic fragility of guinea fowls deprived of water for up to 48 hours when compared with those of guinea fowls that were drinking water *ad libitum*. Erythrocytes of dehydrated camels were osmotically more stable in hypotonic saline solutions than those of hydrated camels (Yagil *et al.*, 1974).

Diseases

A two-fold increase in erythrocyte osmotic fragility was reported in cattle infected with *Anaplasma marginale* which correlated positively with intra-erythrocytic parasitaemia (Silva *et al.*, 1989). Increased erythrocyte osmotic fragility was also observed in cattle affected with trypanosomiasis and theileriosis (Pati *et al.*, 2017). In camels, erythrocyte osmotic fragility increased due to direct attack of reactive oxygen species on the erythrocyte's plasma membrane (Saleh *et al.*, 2009). Rats infected experimentally by *Trypanosoma evansi* (Mijares *et al.*, 2010) and *Trypanosoma brucei* (Ikejiani, 1946; Oyewale, 1987; Kobo *et al.*, 2014) had increased erythrocyte osmotic fragility. Lambs experimentally infected with *Dictyocaulus filariae* had increased erythrocyte osmotic fragility (Sharma *et al.*, 1989). Makinde & Bobade (1994) reported an increase in erythrocyte osmotic fragility in dogs infected with *Babesia canis* and *Ehrlichia canis*. Dogs diagnosed with leptospirosis had decreased erythrocyte osmotic fragility (Santoro *et al.*, 1994).

Erythrocytes of dogs affected with a heritable muscle disorder that clinically resembles a muscular dystrophy had decreased osmotic fragility (Abhold *et al.*, 1983). Erythrocytes from patients with Duchenne muscular dystrophy had increased osmotic fragility (Kim *et al.*, 1980). Cattle with traits of "double muscle" had increased erythrocyte osmotic fragility (Basarab *et al.*, 1980).

There was an increase in osmotic fragility as well as fluctuation in the lipid fraction of erythrocytes in elderly patients with anaemia (Tadano *et al.*, 1981). Human patients diagnosed with hereditary stomatocytosis and haemolytic anaemia had increased autohaemolysis and osmotic fragility (Mutoh *et al.*, 1983). Erythrocyte osmotic fragility decreased in dogs with immune-mediated anaemia (Paes *et al.*, 2013) and humans with sickle cell anaemia (Harris *et al.*, 1956; Dash & Kar, 1999) but increased in cats with anaemia (Kohn *et al.*, 2000; Tritschler *et al.*, 2016). Normal erythrocyte osmotic fragility was reported in a girl diagnosed with

hereditary spherocytosis (Korones & Pearson, 1988) and in dogs with malignant hyperthermia (Cribb *et al.*, 1986). Fansmade (1999) reported that osmotic fragility of erythrocytes was higher in hypertensive than in normotensive patients.

In alloxan-induced diabetic rats, erythrocyte osmotic fragility increased due to oxidative stress which was corrected after placing the rats on a diet containing antioxidants (Kowluru *et al.*, 1996). In humans, increased erythrocyte osmotic fragility was reported in diabetic patients (Agte *et al.*, 2004; Ibang *et al.*, 2005; Kung *et al.*, 2009; Mostafavi *et al.*, 2013). Increased erythrocyte osmotic fragility was reported in rats (Endoh *et al.*, 1992) and rabbits (Yuan *et al.*, 1988) with burnt body surfaces. Erythrocyte osmotic fragility increased in hyperlipidemic and dyslipidemic dogs probably due to alteration of cholesterol content in the membrane (Behling-Kelly & Collins-Cronkright, 2014).

In cervical cancer patients, decreased erythrocyte membrane vitamin E concentrations led to reduced erythrocyte glutathione which in turn led to an increase in erythrocyte osmotic fragility (Kolanjiappan *et al.*, 2002). The fragility of erythrocytes increased in rats inoculated with a variant of Walker 256 rat tumour cell (Cavalcanti *et al.*, 2003) but the fragility decreased in anaemia caused by Walker 256 rat tumour cell (Vido *et al.*, 2000). An increase in erythrocyte fragility was reported in buffaloes with dystocia (Prabhakar *et al.*, 2013).

Chemicals

Wang *et al.* (2010) reported membrane damage, haemolysis, K^+ leakage, alterations in erythrocyte shape, increased erythrocyte osmotic fragility, and inhibition of enzymatic activity after treatment with the melamine-cyanurate complex in humans. Melamine or cyanuric acid alone had no effect on erythrocyte membrane. Potassium bromate ($KBrO_3$) reduced glutathione content, induced oxidative stress on erythrocyte, impaired the anti-oxidant defense system and increased erythrocyte osmotic fragility in humans (Ahmad & Mahmood, 2012; Ahmad *et al.*, 2014). Ethanol created pores on the membrane of human erythrocytes, causing leakage of K^+ from the cell which led to haemolysis and increased osmotic fragility in a dose-dependent fashion (Chi & Wu, 1991; Tyulina *et al.*, 2002).

Organophosphorus insecticides (methylbromphenvinphos, diclorvos, malathion and methyl parathion) increased the resistance of pig erythrocytes to lysis and, thus, decreased their

osmotic fragility (Blasiak *et al.*, 1991). A significant increase was recorded in the erythrocyte osmotic fragility of Wistar rats after chronic exposure to chlorpyrifos when compared with control rats (Ambali *et al.*, 2010). Swiss albino rats exposed to ethanol in drinking water had increased erythrocyte osmotic fragility due to increased cholesterol content and cholesterol to phospholipid ratio in erythrocyte membrane (Sözmen *et al.*, 1994). Exposure of adult male black Nera chickens to iodosteryl (a disinfectant) in drinking water for six weeks increased erythrocyte osmotic fragility (Azeez *et al.*, 2012) due to intravascular haemolysis.

Drugs

A reduction in erythrocyte production and an increase in osmotic fragility was reported in rhesus monkeys treated with broad-spectrum antiviral ribavirin (1-beta-D-ribofuranosyl-1, 2, 4-triazole-3-carboxamide) (Canonica *et al.*, 1984). Pharmacological dosages of methylprednisolone produced a decrease in membrane deformability of erythrocytes in cardiac surgical patients leading to decreased fragility (Rand *et al.*, 1997). A gradual and significant decrease in osmotic fragility with an increase in concentration of saline solutions was observed in dogs under pentobarbitone anaesthesia (Olaifa *et al.*, 1999). Lidocaine protected erythrocytes against oxidative stress by inhibiting potassium efflux and delaying the occurrence of haemolysis (Lenfant *et al.*, 2000). Biltto (1990) reported a significant increase in osmotic fragility after *in-vitro* incubation of human erythrocytes for 30 to 60 minutes in 1, 2 and 4mg/mL of aspirin. Chlorpromazine, a tranquilizer has the ability to protect human erythrocytes from osmotic haemolysis by increasing the mean cellular volume and changing the shape of the cell in an isotonic medium (Freeman & Spirtes, 1963; Kwant & Van Steveninck, 1968). Low concentration of phenothiazine reduced osmotic fragility of erythrocytes in hypotonic sodium chloride solutions (Seeman *et al.*, 1969). Ebselen was reported to reduce erythrocyte osmotic fragility in humans by acting as an inhibitor of haemoglobin glycation after glucose-induced haemolysis (Soares *et al.*, 2014).

Detergents

Triton® X-100 (non ionic detergent) and sodium dodecyl sulfate (anionic detergent) increased erythrocyte osmotic fragility by the formation of pores in various membrane lipid regions, releasing

vesicles from erythrocytes (Bielawski, 1990; Chernitsky & Senkovich, 1997) and time dependence of the opening probability of these pores (Chernitsky & Senkovich, 1998). Parameters of detergent-induced haemolysis are sensitive to the changes of the charge and structural state of erythrocyte membrane (Chernitsky *et al.*, 2001). Triton[®] X-100 caused swelling and pores in the membrane followed by the haemolysis of erythrocytes while sodium dodecyl sulfate fragmented and destroyed erythrocytes. The rate of haemolysis increased with an increase in detergent concentration (Bielawski, 1990). Gaehtgens & Benner (1974) reported that three detergents (Pluronic[®] F 38, F 68, and F 108) exhibited anti-haemolytic effect and decreased osmotic fragility in human erythrocytes.

Metals

Elevated erythrocyte calcium caused a progressive increase in erythrocyte osmotic fragility (Cueff *et al.*, 2010; Costa *et al.*, 2010). Calcium levels of erythrocytes from goats were reduced by blocking calcium channels using diltiazem which decreased the level of oxidative damage (Das & Bhattacharyya, 2010). Erythrocyte deformability, membrane fluidity, and osmotic fragility improved significantly with nifedipine therapy in patients with systemic scleroderma. Impairment of ATP-Ca²⁺ pump occurred in the patients and there was the accumulation of Ca²⁺ in the erythrocytes which increased osmotic fragility (Spengler *et al.*, 2007). Mostafavi *et al.* (2013) reported that intra-erythrocytic calcium reduced erythrocyte osmotic fragility in patients with type 2 diabetes.

Humans diagnosed with iron deficiency anaemia (Kirchgeßner *et al.*, 1994) and iron deficient pregnant rats (Al-Hashimi *et al.*, 2015) had decreased erythrocyte osmotic fragility.

Osmotic fragility of erythrocytes obtained from zinc-deficient rats was increased when compared with that from normal rats (O'Dell *et al.*, 1987; Kraus *et al.*, 1997). Roozbeh *et al.* (2009) reported a significant decrease in osmotic fragility of erythrocytes from humans on haemodialysis administered zinc supplements for 6 weeks.

Karuppasamy *et al.* (2005) reported an increase in erythrocyte fragility of *Channa punctatus* fish exposed to sub lethal dose of cadmium. However, the ingestion of cadmium in food and drinking water by Swiss male rats did not cause any significant change in erythrocyte osmotic fragility (Demir & Öner, 1995).

Erythrocytes from copper-deficient rats had increased membrane lipid peroxidation and lower osmotic fragility than erythrocytes from rats fed diets containing various concentrations of copper (Jain & Williams, 1988). Toxic concentrations of copper in the blood resulted in increased membrane permeability and erythrocyte osmotic fragility in rats (Kirchgeßner *et al.*, 1994). Copper-dextran complex (C-79) in doses of 0.4-0.8mg/kg stabilized the erythrocyte membrane for several hours but lower doses produced a weak stabilizing effect while a dose of 1.6mg/kg increased erythrocyte osmotic fragility in rabbits (Debowy *et al.*, 1985). Hong-Wei *et al.* (2008) reported an increase in erythrocyte osmotic fragility, the rate of haemolysis and damage to erythrocytes when high concentrations of copper sulphate were added to media in which erythrocytes were incubated.

Chronic exposure of people working in lead refining factories to lead (Pb) caused alterations in membrane proteins which decreased the permeability of erythrocyte membrane and erythrocyte osmotic fragility (Karai *et al.*, 1981, 1982a, Fukumoto *et al.*, 1983). *In vitro* treatment of erythrocytes of rats (Levander *et al.*, 1977) and humans (Qazi *et al.*, 1972; Lessler & Walters, 1973; Mrugesh *et al.*, 2011) with Pb decreased osmotic fragility. Decreased mean corpuscular volume (MCV) and erythrocyte osmotic fragility were reported in lead poisoning sequel to intracellular water leakage accompanied with K⁺ loss due to erythrocyte membrane damage caused by the increased cholesterol level recorded in the erythrocyte membrane (Karai *et al.*, 1979; Karai *et al.*, Karao *et al.*, 1982a; Karai *et al.*, 1982b). However, Oyedeki & Alabi (2015) reported no change in erythrocyte osmotic fragility of male Wistar rats after oral administration of lead acetate for five weeks.

Mercury increased erythrocyte osmotic fragility in humans (Lessler & Walters, 1973). Okuda & Tsuzuki (1977) reported decreased erythrocyte osmotic fragility in low doses of methylmercury and no change in osmotic fragility in higher doses in male Wistar rats. Erythrocyte osmotic fragility decreased in female mice fed methyl mercury (10nmol/g feed) (Yamamoto & Suzuki, 1982). Erythrocytes treated with mercuric ions showed resistance to osmotic shock after 5 minutes of incubation but they began to haemolyse when the incubation time was increased (Zolla *et al.*, 1994). *In vitro* treatment of human erythrocytes with mercuric chloride resulted in shrinking of the erythrocytes and conferred protection against osmotic haemolysis (Mel & Reed,

1981). Igbokwe *et al.* (2018) reported an increase in osmotic stability of Sahel goat erythrocytes incubated in saline, glucose or sucrose media after *in vitro* treatment with mercuric chloride due to inhibition of osmotic permeability by blockage of aquaporins or by steric mechanism.

Various concentrations of aqueous solutions of tris acetylacetonate aluminium (III) (Al (acac)₃, tris maltolate aluminium (III) (Al(malt)₃ and tris aluminium lactate (Al(lac)₃) caused erythrocytes to be osmotically fragile (Zatta *et al.*, 1989). Treatment of rat erythrocytes with aluminium chloride caused increased osmotic fragility due to increased lipid peroxidation (Gutteridge *et al.*, 1995; Hernández *et al.*, 2008; Al-Qayim *et al.*, 2014; Oztürk & Ozdemir, 2015; Zhang *et al.*, 2016). However, Bazzoni *et al.* (2005) reported improved stability of erythrocyte membrane and reduced erythrocyte size and aggregation in adult male rats after parenteral treatment with aluminium hydroxide.

In conclusion, many intrinsic and extrinsic factors as seen in this review can affect erythrocyte osmotic fragility or stability. A good understanding of these factors will guide in blood sample collection, preservation and analysis such that inferences can be drawn from erythrocyte osmotic fragility tests with little or no interference from the factors.

References

Abdelhalim MA & Moussa SA (2010). Biochemical changes of hemoglobin and osmotic fragility of red blood cells in high fat diet rabbits. *Pakistan Journal of Biological Sciences*, **13**(2): 73-77.

Abhold RH, Hegreberg GA & Magnuson JA (1983). Decreased osmotic fragility in heritable canine myopathy. *American Journal of Hematology*, **15**(1): 15-22.

Adebiyi OE, Ajayi JO & Olayemi FO (2014). Haematology of Rottweiler dog in a tropical environment. *New York Academy of Sciences Journal*, **7**(9): 1-4.

Adenkola AY, Agbendeh J & Okpe J (2011). Comparative assessment of erythrocyte osmotic fragility of apparently healthy goat and cattle during the hot-dry and harmattan season in Markurdi, Nigeria. *Journal of Animal and Plant Sciences*, **11**(3): 1474-1480.

Adenkola AY & Ayo JO (2009). Effect of road transportation on erythrocyte osmotic fragility of pigs administered ascorbic acid during the harmattan season in Zaria,

Nigeria. *Journal of Cell and Animal Biology*, **3**(1): 4-8.

Adenkola AY, Ayo JO, Sackey AKB & Adelaiye AB (2010a). Erythrocyte osmotic fragility of pigs administered ascorbic acid and transported by road for short-term duration during the harmattan season. *African Journal of Biotechnology*, **9**(2): 226-233.

Adenkola AY, Kaankuka FG, Ikyume TT, Ichaver IF & Yaakugh IDI (2010b). Ascorbic acid effect on erythrocyte osmotic fragility, haematological parameters and performance of weaned rabbits at the end of rainy season in Markurdi, Nigeria. *Journal of Animal and Plant Sciences*, **9**(1): 1077-1085.

Adenkola AY & Oluremi OIA (2014). Erythrocyte osmotic fragility and excitability score in rabbit fed *Hibiscus sabdariffa* in graded level. *Nigerian Journal of Physiological Sciences*, **29**(2): 113-117.

Agte VV, Nagmote RV & Tarwadi KV (2004). Comparative *in vitro* uptake of zinc by erythrocyte of normal Vs type 2 diabetic individuals and the associated factors. *Diabetes Nutrition and Metabolism*, **17**(6): 343-349.

Ahmad I & Suhail M (2002). Protective role of vitamin E in mefenamic acid-induced alterations in erythrocytes. *Biochemistry (Moscow)*, **67**(8): 945-948.

Ahmad MK, Amani S & Mahmood R (2014). Potassium bromate causes cell lysis and induces oxidative stress in human erythrocytes. *Environmental Toxicology*, **29**(2): 138-145.

Ahmad MK & Mahmood R (2012). Oral administration of potassium bromate, a major water disinfection by-product, induces oxidative stress and impairs the oxidant power of rat blood. *Chemosphere*, **87**(7): 750-756.

Akinwande AI, Adeyemi OA & Odiri BO (2007). Effect of aqueous extract of *Alstonia congensis* on glucose transport and Na⁺-K⁺-ATPase activity in the everted and non-everted Guinea pig intestine and erythrocyte fragility. *Nigerian Journal of Health and Biomedical Sciences*, **6**(1): 5-9.

Aldrich KJ, Saunders DK, Sievert LM & Sievert YG (2006). Comparison of erythrocyte osmotic fragility among amphibians, reptiles, birds

- and mammals. *Transactions of the Kansas Academy of Science*, **109**(3): 149-158.
- Al-Hashimi LM, Gambling L & McArdie HJ (2015). The effect of iron deficiency on osmotic sensitivity of red blood cells from neonatal rats and their mothers. *Journal of Membrane Biology*, **248**(6): 1199-1206.
- Alhassan AW, Adenkola AY, Yusuf A, Bauchi ZM, Saleh MI & Ochigbo V (2010). Erythrocyte osmotic fragility of Wistar rats administered ascorbic acid during the hot dry season. *Journal of Cell and Animal Biology*, **4**(2): 29-33.
- Aloni B, Eitan A & Livne A (1977). The erythrocyte membrane site for the effect of temperature on osmotic fragility. *Biochimica et Biophysica Acta*, **465**(1): 46-53.
- Al-Qarawi AA & Ali BH (2003). Variations in the normal activity of esterases in plasma and liver of camels (*Camelus dromedarius*), cattle (*Bos indicus*), sheep (*Ovis aries*) and goats (*Capra hircus*). *Journal of Veterinary Medicine. A Physiology, Pathology, Clinical Medicine Journal*, **50**(4): 201-203.
- Al-Qarawi AA & Mousa HM (2004). Lipid concentrations in erythrocyte membranes in normal, starved, dehydrated and rehydrated camels (*Camelus dromedarius*) and in normal sheep (*Ovis aries*) and goats (*Capra hircus*). *Journal of Arid Environments*, **59**(4): 675-683.
- Al-Qayim MAJ, Ghali LS & Al-Azwai TS (2014). Comparative effects of propolis and malic acid on hematological parameters of aluminium exposed male rats. *Global Journal of Biochemistry and Biotechnology*, **3**(1): 6-11.
- Ambali SF, Ayo JO, Ojo SA & Esievo KAN (2010). Ameliorative effect of vitamin C on chlorpyrifos-induced increased erythrocyte fragility in Wistar rats. *Human and Experimental Toxicology*, **30**(1): 19-24.
- Anosike CA, Obidoa O & Ezeanyika LUS. (2012). Membrane stabilization as a mechanism of the anti-inflammatory activity of methanol extract of garden egg (*Solanum aethiopicum*). *DARU Journal of Pharmaceutical Sciences*, **20**(1): 76-80.
- Anuradha CV, Balakrishnan SD & Venugopal PM (1995). Increased erythrocyte lipid-peroxidation and osmotic fragility in sports people. *Medical Science Research*, **23**(6): 409-412.
- Armsby CC, Stuart-Tilley AK, Alper SL & Brugnara C (1996). Resistance to osmotic lysis in BXD-31 mouse erythrocytes: association with upregulated K-Cl cotransport. *American Journal of Physiology*, **270**(3Pt 1): C866-C877.
- Arora B, Punia RS, Lal P & Arora DR (1994). The effect of pregnancy on erythrocyte osmotic fragility. *Journal of Nepal Medical Association*, **32**(112): 227-230.
- Asri-Rezaei S, Ramin AG & Koukalanifar M (2006). Evaluation of correlation between serum vitamin E status and the erythrocyte osmotic fragility in sheep. *Journal of Faculty of Veterinary Medicine University of Tehran*, **61**(2): 155-159.
- Azeez OI, Olayemi FO & Olanrewaju JR (2011b). Age and sex influences on the haematology and erythrocyte osmotic fragility of the Nigerian turkey. *Research Journal of Veterinary Sciences*, **4**(2): 43-49.
- Azeez OI, Oyagbemi AA & Iji OT (2012). Haematology and erythrocyte fragility indices in domestic chicken following exposure to a polyvalent iodophorus disinfectant. *Jordan Journal of Biological Sciences*, **5**(2): 99-103.
- Azeez OI, Oyagbemi AA & Oyewale JO (2011a). Erythrocyte membrane stability after transportation stress in the domestic chickens as modulated by pretreatment with vitamin C and E. *Journal of Animal and Veterinary Advances*, **10**(10): 1273-1277.
- Azeez OI & Oyewale JO (2010). Effects of swimming exercise on erythrocyte osmotic fragility of the rainbow lizard (*Agama agama*). *Russian Journal of Herpetology*, **17**(3): 18-188.
- Baloyi C, Khosa F, Tembani S & Erlwanger KH (2006). The osmotic fragility of erythrocytes from Pekin ducks deprived of water for 24 hours. *South African Journal of Science*, **102**(1-2): 19-20.
- Basarab JA, Berg RT & Thompson JR (1980). Erythrocyte fragility in "double-muscling" cattle. *Canadian Journal of Animal Science*, **60**(4): 869-876.
- Bautista MLG, Ritulall AW & Wapnir RA (2003). Cord blood red cell osmotic fragility comparison between preterm and full-term new infants. *Early Human Development*, **72**(1): 37-46.
- Bazzoni GB, Bollini AN, Hernández GN, Contini MdelC & Rasia ML (2005). In vivo effect of aluminium upon the physical properties of

- the erythrocyte membrane. *Journal of Inorganic Biochemistry*, **99**(3): 822-827.
- Behling-Kelly E & Collins-Cronkright R (2014). Increases in beta-lipoproteins in hyperlipidemic and dyslipidemic dogs are associated with increased erythrocyte osmotic fragility. *Veterinary Clinical Pathology*, **43**(3): 405-415.
- Beutler E, Kuhl W & West C (1982). Osmotic fragility of erythrocytes after prolonged liquid storage and after reinfusion. *Blood*, **59**(6): 1141-1147.
- Bielawski J (1990). Two types of haemolytic activity of detergents. *Biochimica et Biophysica Acta*, **1035**(2): 214-217.
- Bilto YY (1990). Rheological action of aspirin on human erythrocytes. *Clinical Hemorrhology and Microcirculation*, **20**(3): 159-165.
- Bilto YY, Suboh S, Aburjai T & Abdalla S (2012). Structure-activity relationships regarding the antioxidant effects of the flavonoids on human erythrocytes. *Natural Science*, **4**(9): 740-747.
- Blasiak J, Walter Z & Gawronska M (1991). The changes of osmotic fragility of pig erythrocytes induced by organophosphorus insecticides. *Acta Biochimica Polonica*, **38**(1): 75-78.
- Bobhate SK, Gaikwad ST & Bhaledrao T (2002). NESTROFF as a screening test for detection of beta-thalassemia trait. *Indian Journal of Pathology and Microbiology*, **45**(3): 265-267.
- Boelsterli UA, Shie KP, Brändle E & Zbinden G (1983). Toxicological screening models: drug-induced oxidative hemolysis. *Toxicology Letters*, **15**(2-3): 153-158.
- Bogin E, Massry SG, Levi J, Djaldeti M, Bristol G & Smith J (1982). Effect of parathyroid hormone on osmotic fragility of human erythrocytes. *Journal of Clinical Investigation*, **69**(4): 1017-1025.
- Bondan C, Soares JC, Cecim M, Lopes ST, Graca DL & da Rocha RX (2005). Oxidative stress in the erythrocytes of cattle intoxicated with *Senecio sp.* *Veterinary Clinical Pathology*, **34**(4): 353-357.
- Bowdler AJ, Dougherty RM & Bowdler NC (1981). Age as a factor affecting erythrocyte osmotic fragility in males. *Gerontology*, **27**(4): 224-231.
- Buffenstein R, McCarron HC & Dawson TJ (2001). Erythrocyte osmotic fragility of red (*Macropus rufus*) and grey (*Macropus fuliginosus* and *Macropus giganteus*) kangaroos and free-ranging sheep of the arid regions of Australia. *Journal of Comparative Physiology B*, **171**(1): 41-47.
- Canonico PG, Kastello MD, Spears CT, Brown JR, Jackson EA & Jenkins DE (1984). Effects of ribavirin on red blood cells. *Toxicology and Applied Pharmacology*, **74**(2): 155-162.
- Cavalcanti TC, Gregorini CC, Guimaraes F, Rettori O & Vieira-Matos AN (2003). Changes in red blood cell osmotic fragility induced by total plasma and plasma fractions obtained from rats bearing progressive and regressive variants of the Walker 256 tumor. *Brazilian Journal of Medical and Biological Research*, **36**(7): 887-895.
- Chaudhary R & Katharia R (2012). Oxidative injury as contributory factor for red cells storage lesion during twenty eight days of storage. *Blood Transfusion*, **10**(1): 59-62.
- Chernitsky EA, Rozin W & Senkovich OA (2001). pH-dependence of detergent-induced hemolysis and vesiculation of erythrocytes. *Membrane and Cell Biology*, **14**(4): 529-536.
- Chernitsky EA & Senkovich OA (1997). Erythrocytes hemolysis by detergents. *Membrane and Cell Biology*, **11**(4): 475-485.
- Chernitsky EA & Senkovich OA (1998). Mechanisms of anionic detergent-induced hemolysis. *General Physiology and Biophysics*, **17**(3): 265-270.
- Chi LM & Wu WG (1991). Mechanisms of hemolysis of red cell mediated by ethanol. *Biochimica et Biophysica Acta*, **1062**(11): 46-50.
- Chikezie PC (2007). Osmotic fragility index of HbAA red blood cells in the presence of aqueous extracts of the three medicinal plants (*Aframomum melegueta*, *Garcinia kola* and *Cymbogon citratus*). *Global Journal of Pure and Applied Science*, **13**(4): 496-499.
- Chikezie PC & Uwakwe AA (2011). Membrane stability of sickle erythrocytes incubated in extracts of three medicinal plants: *Anacardium occidentale*, *Psidium guajava*, and *Terminalia catappa*. *Pharmacognosy Magazine*, **7**(26): 121-125.
- Chikezie PC, Uwakwe AA & Monago CC (2010). Comparative osmotic fragility of three erythrocyte genotypes (HbAA, HbAS and HbSS) of male participants administered

- with five anti malarial drugs. *African Journal of Biochemistry Research*, **4**(3): 57-64.
- Coldman NF, Gent M & Good W (1969). The osmotic fragility of mammalian erythrocytes in hypotonic solution of sodium chloride. *Comparative Biochemistry and Physiology*, **31**(4): 605-609.
- Colin-Negrete J, Kiesling HE, Ross TT & Smith JF (1996). Effect of the cottonseed on serum constituents, fragility of erythrocyte cells and reproduction of growing Holstein heifers. *Journal of Dairy Science*, **79**(11): 2016-2023.
- Costa DL, Albuquerque AC, Camacho ACLF, Costa FCH, Ferreira CAA, Caldas ACO, Almeida ACC, Cardoso MEO, Guimaraes-Silva S, Silva AH, Araujo RWN, Lima RC, Souza AS, Neto NCR, Borba HR, Dire GF & Bernardo-Filho M (2010). Studying of the biological effects of stannous chloride on the cell membrane. *New York Science Journal*, **3**(7): 70-77.
- Cribb PH, Olfert EA & Reynold FB (1986). Erythrocyte osmotic fragility testing and the prediction of canine malignant hyperthermia susceptibility. *Canadian Veterinary Journal*, **27**(12): 517-522.
- Cueff A, Seear R, Dyrda A, Bouyer G, Egee S, Esposito A, Skepper J, Tiffet T, Lew VL & Thomas SLY (2010). Effects of elevated intracellular calcium on the osmotic fragility of human red cells. *Cell Calcium*, **47**(1): 29-36.
- Dacie JV & Lewis SM (1995). *Practical Haematology*. Eight edition. Churchill Livingstone, New York. Pp 216-220.
- Dariyerli N, Toplan S, Akyolcu MC, Hatemi H & Yigit G (2004). Erythrocyte osmotic fragility and oxidative stress in experimental hypothyroidism. *Endocrinology*, **25**(1): 1-5.
- Das K & Bhattacharyya J (2010). Effect of calcium and diltiazem on phenylhydrazine-induced oxidative injury in goat erythrocytes. *Health*, **2**(10): 1221-1225.
- Dash BP & Kar BC (1999). Osmotic fragility of normal and sickle haemoglobin containing red blood cells. *Indian Journal of Physiology and Pharmacology*, **43**(2): 267-269.
- Davis JE & Brewer N (1935). Effect of physical training on blood volume, hemoglobin, alkali reserve and osmotic resistance of erythrocytes. *American Journal of Physiology*, **113**: 586-591.
- de Freitas MV, Netto RC, da Costa HJC, de Souza TM, Costa JO, Firmino CB & Penha-Silva N (2008). Influence of aqueous crude extracts of medicinal plants on the osmotic stability of human erythrocytes. *Toxicology In vitro*, **22**(1): 219-224.
- Debowy J, Obminska-Domoradzka B, Switala M & Garbulinski T (1985). The influence of copper-dextran complex (C-79), non-steroid anti-inflammatory drugs and bacterial pyrogen on stabilization of rabbit erythrocyte membrane. *Polish Journal of Pharmacology and Pharmacy*, **37**(1): 33-40.
- Demir S & Öner G (1995). The effects of cadmium on the fragility of red blood cell. *Journal of Islamic World Academy of Sciences*, **8**(2): 73-78.
- Devenuto F, Ligon DF, Friedrichsen DH & Wilson HL (1969). Human erythrocyte membrane uptake of progesterone and chemical alterations. *Biochimica et Biophysica Acta-Biomembranes*, **193**(1): 36-47.
- Dewey MJ, Mrown JL & Nallaseth FS (1982). Genetic differences in red cell osmotic fragility: analysis in allophonic mice. *Blood*, **59**(5): 986-989.
- Durotoye LA & Oyewale JO (1988): A comparative study of erythrocyte osmotic fragility between the guineafowl (*Numida meleagris galeata*) and the Nigerian domestic fowl (*Gallus domesticus*). *Bulletin of Animal Health and Production in Africa*, **36**(1): 73-76.
- Emembolu JO & Mba EC (1994). Red cell osmotic fragility in pregnant Nigerian women. *International Journal of Gynaecology and Obstetrics*, **44**(1): 73-74.
- Endoh Y, Kawakami M, Orringer EP & Meyer AA (1992). Causes and time course of acute haemolysis after burn injury in the rat. *Journal of Burn Care and Rehabilitation*, **13**(2Pt1): 203-209.
- Epps DE, Knechtel TJ, Baczynskyj O, Decker D, Guido DM, Buxser SE, Mathews WR, Buffenbarger SL, Lutzke BS, McCall JM, Oliver LK & Kezdy FJ (1994). Tirilazad mesylate protects stored erythrocytes against osmotic fragility. *Chemistry and Physics of Lipids*, **74**(2): 163-174.
- Etlik O, Tomur A, Tuncer M, Ridvanagaoglu AY & Andac O (1997). Protective effect of antioxidant vitamins on red cell lipoperoxidation induced by SO₂ inhalation. *Journal of Basic Clinical Physiology and Pharmacology*, **8**(1-2): 31-43.

- Fairley NM, Price GS & Meuten DJ (1988). Evaluation of red blood cell fragility in pygmy goats. *American Journal of Veterinary Research*, **49**(9): 1598-1600.
- Fansamade AA (1999). Erythrocyte osmotic fragility in hypertension and during diuretic therapy. *West African Journal of Medicine*, **18**(3): 183-186.
- Ferrali M, Signorini C, Caciotti B, Sugherini L, Ciccoli L, Giachetti D & Comporti M (1997). Protection against oxidative damage of erythrocyte membrane by the flavonoid quercetin and its relation to iron chelating activity. *Federation of European Biochemical Sciences Letters*, **416**(2): 123-129.
- Freeman AR & Spirtes MA (1963). Effects of chlorpromazine on biological membranes-II.: Chlorpromazine-induced changes in human erythrocytes. *Biochemical Pharmacology*, **12**(1): 47-53.
- Fukumoto K, Karai I & Horiguchi S (1983). Effect of lead on erythrocyte membranes. *British Journal of Industrial Medicine*, **40**(2): 220-223.
- Gaehagens P & Benner K (1974). Osmotic behaviour of human red blood cells effect of non-ionic detergents. *Blut*. **29**(2): 123-133. Name in full.
- Gomes de Sou JR, Pereira LP, da Costa Gomes RB, Pereira LD & Santos-Filho SD (2007). Potential pitfalls on the physiological (osmotic fragility) properties of the red blood cells: Action of a homeopathic medicine. *Pharmacognosy Magazine*, **3**(11): 182-186.
- Gueorguiev IP, Gueorguieva TM, Paskalev MD & Trifonova KY (1999). Erythrocytic osmotic fragility in stress-susceptible and stress-resistant pigs. *Revue de Medecine Veterinaire*, **1**(150): 39-42.
- Gutteridge JMC, Quinlan GJ, Clark I & Halliwell B (1995). Aluminium salts accelerate peroxidation of membrane lipids stimulated by iron salts. *Biochimica et Biophysica Acta. (BBA)-Lipid Lipid Metabolism*, **835**(3): 441-447.
- Habibu B, Kawu MU, Makun HJ, Aluwong T, Yaqub LS, Ahmad MS, Tauheed M & Buhari HU (2014). Influence of sex, reproductive status and foetal number on erythrocyte osmotic fragility, haematological and physiological parameters in goats during the hot-dry season. *Veterinary Medicine-Czech*, **59**(10): 479-490.
- Habibu B, Kawu MU, Makun HJ, Buhari HU & Hussaini M (2016). Breed and seasonal variations in erythrocyte fragility of goat kids raised in semi-arid savannah. *Comparative Clinical Pathology*, **25**(6): 1309-1312.
- Habibu B, Yaqub LS, Ahmad IA, Kawu MU, Buhari HU, Tauheed M & Isa HI (2013). Erythrocyte osmotic fragility and haematologic parameters of three breeds of 9-week old broiler chickens. *International Journal of Poultry Science*, **12**(5): 277-279.
- Hagve TA, Johansen Y & Christophersen B (1991) The effect of n-3 fatty acids on osmotic fragility of rat erythrocytes. *Biochimica et Biophysica Acta*, **1084**(3): 251-254.
- Hagve TA, Lie O & Grønn M (1993). The effect of dietary N-3 fatty acids on osmotic fragility and membrane fluidity of human erythrocytes. *Scandinavian Journal of Clinical and Laboratory Investigation Supplementum*, **53**(215): 75-84.
- Hanzawa K (2000). Changes in osmotic fragility of erythrocytes during exercise in athletic horses. *Journal of Equine Science*, **11**(3): 51-61.
- Hanzawa K, Kubo K, Hiraga A & Watanbe S (1996). Correlation between field exercise intensity and osmotic fragility of erythrocytes in thorough breed horses. In: *The Proceedings of the Eight Animal Science Congress of the Asia-Australasian Association of animal Production Societies*. Chiba, Japan. **2**: 574-575.
- Hanzawa K, Kubo K, Kai M, Hiraga A & Watanbe S (1998). Effects of splenic erythrocytes and blood lactate levels on osmotic fragility of circulating red blood cells in thoroughbred horses during exercise. *Journal of Equine Science*, **9**(4): 107-112.
- Hanzawa K, Kubo K, Kai M, Hiraga A & Watanbe S (1999). Effects of splenectomy for osmotic fragility of circulating red cells in thoroughbred horses during exercise. *Japan Journal of Equine Science*, **10**(3-4): 61-65.
- Harris JW, Brewster HH, Ham TH & Castle WB (1956). Studies on the destruction of red blood cells. X. The biophysics and biology of sickle cell disease. *Archives of Internal Medicine*, **97**(2): 145-168.

- Herman R (1969). Osmotic fragility of normal duck erythrocytes as influenced by extracts of *Plasmodium lophurae*, P. lophurae-infected cells and plasma. *Journal of Parasitology*, **55**(3): 626-632.
- Hernández G, Bollini A, Huarte M, Bazzoni G, Piehl L, Chiarotto M, Rubin de Celis E & Rasia M (2008). *In vitro* effect of aluminium upon erythrocyte membrane properties. *Clinical Hemorheology and Microcirculation*, **40**(3): 191-205.
- Hong-Wei T, Jiu-Li Z, Yin-Xia X & Shi-wen X (2008). The contribution of copper stress in erythrocytic osmotic fragility and haemolysis speed of goat. *Chinese Journal of Veterinary Science*, **1**:87-89. **Issue no?**
- Ibanga IA, Usoro CA & Nsonwu AC (2005). Glycaemic control in type 2 diabetics and the mean corpuscular fragility. *Nigerian Journal of Medicine*, **14**(3):304-306.
- Igbokwe NA (2016). Characterization of the osmotic stability of Sahel goat erythrocytes in ionic and non-ionic hypotonic media, PhD Thesis, Department of Physiology, Pharmacology and Biochemistry, Faculty of Veterinary Medicine, University of Maiduguri, Nigeria. Pp 1-219.
- Igbokwe NA & Igbokwe IO (2015). Influence of extracellular media's ionic strength on the osmotic stability of Sahel goat erythrocytes. *Journal of Basic and Clinical Physiology and Pharmacology*, **26**(2): 171-179.
- Igbokwe NA & Igbokwe IO (2016a). Phenotypic variations in osmotic lysis of Sahel goat erythrocytes in non-ionic glucose media. *Journal of Basic and Clinical Physiology and Pharmacology*, **27**(2): 147-154.
- Igbokwe NA & Igbokwe IO (2016b). Phenotypic homogeneity with minor deviance in osmotic fragility of Sahel goat erythrocytes in non-ionic sucrose media during various physiologic states. *Journal of Basic and Clinical Physiology and Pharmacology*, **27**(6): 633-641.
- Igbokwe NA, Ojo NA & Igbokwe IO (2015a). Quantified effects of late pregnancy and lactation on the osmotic stability of Sahel goat erythrocytes. *Nigerian Veterinary Journal*, **32**(1): 1122-1129.
- Igbokwe NA, Ojo NA & Igbokwe IO (2015b). Phenotypic drift in osmotic fragility of Sahel goat erythrocytes associated with variability of median fragility. *Sokoto Journal of Veterinary Sciences*, **13**(2): 6-13.
- Igbokwe NA, Ojo NA & Igbokwe IO (2016). Effects of sex and age on the osmotic stability of Sahel goat erythrocytes. *Comparative Clinical Pathology*, **25**(1): 15-22.
- Igbokwe NA, Sandabe UK, Bokko BP & Igbokwe IO (2018). Inhibition of osmotic permeability of caprine erythrocytes by mercuric chloride in osmotic fragility models. *Sokoto Journal of Veterinary Sciences*, **16**(3):24-32.
- Ikejiani O (1946). Studies in trypanosomiasis. IV. The fragility of erythrocytes in rats during the course of infection with *Trypanosoma lewesi*, *Trypanosoma brucei* and *Trypanosoma equiperdum*. *Journal of Parasitology*, **32**(4): 383-386.
- Islah L, Rita B, Youssef C, Abdarrahmane B, Abdarrahman H & Mohammed EK (2016). Study of incubation conditions for erythrocytes osmotic fragility testing in Dromedary camel (*Camelus dromedarius*). *International Journal of Research in Environmental Science*, **2**(2): 22-32.
- Jacob MH & Parpart AK (1931). Osmotic properties of erythrocytes. II. The influence of pH, temperature and oxygen tension on haemolysis by hypotonic solutions. *Bulletin*, **60**(2): 95-119.
- Jain NC (1973). Osmotic fragility of erythrocytes of dogs and cats in health and in certain hematologic disorders. *Cornell Veterinarian*, **63**(3): 411-413.
- Jain SK (1989). Hyperglycemia can cause membrane lipid peroxidation and osmotic fragility in human red blood cells. *Journal of Biological Chemistry*, **264**(35): 21340-21345.
- Jain SK & Williams DM (1988). Copper deficiency anemia: altered red blood cell lipids and viscosity in rats. *American Journal of Clinical Nutrition*, **48**(3): 637-640.
- Kafka M & Yerimiah T. (1998). The effect of EDTA as an anticoagulant on the osmotic fragility of erythrocytes. *Clinical and Laboratory Haematology*, **20**(4): 213-216.
- Kanias T, Lanteri M, Dereck S, Busch M & Gladwin MT (2013). Red blood cell storage in pediatric transfer bags is correlated with increased level of haemolysis and altered osmotic fragility. *Blood*, **122**(21): 2403-2404.
- Kaplay SS (1978). Lack of association between human erythrocyte membrane

- acetylcholinesterase and osmotic fragility. *Biochemia Medica*, 19(3): 299-304.
- Karai I, Fukumoto K & Horiguchi S (1979). Effects of heavy metal ions on osmotic resistance of human red cells. *Osaka City Medical Journal*, 25(2): 153-157.
- Karai I, Fukumoto K & Horiguchi S (1981). Studies on osmotic fragility of red blood cells determined with a coil plant centrifuge for workers occupationally exposed to lead. *International Archives of Occupational and Environmental Health*, 48(3): 273-281.
- Karai I, Fukumoto K & Horiguchi S (1982a). Relationships between osmotic fragility of red blood cells and various hematologic data in workers exposed to lead. *International Archives of Occupational and Environmental Health*, 50(1): 17-23.
- Karai I, Fukumoto K, Kageyama K & Horiguchi S (1982b). Effect of lead *in vitro* on water metabolism and osmotic fragility of human erythrocytes. *British Journal of Industrial Medicine*, 39(3): 295-299.
- Karuppasamy R, Subathra S & Puvanewari S (2005). Hematological responses to exposure to sublethal concentration of cadmium in air breathing fish, *Channa punctatus*. *Journal of Environmental Biology*, 26(1): 123-126.
- Kaya H & Saito T (1985). Effect of progesterone and its 17 alpha hydroxyl derivative on human erythrocyte membrane. *Japan Journal of Pharmacology*, 39(3): 299-306.
- Kim HD, Luthra MG, Watts RP & Stern LZ (1980). Factors influencing osmotic fragility of red blood cells in Duchenne muscular dystrophy. *Neurology*, 30(7Pt1): 726-731.
- Kirchgessner M, Stangl GI, Reichlmayr-Lais AM & Eder K (1994). The effects of dietary oils on the fatty acid composition and osmotic fragility of rat erythrocytes. *Z. Ernährungswiss*, 33(2): 146-158.
- Kiron V, Takeuch T & Watanabe T (1994). The osmotic fragility of erythrocytes in rainbow-trout under different dietary fatty-acid status. *Fisheries Science*, 60(1): 93-95.
- Kobo PI, Ayo JO, Aluwong T, Zezi AU, Maikai V & Ambali SF (2014). Flavonoid mixture ameliorates increase in erythrocyte osmotic fragility and malondialdehyde concentration induced by Trypanosoma brucei brucei-infection in Wistar rats. *Research in Veterinary Science*, 96(1): 139-142.
- Kogawa H, Yabushita N, Nakajima T & Kageyama K (1998). Studies on *in vitro* effect of free fatty acids on water content and osmotic fragility of rabbit (*Lepus cuniculus*) erythrocytes. *Life Science*, 62(9): 823-828.
- Kohn B, Goldschmidt MH, Hohenhaus AE & Giger U (2000). Anemia, splenomegaly and increased osmotic fragility of erythrocytes in Abyssinian and Somali cats. *Journal of American Veterinary Medical Association*, 217(10): 1483-1491.
- Kolanjippan K, Manohoran S & Kayalvizhi M (2002). Measurement of erythrocyte lipids, lipid peroxidation, antioxidants and osmotic fragility in cervical cancer patients. *Clinica Chimica Acta*, 326(1-2): 143-149.
- Korones D & Pearson HA (1988). Normal erythrocyte osmotic fragility in hereditary spherocytosis. *Journal of Pediatrics*, 114(2): 264-266.
- Koumanov KS, Tessier C, Momchilova AB, Rainteau D, Wolf C & Quinn PJ (2005). Comparative lipid analysis of detergent-resistant membrane raft fractions isolated from human and ruminant erythrocytes. *Archives of Biochemistry and Biophysics*, 434(1): 150-158.
- Kowluru RA, Kern TS, Engerman RL & Armstrong D (1996). Abnormalities of Retinal Metabolism in Diabetes or Experimental Galactosemia. III. Effects of Antioxidants. *Diabetes*, 45(9): 1233-1237.
- Kraus A, Roth HP & Kirchgessner M (1997). Supplementation with vitamin C, vitamin E or β -carotene influences osmotic fragility and oxidative damage of erythrocytes of zinc-deficient rats. *Journal of Nutrition*, 127(7): 1290-1296.
- Krogmeier DE, Mao IL & Bergen WG (1993). Genetic and non-genetic effects on erythrocyte osmotic fragility in lactating Holstein cows and its association with yield traits. *Journal of Dairy Science*, 76(7): 1994-2000.
- Kual P, Sidhu H, Thind SK, Sharma SK & Nath R (1995). Vitamin B₆ deficiency and galactose induced alterations in morphology and osmotic fragility of rat erythrocyte. *Scanning Microscopy*, 9(4): 1127-1135.
- Kuiper PJC, Avinoam L & Meyerstein N (1971). Changes in lipid composition and osmotic fragility of erythrocytes of hamster induced by heat exposure. *Biochimica et Biophysica*

- Acta (BBA) Lipid Lipid Metabolism*, **248**(2): 300-305.
- Kung CM, Tseng ZL & Wang HL (2009). Erythrocyte fragility increases with level of glycosylated hemoglobin in type 2 diabetic patients. *Clinical Hemorheology and Microcirculation*, **43**(4): 345-351.
- Kwant WO & Van Steveninck J (1968). The influence of chlorpromazine on human erythrocytes. *Biochemical Pharmacology*, **17**(10): 2215-2218.
- Legrange SN, Breitschwerdt EB, Grindem CB & Beutler E (1995). Erythrocyte fragility and chronic intermittent pigmenturia in a dog. *Journal of American Veterinary Medical Association*, **206**(7): 1002-1006.
- Lemos GS, Marquez-Bernardes LF, Arvelos LR, Paraiso LF & Penha-Silva N (2011). Influence of glucose concentration on the membrane stability of human erythrocytes. *Cell Biochemistry and Biophysics*, **61**(3): 531-537.
- Lenfant F, Lahet JJ, Vergely C, Volot C, Freysz M & Rochette L (2000). Lidocaine inhibits potassium efflux and hemolysis in erythrocytes during oxidative stress *in vitro*. *General Pharmacology:Vascular System*, **34**(3): 193-199.
- Lessler MA & Walters MI (1973). Erythrocyte osmotic fragility in the presence of lead or mercury. *Proceedings for the Society for Experimental Biology and Medicine*, **142**(2): 548-553.
- Levander OA, Ferretti RJ & Morris VC (1977). Osmotic and peroxidative fragilities of erythrocytes from vitamin E deficient lead-poisoned rats. *Journal of Nutrition*, **107**(3): 373-377.
- Lewis JH & Ferguson EE (1966). Osmotic fragility of premammalian erythrocytes. *Comparative Biochemistry and Physiology*, **18**(3): 589-595.
- Leylek OA, Egur A, Senocak F, Sencan M, Bakir S, Ozdemir H & Songur S (1998). Prophylaxis of the occurrence of hyperbilirubinemia in relation to maternal oxytocin infusion with steroid treatment. *Gynecologic and Obstetric Investigation*, **46**(3): 164-168.
- Livne A, Kuiper PJC & Mevestein N (1972). Differential effects of lipids on the osmotic fragility of hamster erythrocytes. *Biochemica et Biophysica Acta (BBA)-Biomembranes*, **255**(3): 744-750.
- Mafudvaze B & Erlwanger KH (2007). The effect of EDTA, heparin and storage on the erythrocyte osmotic plasma osmolality and haematocrit of adult ostriches (*Struthio camelus*). *Veterinary Archives*, **77**(5): 427-434.
- Mafuvadze B, Nyanungo M, Saina H, Gorejena B, Mashayamombe T & Erlwanger KH (2008). Deprivation of drinking water for up to 48 hours does not affect the osmotic fragility of erythrocytes from captive helmeted guinea fowl (*Numida meleagris*). *International Journal of Poultry Science*, **7**(1): 59-63.
- Magid MS, Perlin M & Gottfried EL (1982). Increased erythrocyte osmotic fragility in pregnancy. *American Journal of Obstetrics and Gynecology*, **144**(8): 910-914.
- Makinde MO (1986) Osmotic fragility of erythrocytes in two breeds of swine. *Animal Technology*, **37**:73-76.
- Makinde MO & Bobade PA (1994). Osmotic fragility of erythrocytes in clinically normal dogs and dogs infected with parasites. *Research in Veterinary Science*, **53**(3): 343-348.
- Makni M, Chtourou Y, Fetoui H, Garoui EM, Barkallah M, Marouani C, Kallel C & Zeghal N (2012). Erythrocyte oxidative damage in rat treated with CCl₄: protective role of vanillin. *Toxicology and Industrial Health*, **28**(10): 908-916.
- Malachi T, Bogin E, Gafter U & Levi J (1986): Parathyroid hormone effect on the fragility of human young and old red blood cells in uremia. *Nephron*, **42**(1): 52-57.
- Manju M, Anil J, Kishor D, Vinky R & Hemant K (2006). Bias-corrected diagnostic performance of the naked eye single-tube red cell osmotic fragility test (NESTROFT): an effective screening tool for beta-thalassemia. *Hematology*, **11**(4): 277-286.
- Marar T (2011). Amelioration of glucose induced hemolysis of human erythrocytes by vitamin E. *Chemico- Biological Interactions*, **193**(2): 149-153.
- March BE, Coates V & Biely J (1966). The effects of estrogen and androgen on osmotic fragility and fatty acid composition of erythrocyte in the chicken. *Canadian Journal of Physiology and Pharmacology*, **44**(3): 379-387.
- Marques MLD, Cazana FJL & Puyol MR (1986). *In vitro* response of erythrocytes to alpha-tocopherol exposure. *International Journal*

- of Vitamin and Nutrition Research, **56**(3): 311-315.
- Martins AP, Maron A, Ciancetta A, Cobo AG, Echevarri M, Moura TF, Re N, Casini A & Soveral G (2012). Targeting aquaporin function; potent inhibition of an aquaglyceroporin-3 by a gold based compound. *Public Library of Science*, **7**(5):e37435.
- Matondi GHM, Masam E, Mpofu IDT & Muronzi FF (2007). Effect of feeding graded levels of cottonseed meal on goat erythrocyte membrane osmotic fragility. *Livestock Research for Rural Development*, **19**(11): 169-172.
- Matsuzawa T & Ikarashi Y (1979). Haemolysis of various mammalian erythrocytes in sodium chloride, glucose and phosphate-buffer solutions. *Laboratory Animals*, **13**(4): 329-331.
- Mel HC & Reed TA (1981). Biophysical responses of red cell-membrane systems to very low concentrations of inorganic mercury. *Cell Biochemistry and Biophysics*, **3**(3): 233-250.
- Mena H, Santos JE, Huber JT, Tarazon M & Calhoan MC (2004). The effects of varying gossypol intake from whole cottonseed and cottonseed meal on lactating dairy cows. *Journal of Dairy Science*, **87**(8): 2506-2518.
- Menges WL (1991). Effects of feeding cotton seed on mohair growth and performance of Angora goats. *MSc Thesis*. Angelo State University, San Angelo, USA. Pp 82-84.
- Mijares A, Vivas J, Abad C, Betancourt M, Pinero S, Proverbio F, Maria R & Portillo R (2010). *Trypanosoma evansi*: Effect of experimental infection on the osmotic fragility, lipid peroxidation and calcium ATPase activity of rat red blood cells. *Experimental Parasitology*, **124**(3): 301-305.
- Mineo H & Hara H (2005). Structure-dependent and receptor-independent increase in osmotic fragility of rat erythrocytes by short-chain fatty acids. *Biochimica et Biophysica Acta-Biomembranes*, **1713**(2): 113-117.
- Minka NS & Ayo JO (2010). Physiological responses of erythrocytes of goats to transportation and the modulatory role of Ascorbic acid. *Journal of Veterinary Medical Science*, **72**(7): 875-881.
- Monteiro MO, Pinto M, Marin PJ, Santos-Filho SD & Bernardo-Filho M (2013). Effects of osmotic fragility of red blood cells of whole blood submitted to vibrations in an oscillating platform. *African Journal of Biotechnology*, **10**(64): 14197-14202.
- Mosior M & Gomulkiewicz J (1985). Effect of phosphate ions on osmotic properties of human and bovine erythrocytes: a relation between the state of glycolysis and critical cell volume. *Studia Biophysica*, **107**(3):169-178.
- Mosior M & Gomulkiewicz J (1988). Osmotic properties of bovine erythrocytes aged in vivo. *General Physiology and Biophysics*, **7**(1): 73-79.
- Mostafavi E, Nakhjavani M, Ghazizadeh Z, Barakati H, Mirmiranpour H & Safari R (2013). Protective role of calcium ion against stress-induced osmotic fragility of red blood cells in patients with type 2 diabetes mellitus. *Clinical Hemorheology and Microcirculation*, **53**(3): 239-245.
- Mrugesh T, Dipa L & Manishika G (2011). Effect of lead on human erythrocytes: an *in vitro* study. *Acta Poloniae Pharmaceutica Drug Research*, **68**(5): 653-656.
- Murphy JR (1967). The influence of pH and temperature on some physical properties of normal erythrocytes and erythrocytes from patients with hereditary spherocytosis. *Journal of Laboratory and Clinical Medicine*, **69**:758-775.
- Mutoh S, Sasaki R, Takaku F, Aoyama M, Moriyama S, Yoshimoto M & Yawata Y (1983). A family of hereditary stomatocytosis associated with normal level of Na⁺-K⁺-ATPase activity of red blood cells. *American Journal of Hematology*, **14**(2): 113-120.
- Nakamura Y (1983). Erythrocyte osmotic resistance in pregnancy. *American Journal of Obstetrics and Gynecology*, **147**(4): 472-473.
- Nirmalan G, Nair SG & Simon KJ (1967). Hematology of the Indian elephant (*Elephas maximus*). *Canadian Journal of Physiology and Pharmacology*, **45**(6): 985-991.
- Norman NK & Dewey MJ (1985). Genetic control of red cell osmotic fragility. *Journal of Heredity*, **76**(1): 31-35.
- O'Dell BL, Browning JD & Reves PG (1987). Zinc deficiency increases the osmotic fragility of rat erythrocytes. *Journal of Nutrition*, **117**(11): 1883-1889.
- Ogunro PS, Ogungbamigbe TO & Muhibi MA (2010). The influence of storage period on the

- antioxidants level of red blood cells and the plasma before transfusion. *African Journal of Medical Science*, **39**(2): 99-104.
- Ogunyemi A D & Olayemi FO (2016). Comparative assesment of erythrocyte osmotic fragility in two breeds of dog: Nigerian indigenouse breed and the German Shepherd. *Comparative Clinical Pathology*, **25**(1): 79-83.
- Okuda J & Tsuzuki Y (1977). Changes of some properties of blood of rats administered with methylmercuric chloride. *Chemical and Pharmaceutical Bullentin*, **25**(2): 209-214.
- Okwusidi JI (2002). Forty-eight hour storage alters human erythrocyte osmotic fragility in Nigerian males. *Tropical Journal of Health Science* **9**(1): 4-7.
- Okwusidi JI (2004). Moderation of osmotic fragility in non-detergent washed human erythrocytes. *Tropical Journal of Health Science*, **11**(1): 37-40.
- Okwusidi JI (2011). Osmotic fragility in stored non-detergent washed human erythrocytes. *African Journal of Biomedical Research*, **14**(2): 143-146.
- Oladele SB, Ayo JO, Ogundipe SO & Esievo KAN (2003). Seasonal and species variations in erythrocytes osmotic fragility of indigenouse poultry species in Zaria, Northern Guinea Savannah zone of Nigeria. *Bullentin of Animal Health and Production in Africa*, **51**(3): 204-214.
- Olaifa AK, Onwuka SK, Ariyibi A & Craig OO (1999). Futher studies in canine erythrocyte fragility in the tropical sub region: The effects of sodium pentobarbitone aneesthesia in Nigerian dogs. *African Journal of Biomedical Research*, **2**(1): 29-31.
- Olaifa F, Ayo JO, Ambali SF & Rekwot PI (2012). Effect of packing on changes in erythrocyte osmotic fragility and malondialdehyde concentration in donkeys administered with ascorbic acid. *Onderstepoort Journal of Veterinaty Research*, **79**(1) doi.org/10.4102/ojvr.v79i1.413 .
- Olayemi FO (2004). Erythrocyte osmotic fragility, haematological and biochemical parameters of the Nigerian White Fulani cattle. *Bullentin of Animal Health and Production in Africa*, **52**(3): 208-211.
- Olayemi FO (2007). The effect of sex on the erythrocyte osmotic fragility of the Nigerian White Fulani and Ndama breeds of cattle. *Tropical Veterinarian*, **25**(3): 106-111.
- Olayemi FO, Azeez IO, Ogunyemi A & Ighagbon, F.O. (2009). Study on erythrocyte values of the Nigerian indigenouse dog. *Folia Veteterinaria*, **53**(2): 65—67.
- Olayemi FO & Oyewale JO (2002). Comparative assesment of the erythrocyte osmotic fragility and of haematological and plasma biochemical values in the NigerianWhite Fulani and N'dama breeds of cattle. *Tropical Animal Health and Production*, **34**(3): 181-187.
- Olorunshola KV, Eze KO & Achie LN (2012). Effect of road transportation stress on erythrocyte fragility (EOF) of healthy young adult Nigerians during the harmattan season. *Nigerian Journal of Physiological Science*, **27**(2): 157-164.
- Olowookorun MO & Makinde MO (1998). Comparative assessment of erythrocyte osmotic fragility, haematological and serum biochemical values in the domestic chicken and ostrich. *Tropical Veterinarian*, **16**:1-7.
- Olusanya SK & Adepoju FO (1979). Osmotic fragility of erythrocyte of some Nigerian species of domestic ruminants. *OAU/STRC Bullentin of Animal Health and Production in Africa*, **27**(4): 237-243.
- Orbach A, Zelig O, Yedgar S & Barshtein G (2017). Biophysical and biochemical markers of red blood cell fragility. *Transfusion Medicine and Hemotherapy*, **44**(3): 183-187
- Oyededeji KO, Abayomi O, & Dele A (2015b). Effects of ethanol extract of *Adenopus breviflorus* (Benth) on erythrocyte osmotic fragility in male Wistar rats. *International Journal of Pharmaceutical Sciences Review and Research*, **35**(1): 152-154.
- Oyededeji KO & Alabi AS (2015). Effects of lead acetate on erythrocyte osmotic fragility in male Wistar rats. *International Journal of Pharmaceutical Sciences Review and Research*, **35**(1): 158-160.
- Oyededeji KO, Deikola A & Abayomi O (2015a). Effects of ethanol extract of *Jatropha gossypifolia* (Pohl) on erythrocyte osmotic fragility in male Wistar rats. *International Journal of Pharmaceutical Sciences Review and Research*, **35**(1): 155-157.
- Oyewale JO (1987). Studies on the erythrocyte osmotic fragility of rats infected with

- Trypanosoma brucei*. *Animal Technology*, **38**(10): 219-228.
- Oyewale JO (1990). The effect of egg-laying on the osmotic fragility of erythrocytes, leucocytes and blood volume in the guinea-hen. *Animal Technology*, **41**(8): 59-64.
- Oyewale JO (1991a). Osmotic fragility of erythrocytes of West African Dwarf sheep and goats: Effects of temperature and pH. *British Veterinary Journal*, **147**(2): 163-170.
- Oyewale JO (1991b). Osmotic fragility of erythrocytes of guinea-fowls at 21 and 156 weeks of age. *Veterinarski Arhiv*, **61**(1): 49-56.
- Oyewale JO (1992). Changes in osmotic resistance of erythrocytes of cattle, pigs, rats and rabbits during variation in temperature and pH. *Zentralbl Veterinarmed A*, **39**(2): 98-104.
- Oyewale JO (1993). Effect of storage of blood on the osmotic fragility of mammalian erythrocytes. *Zentralbl Veterinarmed A*, **40**(4): 258-264.
- Oyewale JO (1994a). Further studies on osmotic resistance of nucleated erythrocytes: observations with pigeon, pea fowls, lizard and toad erythrocytes during changes in temperature and pH. *Zentralbl Veterinarmed A*, **41**(6): 62-71.
- Oyewale JO (1994b). Changes in osmotic fragility of nucleated erythrocytes resulting from blood storage. *Zentralbl Veterinarmed A*, **41**(6): 475-479.
- Oyewale JO & Ajibade HA (1990). The osmotic fragility of erythrocytes of turkeys of two age groups. *Veterinarski Arhiv*, **60**(1):43-48.
- Oyewale JO & Durotoye LA (1988). Osmotic fragility of erythrocytes of two breeds of domestic fowl in the warm humid tropics. *Laboratory Animals*, **22**(3): 250-254.
- Oyewale JO, Dzenda T, Yaqub L, Akanbi D, Ayo J, Owoyele O, Minka N & Dare T (2011). Alterations in the osmotic fragility of camel and donkey erythrocytes caused by temperature, pH and blood storage. *Veterinarski Arhiv*, **81**(4): 459-470.
- Oyewale JO, Olayemi FO & Oke OA (1998b). Haematology of the wild adult African giant rat (*Cricetomys gambianus*, waterhouse). *Veterinarski Arhiv*, **68**(2): 91-99.
- Oyewale JO, Olayemi FO & Rahman SA (1998a). Blood characteristics of the Nigerian local duck (*Anas platyrhynchos*). I. Red blood cell characteristics. *Veterinarski Arhiv*, **68**(6): 199-204.
- Oyewale JO, Sanni AA & Ajibade HA (1991). Effects of temperature, pH and blood storage on osmotic fragility of duck erythrocytes. *Journal of Veterinary Medicine Series A*, **38**(1-10): 261-264.
- Oztürk B & Ozdemir S (2015). Effects of aluminum chloride on some trace elements and erythrocyte osmotic fragility in rats. *Toxicology and Industrial Health*, **31**(12): 1069-1077.
- Paes G, Paepe D, Meyer E, Kristensen AT, Duchateau L, Campos M & Daminet S (2013). The use of the rapid osmotic fragility test as an additional test to diagnose canine immune-mediated haemolytic anaemia. *Acta Veterinaria Scandinavica*, **55**(1): 74.
- Pati S, Panda SK, Behera PC & Panda MR (2017). Assessment of erythrocyte osmotic fragility in cattle due to haemoprotozoan diseases. *International Journal of Science, Environment and Technology*, **6**(2): 1560-1568.
- Peinado VI, Alfaro V, Palomeque J, Palacios L & Viscor G (1993). Erythrocyte osmotic resistance during acute hypothermia in awake unrestrained rats. *Pflügers Archiv*, **424**(5-6): 555-557.
- Penha-Silva N, Firmino BC, Reis FGD, Huss JCD, Souza TMT, de Freitas MV & Netto RdM. (2007). Influence of age on the stability of human erythrocyte membranes. *Mechanisms of Ageing and Development*, **128**(7-8): 444-449.
- Perk K, Frei YF & Herz A (1964a). Osmotic fragility of red blood cells of young and mature domestic and laboratory animals. *American Journal of Veterinary Research*, **25**:1241-1248.
- Perk K, Hort I & Perri A (1964b). The degree of swelling and osmotic resistance in hypotonic solutions of erythrocytes from various domestic animals. *Refuah Veterinaria*, **20**:122-124.
- Pieragostini E, Petazzi F & Luccia AD (2003). The relationship between the presence of extra α -globin gene and blood cell traits in Altamurana sheep. *Genetic Selection Evolution*, **35** (Suppl. 1): S121-S133.
- Pillai, S.R., Steiss, J.E., Traber, M.G., Kayden, H.J. and Wright, J.C. (1992). Comparison of four erythrocyte fragility tests as indicators of

- vitamin E status in adult dogs. *Journal of Comparative Pathology*, **107**(4): 399-410.
- Prabhakar S, Nanda AS & Ghuman SPS (2013). Evaluation of osmotic fragility of erythrocytes in dystocia affected buffaloes. *Indian Journal of Animal Sciences*, **70**(5): 452-454.
- Price GS, Armstrong PJ, McLeod DA, Babineau CA, Metcalf MK & Sellett LC (1988). Evaluation of citrate-phosphate-dextrose-adenine as a storage medium for packed canine erythrocytes. *Journal of Veterinary Internal Medicine*, **2**(3): 126-132.
- Qazi QH, Go SC, Smithwick EM & Madahar DP (1972). Osmotic resistance of abnormal red cells exposed to lead *in vitro*. *British Journal of Haematology*, **23**(5): 631-633.
- Ramana DC, Prasad MH, Reddy TP & Reddy PP (1997). Glycosylation of hemoglobin and erythrocyte membrane proteins mediated changes in osmotic fragility of erythrocytes. *Indian Journal of Medical Science*, **51**(1): 5-9.
- Ranasinghe P, Kaushalya WP, Abeysekera M, Premakumara GAS, Perera YS, Gurugama P & Gunatilake SB (2012). *In vitro* erythrocyte membrane stabilization properties of *Carica papaya* L. leaf extracts. *Pharmacognosy Research*, **4**(4): 196–202.
- Rand PW, Lacombe E, Barker ND & Kallechey GL (1997). Effects of methylprednisolone on the physical properties of human red cell. *Journal of Laboratory Clinical Medicine*, **89**(6): 1241-1250.
- Raz A & Livne A (1973). Differential effects of lipids on the osmotic fragility of erythrocytes. *Biochimica et Biophysica Acta. (BBA)-Biomembranes*, **311**(2): 222-229.
- Richards RS, Wang L & Jelinek H (2007). Erythrocytes oxidative damage in chronic fatigue syndrome. *Archives of Medical Research*, **38**(1): 94-98.
- Rifkind JM, Araki K & Hadley EC (1983). The relationship between the osmotic fragility of human erythrocytes and cell age. *Archives of Biochemistry and Biophysics*, **222**(2): 582-589.
- Roozbeh J, Sharifian M, Karimi M, Hamidian-Jahromi AR & Afshariani R (2009). Effect of zinc supplementation on red blood cell osmotic fragility in hemodialysis patients. *Shiraz E-Medical Journal*, **10**(4): 186-189.
- Salami HA, John AI & Ekanem AU (2012). The effect of aqueous preparation of *Allium cepa* (onion) and *Allium sativa* (garlic) on erythrocyte osmotic fragility in wistar rats: *in vivo* and *in vitro* studies. *Nigerian Journal of Physiology*, **27**(1): 29-34.
- Saleh MA, Al-Salahy MB & Sanousi SA (2009). Oxidative stress in blood of camels (*Camelus dromedarius*) naturally infected with *Trypanosoma evansi*. *Veterinary Parasitology*, **162**(3-4): 192-199.
- Santoro ML, Kogika MM, Hagiwara MK, Miranda RMS & Castela IICG (1994). Decreased erythrocyte osmotic fragility during canine leptospirosis. *Revista do Instituto de Medicina Tropical de São Paulo*, **36**(1): 1-5.
- Schaefer CA & Dewey MJ (1989). Single Locus (*rol*) Control of extreme resistance to red cell osmotic lysis: Intrinsic mode of gene action. *Genetics*, **121**(2): 353-358.
- Schalm OW, Jain NC & Carroll EJ (1975). *Veterinary Hematology*. Third edition. Lee and Febiger. Philadelphia. Pp 66-78.
- Seeman P, Kwant WO, Sauks T & Argent W (1969) Membrane expansion of intact erythrocytes by anaesthetics. *Biochimica et Biophysica Acta*, **183**(3): 490-498.
- Sekar PS & Selvam R (1994). Influence of glycemia on erythrocyte lipid-peroxidation, osmotic fragility and glutathione-dependent enzyme defense in human non-insulin-dependent diabetes-mellitus. *Medical Science Research*, **22**(7): 485-487.
- Senturk UK, Gunduz F, Kuru O, Aktekin MR & Kipmen D (2001). Exercise-induced oxidative stress affects erythrocytes in sedentary rats but not exercise-trained rats. *Journal of Applied Physiology*, **91**(5): 199-204.
- Sharma RL, Bhat TK & Dhar DN (1989). Effect of *Dictyococcus filaria* infection on the osmotic fragility of sheep erythrocytes. *Veterinary Parasitology*, **30**(3): 253-258.
- Sharma OP, Makkar HPS, Pal RN & Negi SS (1981). Fragility of erythrocytes in animals affected by *Lantana* poisoning. *Clinical Toxicology*, **18**(1): 25-35.
- Sharma B, Rai DK, Rai PK, Rizvi SI & Watal G (2010). Determination of erythrocyte fragility as a marker of pesticide-induced membrane oxidative damage. *Advanced Protocols in Oxidative Stress II: Methods. Molecular Biology*, **594**(1):123-128.

- Silva ID & Kuruwita VY (1994). The osmotic fragility of erythrocytes of the Asian elephant (*Elephas maximus*). *GAJAH: Journal of the IUCN/SSC Asian Elephant Specialist Group*, **13**(1): 25-29.
- Silva IM, Hubsch C & Ysern-Caldentey M (1989). Erythrocyte osmotic fragility and cation concentrations during experimentally induced bovine anaplasmosis. *Comparative Biochemistry and Physiology Part A: Physiology*, **94**(3): 455-459.
- Sirichotiyakul S, Tantipalakov C, Sanguanserm Sri T, Wanapirak C & Tongsong T (2004). Erythrocyte osmotic fragility test for screening of alpha-thalassemia-1 and beta-thalassemia trait in pregnancy. *International Journal of Gynecology and Obstetrics*. **86**(3): 347-50.
- Soares JJ, Folmer VV, DaRocha JB & Nogueira CW (2014). Ebselen exhibit glycation-inhibiting properties and protects against osmotic fragility of human erythrocytes *in vitro*. *Cell Biology International*, **38**(5): 625-630.
- Soliman MK & Amrousi SE (1966). Erythrocyte fragility of healthy fowl, dog, sheep, cattle, buffalo, horse and camel blood. *Veterinary Record*, **78**(12): 429-430.
- Sözmen EY, Tanyalçin T, Onat T, Kutay F & Erilaçin S (1994). Ethanol induced oxidative stress and membrane injury in rat erythrocyte. *European Journal of Clinical Chemistry and Clinical Biochemistry*, **32**(10): 741-744.
- Spengler MI, Leroux MB, Svetaz MJ, Contesti JF, Parente FM & Bertoluzzo SM (2007). Nifedipine effect on red cell rheological properties in patients with systemic scleroderma. *Clinical Hemorheology and Microcirculation*, **36**(2): 105-110.
- Srouf MA, Bilto YY & Juma M (2000a). Susceptibility of erythrocytes from non-insulin-dependent diabetes mellitus and hemodialysis patients, cigarette smokers and normal subjects to *in vitro* oxidative stress and loss of deformability. *Clinical Hemorheology and Microcirculation*, **22**(3): 173-180.
- Srouf MA, Bilto YY, Juma M & Irhimeh MR (2000b). Exposure of human erythrocytes to oxygen radical causes loss of deformability; increased osmotic fragility, lipid peroxidation and protein degradation. *Clinical Hemorheology and Microcirculation*, **23**(1): 13-21.
- Stevenson LM & Jones DG (1989). Relationships between vitamin E status and erythrocyte stability in sheep. *Journal of Comparative Pathology*, **100**(4): 359-368.
- Suboh SM, Bilto YY & Aburja TA (2004). Protective effects of selected medicinal plants against protein degradation, lipid peroxidation and deformability loss of oxidatively stressed human erythrocyte. *Phytotherapy Research*, **18**(4): 280-284.
- Suhail M & Ahmad I (1995). *In vitro* effects of acetaminophen on rat RBC and role of vitamin E. *Indian Journal of Experimental Biology*, **33**(4): 269-271.
- Suhail M, Patil S, Khan S & Siddiqui S (2010). Antioxidant Vitamins and Lipoperoxidation in Non-pregnant, Pregnant, and Gestational Diabetic Women: Erythrocytes Osmotic Fragility Profiles. *Journal of Clinical Medicine and Research*, **2**(6): 266-273.
- Tadano J, Niwa M, Ueda H & Shibuya M (1981). Study on erythrocyte lipids and osmotic fragility in elderly people with anemia. *Tokai Journal of Experimental and Clinical Medicine*, **6**(1): 35-40.
- Thomas-George N, Jayanthi B & Krishna-Murthy S (1979). Studies on hemolysis of human erythrocytes by linoleic acid. *Journal of Biosciences*, **1**(4): 385-392.
- Thool AA, Walde MS, Shrikhande AV & Talib VH (1998). A simple screening test for detection of heterozygous beta thalassemia. *Indian Journal of Pathology and Microbiology*, **41**(4): 423-426.
- Tritschler C, Mizukami K, Raj K & Giger U (2016). Increased erythrocyte osmotic fragility in anemic domestic shorthair and purebred cats. *Journal of Feline Medicine and Surgery*, **18**(6): 462-470.
- Troiano JC, Althaus R, Scaglione MC & Scaglione LM (2000). Osmotic fragility and size of erythrocytes in *Caiman latirostris* and *Caiman crocodylus jacare* (*Crocodylia-alligatoridae*) under captive conditions. *Comparative Haematology International*, **8**(1): 50-52.
- Tyulina OV, Prokopieva VD, Boldyrev AA & Johnson P (2002). *In vitro* effects of ethanol, acetaldehyde and fatty acid ethyl esters on human erythrocytes. *Alcohol and Alcoholism*, **37**(2): 179-186.
- Uchendu C, Ambali SF, Yakub SL, Lasisi OI & Umosen AJ (2011). Comparative effects of vitamin C

- and acetyl-L-carnitine on sub-acute chlorpyrifos-induced erythrocyte osmotic fragility in wistar rats. *Advances in Applied Science Research*, **2**(5): 297-302.
- Uzum A, Toprak O, Gumustas MK, Ciftci S & Sen S (2006). Effect of vitamin E therapy on oxidative stress and erythrocyte osmotic fragility in patients on peritoneal dialysis and hemodialysis. *Journal of Nephrology*, **19**(6): 739-745.
- Van de Watering L (2011). Red cell storage and prognosis. *Vox Sanguinis*, **100**(1): 36-45.
- Van der Walt JH & Russell WJ (1978). Effect of heating on the osmotic fragility of stored blood. *British Journal of Anaesthesia*, **50**(8): 815-820.
- Veale MF, Healey G & Sparrow RL (2011). Effect of additive solutions on red blood cell (RBC) membrane properties of stored RBCs prepared from whole blood held for 24hrs at room temperature. *Transfusion*, **51**(supplement): 25-33.
- Vido AA, Cavalcanti TC, Guimareães F, Vieira-Matos A & Rettori O (2000). The hemolytic component of cancer anemia: Effects of osmotic and metabolic stress on the erythrocytes of rats bearing multifocal inoculations of the Walker 256 tumor. *Brazilian Journal of Medical and Biological Research*, **33**(7): 815-822.
- Viscor G & Palomeque J (1982). Method of determining the osmotic fragility curves of erythrocytes in birds. *Laboratory Animals*, **16**(1): 48-50.
- Wahab AA, Mabrouk MA, Ayo JO, Sullaiman AF, Muftau S, Yahaya A & Olutobi SE (2010). Effects of co-administration of antioxidants on erythrocyte osmotic fragility of wistar rats during the hot-dry season. *European Journal of Scientific Research*, **46**(1): 73-79.
- Wang C, Qin X, Huang D, He F & Zeng C (2010). Hemolysis of human erythrocytes induced by melamine-cyanurate complex. *Biochemical and Biophysical Research Communications*, **402**(4): 773-777.
- Wyse SJ, Velez JS, Stahringer RC, Greene LW & Randel RD (1991). Effects of diets containing free gossypol on erythrocyte fragility and packed cell volume in cattle. *Journal of Animal Science*, **69**(Suppl 1): 43-47.
- Yagil R, Luchtenstein C & Meyerstein MD (1976). Haemoconcentration and erythrocyte fragility in chickens exposed to heat and dehydration. *American Journal of Veterinary Research*, **37**(2): 103-106.
- Yagil R, Sod-Moriah UA & Meyerstein N (1974). Dehydration and camel blood. III. Osmotic fragility, specific gravity and osmolality. *American Journal of Physiology*, **226**(2): 305-308.
- Yamamoto R & Suzuki T (1982). Decreased membrane fragility of mouse erythrocytes by small doses of methylmercury and its restoration by coadministered selenite. *Tohoku Journal of Experimental Medicine*, **137**(3): 297-303.
- Yamamura H, Miyhara M & Kimura T (1991). Successful 0°C liquid preservation of red blood cells. *International Journal of Hematology*, **54**(3): 189-194.
- Yoong WC, Tuck SM & Micheal AE (2003). Binding of ovarian steroids to erythrocytes in patients with sickle cell disease: Effects on cell sickling and osmotic fragility. *Journal of Steroid Biochemistry and Molecular Biology*, **84**(1): 71-78.
- Yuan Y, Fang ZY & Zang ZH (1988). Changes in the rate of haemolysis during the early stage after burns in the rabbit. *Burns Including Thermal Injuries*, **14**(5): 365-368.
- Yusof A, Leithauser RM, Roth HJ, Finkernagel H, Wilson MT & Beneke R (2007). Exercise-induced hemolysis is caused by protein modification and most evident during the early phase of an ultraendurance race. *Journal of Applied Physiology*, **102**(2): 582-586.
- Zatta P, Perazzolo M & Corain B (1989). Tris acetylacetonate aluminium (III) induces osmotic fragility and acanthocyte formation in suspended erythrocytes. *Toxicology Letters*, **45**(1): 15-21.
- Zhang Q, Cao Z, Sun X, Zuang C, Huang W & Li Y (2016). Aluminium trichloride induces hypertension and disturbs the function of erythrocyte membrane in male rats. *Biology Trace Element Research*, **171**(1): 116-123.
- Zolla L, Lupidi G & Amiconi G (1994). Effect of mercuric ions on human erythrocytes. 1. Rate of haemolysis induced by osmotic shock as a function of incubation time. *Toxicology In vitro*, **8**(3): 483-490.