

D-Cycloserine enhances spatial learning performances of Rats chronically exposed to lead during the developmental period

Khaled Kahloula^{1,2}, Miloud Slimani², Martine Dubois¹, Jean-Jacques Bonnet¹

¹) *Unité de neuropsychopharmacologie de la dépression, CNRS FRE 2735, IFRMP 23, 22 Bd Gambetta, 76183 Rouen Cedex, France.*

²) *Laboratoire de Biochimie, Département de Biologie, Faculté des Sciences, Université d'Es-Senia, Oran 31000, Algérie.*

Accepté le 08/04/2009

إن الدراسات التي أجريت على الإنسان و الحيوان بينت أن التسمم المزمن بالرصاص إثر التطور الجنيني يؤدي إلى اضطرابات في السلوك. كما تبين أن المستقبلات من نوع NMDA تعد هدفا للرصاص و لنظام glutamate الذي يعتبر نظام هام و أساسي لظاهرة التعلم و التذكر. لوحظ في السنوات الأخيرة أن D CycloSérine (DCS) يحفز مستقبلات NMDA ، و قد يستطيع بدوره تحسين الوظائف السلوكية. إن عملنا يقتصر على الدراسة التطبيقية لمعرفة تأثير أسيتات الرصاص و ذلك بتركيز 0,2 % في ماء الشرب خلال فترة الحمل والإرضاع (GL) لدى الجرذان ، و ذلك لملاحظة وظائف التعلم و التذكر. و قد قيمنا قدرة D cyclosérine بتركيز 30ملغ/مل بحقن يومي للجرذان لمدة 15 يوما بعد الفطام . بعدها لاحظنا أن هذا المحلول قد خفف من أثر الرصاص على سلوك الحيوان. بينت النتائج أن الجرذان المعرضة للتسمم بالرصاص إثر فترة الحمل و الإرضاع تستغرق وقت أكثر و تقطع مسافة أكبر في التعلم ، و ذلك للوصول إلى السطح المغمور في مسبح "موريس" مقارنة مع الحيوانات غير المسممة. أما الحيوانات التي تعرضت للتسمم خلال التطور الجنيني و تحصلت على حقنة DCS ، نلاحظ أن لديها تحسن بليغ في التمرن و هذا مقارنة مع الحيوانات التي لم تحصل على حقنة DCS. و عليه فإن تحسين عمل المستقبلات NMDA من طرف DCS التي كانت تحت تأثير الرصاص يستطيع أن يكون إستراتيجية ناجحة و فعالة لأجل إصلاح العجز في التعلم الناتج عن التسمم بالرصاص.

الكلمات المفتاحية: جرذان؛ تسمم عصبي؛ رصاص؛ تطور جنيني؛ D cyclosérine (DCS).

Résumé

Des études faites chez l'homme et l'animal montrent qu'une exposition chronique au plomb durant le développement provoque des altérations des fonctions cognitives. Les récepteurs au glutamate de type N-methyl-D-aspartate (NMDA), importants dans les mécanismes synaptiques impliqués dans l'apprentissage et la mémoire, sont les principales cibles du plomb. Il a été montré que le D-cyclosérine, un agoniste partiel du site glycine associé aux récepteurs NMDA, peut améliorer les fonctions cognitives. Nous avons étudié les effets possibles d'une exposition au plomb (acetate de plomb 0.2% dans l'eau de boisson) pendant la gestation et la lactation (GL), sur les performances d'apprentissage spatial et de mémorisation chez des rats âgés de 32 jours. Nous avons également évalué la capacité du D-cyclosérine (30 mg/ml), administré quotidiennement 24h après le sevrage durant 15 jours, à atténuer les déficits cognitifs induits par le plomb. Nos résultats indiquent que les rats exposés au plomb pendant la gestation et la lactation mettent significativement plus de temps et parcourent une plus grande distance que les animaux contrôles pour trouver la plateforme immergée au cours de la phase d'apprentissage de la piscine de Morris. Toutefois, l'administration de DCS aux animaux GL améliore significativement leurs performances d'apprentissage comparées à celles des animaux GL non traités au DCS. Aucune différence de motivation ou des capacités de guidage visuo-moteur ne sont présents chez les GL. Donc, l'amélioration de la fonction des récepteurs NMDA pourrait être une stratégie efficace pour améliorer les déficits d'apprentissage spatial associés à la neurotoxicité au plomb.

Mots clés : Neurotoxicité; Plomb; Développement; D-Cyclosérine; Apprentissage; Rat.

Abstract

Chronic developmental lead (Pb) exposure has long been associated with cognitive dysfunction in children and animals. N-methyl-D-aspartate (NMDA) receptors, important in the synaptic mechanisms involved in learning and memory, are key target of lead toxicity. D-cycloserine (DCS), a partial agonist of the NMDA-

Auteur correspondant: bombixc2@yahoo.fr (Khaled Kahloula)

associated glycine site, has been recognised as a potential cognitive enhancer. We investigated the potential effects of Pb exposure (lead acetate 0.2% through the drinking water) during gestation and lactation (GL), on the spatial learning and memory capacities of PN32 rats. We also evaluate the ability of DCS (30 mg/ml), administered daily 24h after weaning during 15 days, to attenuate Pb neurotoxicity-induced cognitive deficits. Results indicate that rats exposed to lead during gestation and lactation have a significantly increased latency to find the hidden platform and cover a significant longer distance compared to control-vehicle in the learning phase of the Morris water maze. However, the administration of DCS to GL animals improved significantly their learning performances compared with GL-vehicle. In contrast, there is no significant difference between all groups during the probe test and the visual cue test. In conclusion, DCS enhancement of the NMDA receptor function is an effective strategy to ameliorate neurotoxicity lead-associated spatial learning deficits.

Key words: Neurotoxicity; Lead; Development; D-Cycloserine; Learning; Rat.

1. INTRODUCTION

Lead (Pb) is a common environmental neurotoxic agent that is a major public health problem. Lead poisoning affects particularly the developing central nervous system [1-3]. Exposure to low levels of lead during early development is sufficient to cause long lasting behavioural abnormalities and cognitive deficits in human and experimental animals [4,5]. Pb-toxicity induces diminished IQ scores and reduced learning and memory capacities in children exposed early in life [6,7]. Rodents exposed to lead are impaired in cognitive tasks such as the step-down avoidance test, the contextual fear conditioning, the radial maze, the Morris water maze (MWM) [8-11].

Lead alters glutamatergic function and also interferes with cholinergic and dopaminergic neurotransmission [12-15]. This neurotoxicant impairs neurogenesis, synaptic plasticity, long-term potentiation (LTP) and cognitive function in experimental animals [9,16,17]. Several studies have indicated impairment in the induction, expression and maintenance of LTP in the hippocampus of animal exposed to lead during the early developmental period [18-21]. The detrimental effects of lead on LTP and spatial learning have been attributed in one hand to the effect of lead on glutamate release [22,23] and in an other hand, to the fact that lead is a potent inhibitor of N-methyl-D-aspartate (NMDA) receptors [24,25]. Further,

chronic exposure to lead alters the expression of NMDA subunit in the rat hippocampus [22,26,27], decreasing particularly the NR2A subunit expression, thus affecting NMDA complex composition, which is important in hippocampal development and maturation. Pb-exposition also leads to an alteration of downstream calcium signalling. Recent findings demonstrated that early lead exposure disrupts expression and phosphorylation of the cAMP-response element binding protein (CREB), a transcription factor important in synaptic plasticity and in learning and memory [28,29]. Therefore, there is compelling experimental evidence that chronic lead exposure alters NMDA function in the rat hippocampus and this effects is associated with impairments in LTP and spatial learning [17,30].

An improvement of performances in various learning tasks has been described after administration of D-cycloserine (DCS), a compound originally used as an antibiotic agent for the treatment of tuberculosis and later characterised as a partial agonist at the glycine modulatory site of the glutamatergic NMDA receptor [31,32]. DCS is a cognitive enhancer in learning tests, such as the trace eye blink conditioning and the MWM task that are closely dependent on the hippocampus [33-35]. Moreover, DCS improves spatial performances of aged memory-deficient rats [36,37].

Based on the above considerations, the

aims of the present work were to study the effects of a chronic lead exposure during gestation and lactation on spatial learning and memory abilities of Wistar rats and to evaluate the effect of DCS on their performances in the MWM.

2. MATERIAL AND METHODS

2.1 Animals and treatment

Experiments were carried out on Wistar rats (obtained from Charles River) weighing 200-350 g. Animals were housed with free access to water and food in an animal room with 12/12h light/dark cycle, at 22 ± 2 °C. They were mated one week after their arrival (three females and one male per cage). On pregnancy day 0 the dams were divided into 2 groups: one group received 0.2% lead acetate (Sigma-Aldrich Incorporated, France) in drinking distilled water during gestation and lactation (GL), whereas rats of the control group received distilled water without lead acetate (C), as described previously [15]. At birth, GL pups continued to receive lead acetate during lactation until postnatal day (PN) 21 and rats were weaned this day. Immediately after the behavioural evaluation, the rats were decapitated and blood samples were taken for Pb blood determination. Number and suffering of animals were minimised in accordance with the guidelines of the European Council Directive (86/609/EEC).

2.2 Drug preparation

D-Cycloserine (DCS) was obtained from Sigma-Aldrich Incorporated (France). DCS was dissolved in distilled water and administered (i.p.) with a constant injection volume of 1 ml/kg. In order to maintain stability of the solution across days, DCS was frozen (-20 °C) in 1 ml aliquots. Each day, DCS was thawed 15 min prior to injection.

2.3 Experimental design: chronic administration prior to assessment

Animals were exposed to Pb during gestation and lactation (GL) and controls animals receive distilled water. In order to test the ability of DCS to attenuate Pb neurotoxicity induced cognitive deficits, the drug therapy was administered beginning 24 h after weaning. Randomly chosen animals of each group were injected (i.p.) with 30 mg/ml DCS (n=5; n=7, respectively) or distilled water (vehicle) (n=8; n=5, respectively). All animals were injected once daily on days 1-15 after weaning. During cognitive assessment (PN days 32-36) all subjects were given their injection 30 min prior to begin the MWM task.

2.4 Morris Water Maze Test

The Morris water maze task was used to assess cognitive function, in particularly spatial learning and memory abilities [38]. In PN 32 rats, the test was conducted in a round white pool 90 cm in diameter with a 50 cm high wall. The pool was filled to a depth of 30 cm with water made opaque with white, non toxic water based paint. Water temperature was maintained at 21 ± 1 C° with an aquarium heater. The platform of 10 cm in diameter and 28 cm high (i.e., 2 cm below the water's surface) was used as the hidden goal. The pool was located in a room with numerous extra-maze cues that remained constant throughout the experiment. Animals were tested on days 11-15 after weaning, with four trials per day for five consecutive days. On each trial, the animal was placed in the pool, facing the wall, at one of the four start locations (north (N), south (S), east (E), and west (W)). Start location sequences were randomly assigned each day prior to the start of testing. Rats were allowed to swim to the submerged platform placed in the NW quadrant during 60 s. If the platform was not found within the allowed time, the

rat was manually placed onto it. Rats were allowed to rest on the platform for 20 s between each trial. A session of four trials was conducted each day from 9:00 to 11:00 A.M.. The experimenter was hidden from the view of the animals. The escape latency (time to find the platform) and path length (distance travelled to the hidden platform) were recorded by using a videotrack system (Viewpoint, France). A probe test was conducted the day following the end of the training session to further characterise swim task performance. In this test, the platform was removed and the rat was placed next to and facing the S side. Animals were allowed to swim freely for the original training session length of time (60 s) for a single trial. The time spent by animals in the previously correct quadrant (NW) was measured. Visual cue test was performed two hours after the end of the probe test in order to evaluate the visuo-motor abilities and the motivation of the animals. It was conducted by extending a large black flag above the water level from the submerged platform. The maximum time allowed was the same as the original training sessions. The escape latency and path length were measured during a single session of four trials.

2.5 Determination of blood lead level

Blood samples were collected immediately after decapitation of animals in tubes pre-treated with 10 µl of heparin. The lead concentration in the blood was determined by using atomic absorption spectroscopy (model 800, Perkin Elmer).

2.6 Statistical analysis

Results are expressed as mean \pm standard error of the mean (SEM). Lead blood levels were compared between GL-vehicle and C-vehicle with a t-test. MWM data were analysed by two-way analyses of variance (ANOVAs), with the intoxication factor (GL or controls) and

the treatment factor (DCS or vehicle), or a three-way ANOVA with day as the repeated measure. When a significant difference was found, the Student-Newman-Keuls post-hoc test was conducted. For all analyses, a difference was considered significant at $p \leq 0.05$.

3. RESULTS

A student test reveals that lead concentration in blood of rats exposed to lead during gestation and lactation is significantly higher than the one of control rats [$t(13)=11.78$, $p < 0.001$]. Lead concentrations in blood are 30.13 (± 23.57) µg/dl for GL (n=8) and 0.30 (± 0.61) µg/dl for control (n=7) rats.

A three-way ANOVA with repeated measures comparing MWM cognitive performances during the training phase reveals a significant interaction between the treatment, intoxication and the repeated measure factors for latency to find the hidden platform, which means that animals perform differently according to their intoxication and treatment across days [$F(3,63)=6.23$, $p < 0.001$]. An intoxication x treatment x day interaction also appears for distance crossed [$F(3,63)=4.41$, $p < 0.01$].

A two-way ANOVA was conducted to compare the learning performances of rats day by day. There is no significant difference between all groups during the first and second days of training in terms of latency to find the hidden platform and swimming distance ($p > 0.05$). On the third day of learning, assessment the ANOVA reveals a significant treatment effect for escape latency [$F(1,21) = 4.65$, $p < 0.05$] and distance crossed [$F(1,21) = 5.79$, $p < 0.05$]. Rats treated with DCS have significant better learning performances than control rats. A significant intoxication by treatment interaction is observed for the fourth day of training in terms of escape latency and path length ([$F(1,21) = 5.65$, $p < 0.05$], [$F(1,21) = 6.02$, $p < 0.05$] respectively). Post-hocs reveals

that rats exposed to lead during gestation and lactation and treated with vehicle have a significant increased escape latency ($p < 0.01$) and cover a significant longer distance ($p < 0.05$) compared to control-vehicle. The administration of DCS to GL animals improves significantly their learning performances compared with GL-vehicle in terms of latency to find the hidden platform ($p < 0.01$) and distance crossed ($p < 0.01$). The performances of GL rats treated with DCS do not differ significantly from those of control-DCS rats ($p > 0.05$).




Figure 1. Escape latency (A) and distance crossed before finding the platform (B) during the training phase of the Morris water maze task of rats exposed to lead during gestation and lactation (GL) and control rats (C), treated with D-cycloserine (DCS) or vehicle. Means \pm SEM. *: $p < 0.05$, **: $p < 0.01$: GL-vehicle vs C-vehicle; #: $p < 0.01$: GL-vehicle vs GL-DCS (ANOVA followed by Student-Newman-Keuls post-hoc).

During the probe test, there is no significant difference between all groups for the time spent in the NW quadrant ($p > 0.05$).




Figure 2. Time spent in the training quadrant (NW) during the probe test by rats exposed to lead during gestation and lactation (GL) and control rats (C), treated with D-cycloserine (DCS) or vehicle. Means \pm SEM. No significant difference is observed between the different groups.

During the visual cue test, GL groups (treated with DCS or vehicle) do not present significantly different performances compared with control group in terms of escape latency and distance crossed ($p > 0.05$).




Figure 3. Escape latency (A) and distance crossed before finding the platform (B) during the cue phase of the Morris water maze task of rats exposed to lead during gestation and lactation (GL) and control rats (C), treated with D-cycloserine (DCS) or vehicle. Means \pm SEM. No significant difference is observed between the different groups.

4. DISCUSSION

The present study shows that, on one hand, a chronic lead intoxication of Wistar rats during gestation and lactation resulted in decreased learning performances in the MWM. On the other hand, the chronic administration of DCS at the dose of 30 mg/kg, on days 1-15 after weaning ameliorates significantly the cognitive impairments of rats exposed to lead until weaning.

Several studies have revealed that Pb exposure produces neurological damage and behavioural disruptions in human beings and in experimental animals [39-43]. The lowest Pb blood level at which behavioural abnormalities are detectable is 15–20 µg/dl in rodents and 10–15 µg/dl in children, as estimated by [44]. In our study, the blood lead level of animals exposed during the gestation and lactation is superior to the lowest Pb blood level at which are observed modification of behaviour and is close to the one described previously [8].

The present findings show that a Pb exposure during the gestation and lactation period causes learning deficits in offspring when blood Pb concentrations are increased than at control levels, as shown by an increase of escape latency and distance crossed before finding the hidden platform in the MWM test. Altered performances in water maze task have been described in rats exposed to Pb during gestation and lactation [10,45]. The period of lead exposure to observe cognitive abnormalities is important as a lead exposition during post-weaning period [46,47] or during one part of the gestation period [16,48] appears not to be sufficient to cause learning deficits in the MWM. We observe no difference in terms of time spent in the target quadrant (NW) between all groups during the probe test. This result is in good agreement with those of Kuhlmann et al. [45]. Moreover, during the visible platform phase no significant difference is observed between

all the groups. This indicates that deficits of GL-vehicle rats during the training phase are not due to visual or motivational troubles.

The glutamatergic system, specifically the NMDA receptor complex, is a key target site of lead neurotoxic action [1]. Lead exposure induces NMDA receptor functions changes [49], induces a wide variety of changes of intracellular signalling [50] and synaptic plasticity in hippocampus [30,51] that are known to interfere with learning and memory.

One pharmacological agent that has received attention as a promising cognitive-enhancer is the NMDA-glycine site partial agonist, D-cycloserine. Various studies indicate that DCS is an effective cognitive enhancer in multiple paradigms of cognitive impairment, including models of ageing and neurological lesions [37,52,53]. The aim of our work was to evaluate the possible effect of DCS in the case of a lead intoxication during development. Our results show that the learning deficits of rats exposed to lead during the gestation and lactation could be corrected by a chronic administration of DCS, as there is no significant difference between the performances of GL-DCS rats and controls rats in the learning phase of the MWM.

DCS has already been shown to ameliorate the cognitive alterations of brain-injured rats and muscarinic antagonists treated rats in the MWM [54-56]. While the mechanism by which DCS improve spatial learning performances in animals is unknown, it is possible that the cognitive-enhancing effect of DCS is the result of several different physiological actions. It has been suggested that DCS function to improve cognitive performances by activating the glycine site and accelerating recovery of the NMDA receptor from desensitization [57,58].

In conclusion, the results from the present study show that learning deficits induced by a chronic Pb intoxication

during gestation until weaning can be corrected by a chronic administration of DCS. Understanding the cellular action of DCS can help to find a strategy to prevent and treat the effect of Pb intoxication during development.

Acknowledgments

This work was supported by a Franco-Algerian grant program. We would like to thank Pr Jean Costentin. We would like to express our sincere gratitude to Pr Isabelle Leroux for her advice. We also thank the "Laboratoire de Toxicologie, CHU de Rouen" for the blood lead concentrations measurements.

References

- [1] D.P. Alfano, J.C. Leboutillier et T.L. Petit, *Hippocampal mossy fiber pathway development in normal and postnatally lead-exposed rats*, *Exp Neurol*, Vol 75, 1982, pp.308-19.
- [2] R.A. Goyer, *Lead toxicity: current concerns*, *Environ. Health. Perspect.*, Vol. 100, 1993, pp. 177-87.
- [3] E.K. Silbergeld, *Mechanisms of lead neurotoxicity, or looking beyond the lamppost*, *FASEB J.*, Vol. 6, 1992, pp. 3201-6.
- [4] E.G. Moreira, I. Vassilieff, V.S. Vassilieff, *Developmental lead exposure: behavioral alterations in the short and long term*. *Neurotoxicol. Teratol.*, Vol. 23, 2001, pp. 489-95.
- [5] K.J. Murphy, C.M. Regan, *Low-level lead exposure in the early postnatal period results in persisting neuroplastic deficits associated with memory consolidation*, *J. Neurochem.*, Vol. 72, 1999, pp. 2099-104.
- [6] D. Bellinger, Leviton A, M. Rabinowitz, E. Allred, H. Needleman et S. Schoenbaum, *Weight gain and maturity in fetuses exposed to low levels of lead*, *Env. Res*, Vol. 54, 1991, p. 151-8.
- [7] R.L. Canfield, D.A. Kreher, C. Cornwell, C.R.J. Henderson, *Low-level lead exposure, executive functioning, and learning in early childhood*, *Child Neuropsychol*, Vol.9, 2003, pp. 35-53.
- [8] H. Chen, T. Ma et I. Ho, *Effects of developmental lead exposure on inhibitory avoidance learning and glutamate receptors in rats*, *Environ Toxicol Pharmacol*, Vol. 9, 2001, pp. 185-191.
- [9] K. Jaako-Movits, T. Zharkovsky, O. Romantchik, M. Jurgenson, E. Merisalu, L. Heidmets et al., *Developmental lead exposure impairs contextual fear conditioning and reduces adult hippocampal neurogenesis in the rat brain*, *Int. J. Dev. Neurosci.*, Vol. 23, 2005, pp. 627-35.
- [10] D.A. Jett, A.C. Kuhlmann, S.J. Farmer et T.R. Guilarte, *Age-dependent effects of developmental lead exposure on performance in the Morris water maze*, *Pharmacol. Biochem. Behav.*, Vol. 57, 1997, pp. 271-9.
- [11] C. Munoz, K. Garbe, H. Lilienthal, G. Winneke, *Neuronal depletion of the amygdala resembles the learning deficits induced by low level lead exposure in rats*, *Neurotoxicol. Teratol.*, Vol. 11, 1989, pp. 257-64.
- [12] H. Bielarczyk, J.L. Tomsig et J.B. Suszkiw, *Perinatal low-level lead exposure and the septo-hippocampal cholinergic system: selective reduction of muscarinic receptors and cholineacetyltransferase in the rat septum*, *Brain Res*. Vol. 643, 1994, pp. 211-7.
- [13] D.A. Cory-slechta, *Relationships between lead-induced learning impairments and changes in dopaminergic, cholinergic, and glutamatergic neurotransmitter system functions*, *Annu. Rev. Pharmacol. Toxicol.*, Vol. 35, 1995, pp. 391-415.
- [14] S.M. Lasley, M.E. Gilbert, *Glutamatergic components underlying*

lead-induced impairments in hippocampal synaptic plasticity, Neurotoxicology, Vol. 21, 2000, pp. 1057-68.

[15] D. Noureddine, S. Miloud et A. Abdelkader, *Effect of lead exposure on dopaminergic transmission in the rat brain*, Toxicol., Vol. 207, 2005, pp. 363-8.

[16] M.E. Gilbert, M.E. Kelly, T.E. Samsam et J.H. Goodman, *Chronic developmental lead exposure reduces neurogenesis in adult rat hippocampus but does not impair spatial learning*, Toxicol. Sci., Vol. 86, 2005, pp. 365-74.

[17] C.D. Toscazo et T.R. Guilarte, *Lead neurotoxicity: from exposure to molecular effects*, Brain Res. Rev., Vol. 49, 2005, pp. 529-54.

[18] L. Altmann, F. Weinsberg, K. Sveinsson, H. Lilienthal, H. Wiegand et G. Winneke, *Impairment of long-term potentiation and learning following chronic lead exposure*. Toxicol. Lett. 66, 1993, pp. 105-12.

[19] M.E. Gilbert, C.M. Mack, *Chronic lead exposure accelerates decay of long-term potentiation in rat dentate gyrus in vivo*, Brain Res. V. 789, 1998, pp.139-49.

[20] M.E. Gilbert, C.M. Mack, S.M. Lasley, *Chronic developmental lead exposure increases the threshold for long-term potentiation in rat dentate gyrus in vivo*, Brain Res., V. 736, 1996, p. 118-24.

[21] D.Y. Ruan, J.T. Chen, C. Zhao, Y.Z. XU, M. Wang et W.F. Zhao, *Impairment of long-term potentiation and paired-pulse facilitation in rat hippocampal dentate gyrus following developmental lead exposure in vivo*, Brain Res., Vol. 806, 1998, pp. 196-201.

[22] S.M. Lasley, M.E. Gilbert, *Presynaptic glutamatergic function in dentate gyrus in vivo is diminished by chronic exposure to inorganic lead*, Brain Res., Vol. 736, 1996, pp. 125-34.

[23] S.M. Lasley, M.E. Gilbert, *Rat hippocampal glutamate and GABA release exhibit biphasic effects as a function of chronic lead exposure level*, Toxicol. Sci., Vol. 66, 2002, pp. 139-47.

[24] P. Gavazzo, A. Gazzoli, M. Mazzolini et C. Marchetti, *Lead inhibition of NMDA channels in native and recombinant receptors*, Neuroreport., Vol. 12, 2001, pp. 3121-5.

[25] T.R. Guilarte, R.C. Miceli, *Age-dependent effects of lead on [3H]MK-801 binding to the NMDA receptor-gated ionophore: in vitro and in vivo studies*, Neurosci. Lett., Vol. 148, 1992, pp. 27-30.

[26] T.R. Guilarte, J.L. Mcglothan, M.K. Nihei, *Hippocampal expression of N-methyl-D-aspartate receptor (NMDAR1) subunit splice variant mRNA is altered by developmental exposure to Pb(2+)*, Mol Brain Res., Vol. 76, 2000, pp. 299-305.

[27] M.K. Nihei et T.R. Guilarte, *Nmdar-2a subunit protein expression is reduced in the hippocampus of rats exposed to Pb2+ during development*, Mol. Brain Res., Vol. 66, 1999, pp. 42-9.

[28] C.D. Toscano, H. Hashemzadeh-Gargari, J.L. Mcglothan et T.R. Guilarte, *Developmental Pb2+ exposure alters NMDAR subtypes and reduces CREB phosphorylation in the rat brain*. Dev. Brain Res., Vol. 139, 2002, pp. 217-26.

[29] C.D. Toscano, J.L. Mcglothan, T.R. Guilarte, *Lead exposure alters cyclic-AMP response element binding protein phosphorylation and binding activity in the developing rat brain*, Dev. Brain Res., Vol. 145, 2003, pp. 219-28.

[30] M.K. Nihei, N.L. Desmond, J.L. Mcglothan, A.C. Kuhlmann et T.R. Guilarte, *N-methyl-D-aspartate receptor subunit changes are associated with lead-induced deficits of long-term potentiation and spatial learning*, Neuroscience, Vol. 99, 2000, pp. 233-42.

- [31] W.F. Hood, R.P. Compton, J.B. Monahan, *D-cycloserine: a ligand for the N-methyl-D-aspartate coupled glycine receptor has partial agonist characteristics*, *Neurosci. Lett.*, Vol. 98, 1989, pp. 91-5
- [32] G.B. Watson, M.A. Bolanowski, M.P. Baganoff, C.L. Deppeler et T.H. Lanthorn, *D-cycloserine acts as a partial agonist at the glycine modulatory site of the NMDA receptor expressed in Xenopus oocytes*, *Brain Res.*, Vol. 510, 1990, pp. 158-60.
- [33] V. Lelong, F. Dauphin, M. Boulouard, *RS 67333 and D-cycloserine accelerate learning acquisition in the rat*, *Neuropharmacology*, Vol. 41, 2001, pp. 517-22.
- [34] D. Quartermain, J. Mower, M.F. Rafferty, R.L. Herting et T.H. Lanthorn, *Acute but not chronic activation of the NMDA-coupled glycine receptor with D-cycloserine facilitates learning and retention*, *Eur. J. Pharmacol.*, Vol. 257, 1994, pp. 7-12.
- [35] L.T. Thompson, J.R. Moskal et J.F. Disterhoft, *Hippocampus-dependent learning facilitated by a monoclonal antibody or D-cycloserine*, *Nature*, Vol. 359, 1992, pp. 638-41.
- [36] J. Aura et P.J. Riekkinen, *Pre-training blocks the improving effect of tetrahydroaminoacridine and D-cycloserine on spatial navigation performance in aged rats*, *Eur J Pharmacol*, Vol. 390, 2000, pp. 313-8.
- [37] M.G. Baxter, T.H. Lanthorn, K.M. Frick, S. Golski, R.Q. Wan et D.S. Olton, *D-cycloserine, a novel cognitive enhancer, improves spatial memory in aged rats*, *Neurobiol Aging*, Vol. 15, 1994, pp. 207-13.
- [38] R. Morris, *Developments of a water-maze procedure for studying spatial learning in the rat*, *J. Neurosci. Methods*, Vol. 11, 1984, pp. 47-60.
- [39] D.P. Alfano, J.C. Leboutillier et T.L. Petit, *Hippocampal mossy fiber pathway development in normal and postnatally lead-exposed rats*, *Exp Neurol*, Vol 75, 1982, pp.308-19.
- [40] D.A. Cory-slechta et D.V. Widzowski, *Low level lead exposure increases sensitivity to the stimulus properties of dopamine D1 and D2 agonists*, *Brain Res.*, Vol. 553, 1991, pp. 65-74.
- [41] M.E. Gilbert, C.M. Mack, S.M. Lasley, *The influence of developmental period of lead exposure on long-term potentiation in the adult rat dentate gyrus in vivo*, *Neurotoxicology*, Vol 20, Issue a, 1999, pp.57-69.
- [42] E. Kiraly et D.G. Jones, *Dendritic spine changes in rat hippocampal pyramidal cells after postnatal lead treatment: a golgi study*, *Exp. Neurol.*, Vol. 77, 1982, pp. 236-9.
- [43] B.P. Lanphear, K. Dietrich, P. Auinger, C. Cox, *Cognitive deficits associated with blood lead concentrations <10 microg/dl in US children and adolescents*, *Public. Health. Rep.*, Vol. 115, 2000, pp. 521-9.
- [44] J.M. Davis, D.A. Otto, D.E. Weil et L.D. Grant, *The comparative developmental neurotoxicity of lead in humans and animals*, *Neurotoxicol. Teratol.*, Vol. 12, 1990, pp. 215-29.
- [45] A.C. Kuhlmann, J.L. Mcglothan et T.R. Guilarte, *Developmental lead exposure causes spatial learning deficits in adult rats*, *Neurosci. Lett.*, Vol. 233, 1997, pp. 101-4.
- [46] G. Garcia-Arenas, V. Ramirez-Amaya, I. Balderas, J. Sandoval, M.L. Escobar, C. Rios et al., *Cognitive deficits in adult rats by lead intoxication are*

related with regional specific inhibition of cNOS, *Behav Brain Res.*, Vol. 2004, pp. 49-59.

[47] J.S. Schneider, M.H. Lee, D.W. Anderson, L. Zuck et T.I. Lidsky, *Enriched environment during development is protective against lead-induced neurotoxicity*, *Brain Res.*, Vol. 896, 2001, pp. 48-55.

[48] H.M. Newman, R.S.H. Yang, K.R. Magnusson, *Effects of developmental exposure to lead, magnesium and zinc mixtures on spatial learning and expression of NMDA receptor subunit mRNA in fischer 344 rats*, *Toxicol. Lett.*, Vol. 126, 2002, pp. 107-19.

[49] D.A. Cory-slechta, C.L. Flaugh, S.B. Evans, M.J. Pokora et J.T. Greenamyre, *Susceptibility of adult rats to lead-induced changes in NMDA receptor complex function* *Neurotoxicol Teratol*, Vol. 19, 1997, pp. 517-30.

[50] G.J. Flora et P.K. Seth, *Alterations in some membrane properties in rat brain following exposure to lead*, *Cytobios.*, Vol.103, 2000, pp. 103-9.

[51] M.E. Gilbert, C.M. Mack, S.M. Lasley, *Chronic developmental lead exposure and hippocampal long-term potentiation: biphasic dose-response relationship*, *Neurotoxicology* Vol 20, Issue b, 1999, pp. 71-82.

[52] J.F. Flood, J.E. Morley et T.H. Lanthorn, *Effect on memory processing by D-cycloserine, an agonist of*

the NMDA/glycine receptor, *Eur. J. Pharmacol.*, Vol. 221, 1992, pp. 249-54.

[53] G.M. Schuster et W.J. Schmidt, *D-cycloserine reverses the working memory impairment of hippocampal-lesioned rats in a spatial learning task*, *Eur. J. Pharmacol.*, Vol. 224, 1992, pp. 97-8.

[54] P.J. Riekkinen, S. Ikonen et M. Riekkinen, *Tetrahydroaminoacridine, a cholinesterase inhibitor, and D-cycloserine, a partial NMDA receptor-associated glycine site agonist, enhances acquisition of spatial navigation*, *Neuroreport*, Vol. 9, 1998, pp. 1633-7.

[55] J. Sirviö, T. Ekonsalo, P.J. Riekkinen, H. Lahtinen et P.S. Riekkinen, *D-cycloserine, a modulator of the N-methyl-D-aspartate receptor, improves spatial learning in rats treated with muscarinic antagonist*, *Neurosci. Lett.*, Vol. 146, 1992, pp. 215-8.

[56] M.D. Temple et R.J. Hamm, *Chronic, post-injury administration of D-cycloserine, an NMDA partial agonist, enhances cognitive performance following experimental brain injury*, *Brain Res.*, Vol. 741, 1996, pp. 246-51.

[57] A.C. Foster, J.A. Kemp, *Neurobiology, Glycine maintains excitement*, *Nature*, Vol. 338, 1989, pp. 377-8.

[58] M.L. Mayer, L.J. Vyklicky, *Clements J. Regulation of NMDA receptor desensitization in mouse hippocampal neurons by glycine*, *Nature*, Vol. 338, 1989, pp. 425-7.