

Successful pregnancy in a unicornuate uterus

OSEMEN EHIDIAMEN OKOJIE, BABASOLA O OKUSANYA, EDIRIN EDWARD AGADAGBA¹, ADESUMBO KAYINSOLA ODESEYE

Department of Obstetrics and Gynaecology, Lagos University Teaching Hospital, Idi-Araba, Lagos State, ¹Department of Obstetrics and Gynaecology, Irrua Specialist Teaching Hospital, Irrua, Edo State, Nigeria

ABSTRACT

Uterine anomalies are generally rare and are usually asymptomatic. When present, these anomalies are implicated in complications such as recurrent miscarriages, preterm labor, malpresentation, intrauterine growth restriction, uterine rupture, and a variety of menstrual disturbances. We report a case of a pregnancy carried to term in a unicornuate uterus with an accessory horn, diagnosed intraoperatively during an emergency cesarean section with favorable outcome to the mother and baby.

Key words: Cesarean section; successful pregnancy; unicornuate uterus; uterine anomaly.

Introduction

Failure of development and fusion of the Mullerian (paramesonephric) ducts result in the various uterine anomalies.^[1] These anomalies have various complications that may arise in pregnancy. Unicornuate uterus is classified as Type II by the American Society for Reproductive Medicine Classification of uterine anomalies [Figure 1].^[2] Further classification is made into communicating, no cavity, and no horn.^[2] In majority of cases, the rudimentary horn is noncommunicating and has a cavity.^[2] From literature, unicornuate uterus is reported to account for 2.4%–13% of all Mullerian anomalies.^[3] Pregnancy in the rudimentary horn is usually very rare and may be complicated by uterine rupture, usually toward the end of the second trimester.^[4]

Case Report

A 28-year-old booked G3P0 + 2 female presented with labor pains at 40 weeks and 6 days of gestational age. On examination, her general condition was satisfactory. On abdominal examination, the abdomen was enlarged with an unusual swelling in the left iliac fossa region, measuring about 10 cm × 12 cm, which had not been

previously noticed during the antenatal period. She had regular uterine contractions and the fetus was in oblique lie and shoulder presentation, with regular fetal heart tones. An emergency bedside ultrasound scan made a suspicion of an ovarian cyst with abnormal lie, and she was subsequently counseled and scheduled for an emergency cesarean section and possible ovarian cystectomy. Findings at surgery revealed a right unicornuate uterus with its own fallopian tube and ovary and a well-formed lower uterine segment and accessory horn with its own tube and ovary as shown in Figures 2 and 3. A live female neonate weighing 3.6 kg with good APGAR scores was delivered and no other abnormality was noticed on exploration of the abdominal cavity. Estimated blood loss was 600 ml. Mother had no postsurgery complications.

Discussion

The diagnosis of unicornuate uterus may be missed in pregnancy, especially if the patient is asymptomatic. Even

Address for correspondence: Dr. Osemen Ehidiامن Okojie, Department of Obstetrics and Gynaecology, Lagos University Teaching Hospital, Idi-Araba, Lagos State, Nigeria. E-mail: dr.oeokojie@gmail.com

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: reprints@medknow.com

How to cite this article: Okojie OE, Okusanya BO, Agadagba EE, Odeseye AK. Successful pregnancy in a unicornuate uterus. Trop J Obstet Gynaecol 2018;35:364-6.

Access this article online	
Website: www.tjogonline.com	Quick Response Code 
DOI: 10.4103/TJOG.TJOG_10_18	




Figure 1: Classification of congenital uterine anomalies (American Society for Reproductive Medicine Classification)




Figure 2: Repaired right unicornuate uterus and left accessory horn with its tube and ovary



Figure 3: Right unicornuate uterus, its tube, and ovary

with ultrasound in the most experienced of hands, diagnosis is often not made. The woman presented had several ultrasound scans done which did not pick up the anomaly, including the scan done in labor, which revealed the mass as ovarian in origin. It may be that with contractions, the accessory horn, which was not visible initially, rotated to

the left lower abdomen. This then resulted in the abnormal lie and malpresentation. The patient's antenatal period was essentially uneventful from an obstetric point of view. Pregnancy was carried to term without any incidence of abdominal pain, preterm contractions, growth restriction or uterine rupture.

Forty percent of women with unicornuate uterus have renal abnormalities, which was not diagnosed in this case.^[5] A unicornuate uterus may cause an ectopic pregnancy and uterine rupture when the pregnancy is in the rudimentary horn. Therefore, it is advised to have the rudimentary horn excised if detected in the preconception period.^[6] Excision of the accessory horn could not be done in this patient because diagnosis was made at surgery, and blood loss would have been more. More so, consent had not been obtained for the procedure. The patient was counseled in the postoperative period on the findings at surgery. She was also counseled on the need to have a hysterosalpingography done and subsequently interval excision of the accessory horn.

Conclusion

This report highlights the challenges of making a diagnosis of unicornuate uterus in pregnancy. A high index of clinical suspicion and advanced imaging is needed to make a diagnosis, to institute timely obstetric management and prevent complications to mother and baby.

Declaration of patient consent

The authors certify that they have obtained all appropriate patient consent forms. In the form the patient(s) has/have given his/her/their consent for his/her/their images and other clinical information to be reported in the journal. The patients understand that their names and initials will not be published and due efforts will be made to conceal their identity, but anonymity cannot be guaranteed.

Financial support and sponsorship

Nil.

Conflicts of interest

There are no conflicts of interest.

References

1. Souvizi B, Jafarzadeh Esfehni R. A case of successful pregnancy in a complete bicornuate uterus. *J Midwifery Reprod Health* 2016;4:720-2.
2. Thakur S, Sood A, Sharma C. Ruptured noncommunicating rudimentary horn pregnancy at 19 weeks with previous cesarean delivery: A case report. *Case Rep Obstet Gynecol* 2012;2012:308476.
3. Caserta D, Mallozzi M, Meldolesi C, Bianchi P, Moscarini M. Pregnancy in a unicornuate uterus: A case report. *J Med Case Rep* 2014;8:130.

4. Rackow BW, Arici A. Reproductive performance of women with müllerian anomalies. *Curr Opin Obstet Gynecol* 2007;19:229-37.
5. Fedele L, Bianchi S, Agnoli B, Tozzi L, Vignali M. Urinary tract anomalies associated with unicornuate uterus. *J Urol* 1996;155:847-8.
6. Jayasinghe Y, Rane A, Stalewski H, Grover S. The presentation and early diagnosis of the rudimentary uterine horn. *Obstet Gynecol* 2005;105:1456-67.