

Original Research Article

Assessment of Tigecycline Prescription and Patients' Outcomes at Three Different Hospitals in Saudi Arabia

Mansour Tobaigy^{1,2*}, Saad Al Humaid³, Derek Stewart⁴, Fayez Omeir Alotaibi⁵, Kamal Ahmad Qureshi⁶, Katie MacLure⁴, Fahad Algharib³, Ahmed Alsameti³, Ahmed Alsaqer³ and Ahmad Almeman⁷

¹Patient Safety, Maternity and Children's Hospital, Ministry of Health, Jeddah, ²Department of Pharmacology, Faculty of Medicine, University of Jeddah, Jeddah, ³Administration of Pharmaceutical Care, Alhassa, Saudi Arabia, ⁴School of Pharmacy and Life Sciences, Robert Gordon University, Aberdeen, United Kingdom, ⁵Department of Inpatient Pharmacy, King Fahad Jeddah Hospital, Jeddah, ⁶Department of Microbiology, College of Pharmacy Unizah, ⁷Department of Pharmacology & Therapeutics, College of Medicine, University of Qassim, Qassim, Saudi Arabia

*For correspondence: **Email:** m.tobaigy@moh.gov.sa

Received: 15 May 2015

Revised accepted: 6 September 2015

Abstract

Purpose: To investigate tigecycline prescription and patient outcomes in the Kingdom of Saudi Arabia (KSA).

Methods: A retrospective observational study was conducted in three KSA government hospitals, between January, 2013 and May, 2014. The patients were identified from electronic prescription records; data were retrieved by trained researchers.

Results: Thirty-seven patients who received tigecycline were included (mean age, 52.5 years; range, 17–92); 51.4 % were female. Tigecycline was prescribed for sepsis (59.5 %), pneumonia (21.6 %), and/or intra-abdominal infections (13.5 %). The majority of the patients (86.5 %) were prescribed tigecycline in intensive care unit (ICU) and the remaining patients were in the general medical ward. APACHE II score at the beginning of treatment was 16.8 ± 4.3 , indicating severe disease. Susceptibility testing revealed 22 different bacterial pathogens, most commonly *Acinetobacter baumannii* (20 patients) and *Klebsiella pneumoniae* (14 patients). A significant proportion (56.7 %) was polymicrobial and 16.2 % involved suspected resistant pathogens. Sixteen patients recovered (5 on tigecycline alone, 5 with additional antimicrobials, and six switched to alternatives) while 21 patients died (nine on tigecycline alone, 12 with additional antimicrobials).

Conclusions: The study revealed that tigecycline prescription was conducted according to marketing authorizations and national guidelines. Infection severity/stage and comorbidities may influence patients' response, and explain some of the poor outcomes.

Keywords: Kingdom of Saudi Arabia, Prescription patterns, Mortality, Tigecycline, Antimicrobial

Tropical Journal of Pharmaceutical Research is indexed by Science Citation Index (SciSearch), Scopus, International Pharmaceutical Abstract, Chemical Abstracts, Embase, Index Copernicus, EBSCO, African Index Medicus, JournalSeek, Journal Citation Reports/Science Edition, Directory of Open Access Journals (DOAJ), African Journal Online, Bioline International, Open-J-Gate and Pharmacy Abstracts

INTRODUCTION

Bacterial infections present an increasing and well-recognized challenge to clinical practice for several reasons: the rapid emergence of resistant strains [1]; medication over-prescription in all healthcare settings [2,3]; and ethical issues,

for example inappropriate patient demands [4]. Furthermore, the empirical antibacterial therapy without sufficient intelligence in relation to pathogen susceptibility has been linked to increased fatality in patients with sepsis [5]. The discovery and development of truly novel anti-bacterials by the pharmaceutical industry has

declined markedly, primarily due to scientific complexity and resource implications [6]. Tigecycline is one of the very few unique agents launched in the last decade, with initial introduction in the United States (US) in 2005 [7]. The US Food and Drug Administration (FDA) licensed tigecycline for the treatment of patients with skin and soft tissue infections, complicated intra-abdominal infections, and community-acquired pneumonia [8]; followed by the European Medicine Agency (EMA) licensing in 2006 [9].

Tigecycline circumvents mechanisms of bacterial resistance via a modification of the central structure of minocycline by the addition of a 9-*t*-butyl-glycylamido side chain. This renders tigecycline effective against a wide range of gram-positive and gram-negative organisms, notably highly resistant pathogens such as methicillin-resistant *Staphylococcus aureus*, *Staphylococcus epidermidis*, vancomycin-resistant *Enterococcus* species, extended spectrum β -lactamase producers (ESBL), *Clostridium difficile* and many carbapenemase-producing Enterobacteriaceae; however, it has no activity against *Proteus* or *Pseudomonas* species [10,11]. It exhibits a linear pharmacokinetic profile, a large volume of distribution V_d (7 - 10 L), a half-life of 37 - 67 h, and clearance of 0.2 - 0.3 L/h/kg [12,13]. In adults, it is administered intravenously as a loading dose of 100 mg, followed by 50 mg every 12 h, with dose-reduction recommended in cases of severe hepatic impairment as in Child-Pugh Class C. Adverse drug reactions include: nausea (26 %), vomiting (18 %), diarrhea (12 %), abdominal pain (6 %), headache (6 %), and increased serum glutamic pyruvic transaminase levels (< 2 %) [14].

Two main concerns of tigecycline have become apparent recently: the emerging clinical resistance [15] and life-threatening adverse events [16] which have been confirmed by independent meta-analyses of all-cause mortality data [17,18]. Studies conducted in Europe [19] and Taiwan [20] have demonstrated largely, an appropriate use of tigecycline in terms of its licensed indications and dosage, highlighting the need to balance patients' risks and benefits.

Tigecycline was introduced to the Saudi market in 2008 to treat skin, soft-tissue, and intra-abdominal infections. In August, 2011, the Saudi Food and Drug Authority (SFDA) issued a safety notification advising that tigecycline should only be prescribed for licensed indications and as a last resort, where alternative antimicrobials were deemed inappropriate [21].

METHODS

Study design

A retrospective observational study was conducted in several hospitals. This study was approved by the Ethical Committee of the Ministry of Health Research Centre in Jeddah, KSA. All the three hospitals must comply with local prescription guidelines for tigecycline: the prescription should be initiated by a medical consultant; pathogen susceptibilities should be determined by laboratory analyses; and the prescription should be authorized by an infectious diseases physician, a microbiologist, and a pharmacist.

Study setting and participants

The study was conducted in three main KSA governmental hospitals: King Fahad Hofuf Hospital (KFHH); King Fahad Jeddah Hospital (KFJH); and King Fahad Specialist Hospital (KFSH), located in the regional provinces of Makkah, Al-Ahsa, and Qassim, respectively. Each hospital has 500-900 beds and mainly serves adult patients. All patients prescribed tigecycline between January 2013 and May 2014 were included in the study, with no exclusions.

Data collection and analysis

The patients prescribed tigecycline were identified from electronic prescription data. A data collection template was developed and piloted to record the following: patient demographics (age, sex, weight, hospital); past medical history; adverse drug reactions, allergies, medication history (including current medication); reason for admission to hospital (signs, symptoms); Acute Physiology and Chronic Health Evaluation (APACHE II) disease severity score [22]; tigecycline indication and regimen (dose, route, duration); documented adverse events; renal and hepatic function; microbiological analyses of pathogen susceptibility; ongoing monitoring (documented clinical response); and clinical safety endpoints (documented recovery, surgery, readmission). The data were entered into SPSS (SPSS Inc., Cary, NC version 22.0) and analyzed using descriptive statistics. Missing data were recorded for all variables considered. Three hospital pharmacists attended researcher training sessions to receive information about the purpose of the research and the data collection process. The data were collected from the patients' electronic medical records in KFSH and KFHH, and from paper-based medical records in

KFJH, where electronic records had not been fully implemented at the time of this study.

RESULTS

Thirty-seven patients were included: 18 in KFHH; 11 in KFJH; and 8 in KFBH. The mean patient

age was 52.5 years (SD \pm 21.4, range 17–92) and 19 (51.4 %) were female. The demographics at the start of tigecycline therapy, disease severity scores, and comorbidities, as shown in Table 1. Thirty-two patients (86.5 %) were

Table 1: Summary of patients' characteristics and reasons for tigecycline use

Characteristics	All patients	Improved^b	Died^b
Number of patients	37 (100%)	16 (43.2%)	21 (56.8%)
Mean Age (SD)	52.5 (21.4)	56.18 (16.37)	52.7 (24.01)
Women, n (%)	19 (51.4%)	5 (26.3%)	14 (73.7%)
<i>Comorbidities, n (%)</i>			
HTN	17 (17.3%)	4 (23.5%)	13 (76.5%)
DM	16 (16.3%)	6 (37.5%)	10 (62.5%)
Heart (CHF, IHD, MVS, MVR, LVH, LBBB)	13 (13.3%)	3 (23.1%)	10 (76.9%)
Renal (ARF, CKD, ESRD, other)	12 (12.3%)	3 (25%)	9 (75%)
Old CVA	6 (6.2%)	1 (16.7%)	5 (83.3%)
Other (amputated leg, bed-ridden, psychosis, piles, thrombocytopenia, warfarin overdose)	6 (6.2%)	2 (33.3%)	4 (66.6%)
Neurological (brain atrophy, multiple brain contusions, PD, CP)	4 (4.1%)	2 (50%)	2 (50%)
ASP	4 (4.1%)	1 (25%)	3 (75%)
Epilepsy	3 (3.1%)	2 (66.6%)	1 (33.3%)
Respiratory (asthma, COPD, IPF)	3 (3.1%)	1 (33.3%)	2 (66.6%)
PE	2 (2.0%)	-	2 (100%)
MR	2 (2.0%)	-	2 (100%)
SCA	2 (2.0%)	-	2 (100%)
Muscular (DMD, rhabdomyolysis)	2 (2.0%)	1 (50%)	1 (50%)
Autoimmune/inflammatory (RA, SLE)	2 (2.0%)	1 (50%)	1 (50%)
Infectious (TB, HCV)	2 (2.0%)	-	2 (100%)
Vascular (DIC)	1 (1.0%)	1 (100%)	-
Cancer	1 (1.0%)	1 (100%)	-
<i>Severity/organ dysfunction scores</i>			
APACHE II score, n	43	11.3	31.7
mean \pm standard deviation (range)	16.8 \pm 4.3 (0-28)	8.7 \pm 5.5 (0-11)	22.1 \pm 7.6 (4-28)
\leq 15, n (%)	11 (25.6)	7 (61.9)	5 (15.8)
\leq 15, mean (median)	7 (8.8)	12.4 (13.7)	11.6 (14.3)
$>$ 15, n (%)	23 (67.6)	3 (8.8)	20 (58.8)
$>$ 15, mean (median)	8.3 (9.7)	11.1 (12.6)	16.1 (18.4)
missing/unknown, n	3	1	2
<i>Reasons for tigecycline use, alone or in combination, in all patients</i>			
Reason, n (%) ^c			
Failure of previous therapy ^d	5 (13.5%)	1 (20%)	4 (80%)
Suspected resistant pathogens	6 (16.2%)	3 (50%)	3 (50%)
Need broad-spectrum coverage/polymicrobial infection	21 (56.7%)	9 (42.9%)	12 (57.1%)
Allergy to/intolerance of previous antibacterial	2 (5.4%)	1 (50%)	1 (50%)
Renal impairment	3 (8.1%)	2 (66.7%)	1 (33.3%)
Others	NA	-	-

Abbreviations: APACHE II, Acute Physiology and Chronic Health Evaluation score; ARF, acute renal failure; ASP, aspiration pneumonia; CHF, congestive heart failure; CKD, chronic kidney disease; COPD, chronic obstructive pulmonary disease; CP, cerebral palsy; DIC, disseminated intravascular coagulopathy; DM, diabetes mellitus; DMD, Duchenne muscular dystrophy; ESRD, end-stage renal disease; HTN, hypertension; HCV, hepatitis C virus; IHD, ischemic heart disease; IPF, interstitial pulmonary fibrosis; LBBB, left bundle branch block; LVH, left ventricular hypertrophy; MR, mental retardation; MVR, mitral valve stenosis; MVS, mitral valve regurgitation; NA, not available; PD, Parkinson's disease; PE, pulmonary edema; RA, rheumatoid arthritis; SCA, sickle-cell disease; SLE, systemic lupus erythematosus; TB, tuberculosis. ^aImprovement was defined as a clinical cure or response without additional antibiotic. ^bDeath was defined as failure or no improvement with additional antibiotic. ^cPatients could have more than one reason. ^dPrevious therapy included all treatments that were given prior to tigecycline

Use of tigecycline in 3 KSA hospitals and the treatment outcome




Figure 1: Treatment and outcomes of patients prescribed tigecycline (n = 37)

Table 2: Tigecycline indications (n = 37)

Indication	% patients
Sepsis	59.5
Community-acquired bacterial pneumonia	21.6
Intra-abdominal infection	13.5
Suspected community-acquired bacterial pneumonia	10.8
Wound infections (abscess, gangrene, diabetic foot and bed sores)	10.8
Urinary tract infection	2.7
Cystic fibrosis	2.7

Table 3: Identified bacterial pathogens (n = 74)

Gram-positive pathogens	n
<i>Enterococcus faecalis</i>	3
<i>Staphylococcus aureus</i>	1
<i>Staphylococcus epidermidis</i>	1
Methicillin-resistant <i>Staphylococcus aureus</i>	1
<i>Streptococcus anginosus</i>	1
<i>Staphylococcus saprophyticus</i>	1
<i>Viridans streptococci</i>	1
Gram-negative pathogens	
<i>Acinetobacter baumannii</i>	20, 3 resistant
<i>Klebsiella pneumonia</i>	14
<i>Pseudomonas aeruginosa</i>	11, 2 resistant
<i>Escherichia coli</i>	4, 2 resistant
<i>Proteus mirabilis</i>	4
ESBL-producing <i>Enterobacteriaceae</i>	3
<i>Serratia marcescens</i>	2
<i>Providencia stuartii</i>	1
<i>Morganella morganii</i>	1
ESBL-producing <i>Klebsiella pneumoniae</i>	1
<i>Klebsiella oxytoca</i>	1
<i>Enterobacter cloacae</i>	1
<i>Pseudomonas luteola</i>	1
<i>Citrobacter freundii</i>	1

treated with tigecycline in the intensive care unit (ICU) and the remaining proportion of patients were in the medical wards. Biochemical monitoring prior to commencing tigecycline, identified three patients with impaired renal function; there was no evidence of impaired hepatic function.

The documented indications for tigecycline are given in Table 2, with some patients having more than one indication. None of the patients had been treated previously for similar infections.

DISCUSSION

Tigecycline is a very few unique antimicrobial agents which was launched in the last decade, with initial introduction to the US in 2005. Two main concerns of tigecycline have become apparent since its launch; emerging clinical resistance and life-threatening adverse events. Tigecycline was introduced to the KSA market in 2008.

This study showed that tigecycline prescription patterns within these hospitals in the KSA were mostly in line with the guidelines [7,9]; it was reserved and prescribed for clearly defined indications at appropriate initiation and maintenance doses [7,11,22]. This is in line with the previous studies in Taiwan [19] and Europe [20]. Tigecycline was initiated under the instruction of a consultant in according to the recommended dosing schedule of 100 mg intravenously initially, followed by 50 mg every

12 hours (except one patient with renal insufficiency received 50 mg followed by 25 mg every 12 h). The treatments and patients' outcomes are shown in Figure 1.

However, we identified tigecycline usage for some unapproved indications that both the SFDA and FDA have issued warnings against; these included diabetic foot, urinary tract infections, undefined pneumonia, and septic shock.

These dangerous pathogens are acknowledged to pose the greatest therapeutic challenges and, while often drug-resistant, may respond to tigecycline [23,24]. Other studies in the KSA reported that 20 % of *Acinetobacter baumannii* cases showed an increased resistance in 2011, as compared to 10 % in 2010 [25,26]. A range of pathogens were isolated, reflecting the complexity of the required therapeutic plans. Interestingly, tigecycline was also effective in patients infected with *Serratia marcescens*, previously reported to be resistant to tigecycline [27], and similarly in a patient infected with *Klebsiella pneumoniae* [28,29].

Of the 37 patients prescribed tigecycline, 16 recovered (five on tigecycline alone, five with additional antimicrobials, and six who switched to alternatives) and 21 died, 14 were confirmed with *Klebsiella pneumoniae*, (nine on tigecycline alone, twelve with additional antimicrobials). In 2011, the SFDA warned of a high death risk if tigecycline is used for pneumonia, or any unapproved indications [21]. A further

confirmation by the FDA was released in 2013 warning against the increased risk of death in approved and unapproved indications, stating that tigecycline should only be used when necessary [16]. Ten out of the 16 patients, continued on tigecycline therapy, and the other 6 were changed to other antibiotics because tigecycline therapy had failed and/or resistant pathogens were present. In the ten recovered patients who continued on tigecycline, improvement was documented as occurring within two to seven days, with full recovery observed between six and 22 days.

Tigecycline's adverse events include: nausea and vomiting (n = 4); diarrhea (n = 2); and hypotension (n = 1). These mostly appeared on the first or second day of tigecycline treatment. None of these adverse drug reactions led to tigecycline discontinuation or treatment change. Patients' clinical outcomes varied and were extremely difficult to associate with tigecycline efficacy because of the influences of confounding factors, including severity and stage of infection, comorbidities, and concurrent antimicrobial therapy. The documented adverse events were

Table 4: Indication, treatment, and outcomes of patients commenced and continued on tigecycline (n = 10)

Documented indication(s)	Pathogen(s)	Antimicrobials added to tigecycline	Days to documented symptom improvement	Days to documented recovery
Severe pneumonia	<i>Klebsiella pneumoniae</i>	Vancomycin, amikacin, meropenem, ribavirin, and interferon α -2an	5	7
Intra-abdominal infection	<i>Acinetobacter baumannii</i> , <i>Escherichia coli</i>	Meropenem and caspofungin	6	22
Septic shock, bilateral pneumonia	<i>Serratia marcescens</i>	No	7	14
Acute respiratory failure with CO ₂ narcosis	Multi-drug resistant <i>Acinetobacter baumannii</i>	No	7	14
Interstitial pulmonary fibrosis, septic shock	<i>Proteus mirabilis</i> , <i>Acinetobacter baumannii</i> , <i>Pseudomonas aeruginosa</i>	No	5	8
Pneumonia	<i>Providencia stuartii</i> , <i>Serratia marcescens</i>	No	5	8
Diabetic foot infection	<i>Acinetobacter baumannii</i> , <i>Escherichia coli</i>	Meropenem and metronidazole	2	22
Fever, malignant neoplasm of the nasopharynx	Coagulase-negative <i>Staphylococci</i> , non-albicans <i>Candida</i>	Colistin	3	6
Urinary tract infection, bed sores	<i>Morganella morganii</i> , <i>Klebsiella oxytoca</i>	Amikacin	5	12
Septic shock, catheter-related	Multi-drug resistant <i>Acinetobacter baumannii</i>	No	4	7

of low prevalence and severity, and were similar to those reported previously [6,14]. While life-threatening adverse events have been noted with tigecycline [16,18] and have resulted in safety warnings, it is unlikely that the high mortality observed in this study was directly attributable to tigecycline. Many patients included in this study died, irrespective of treatment, and few patients continued on tigecycline as the sole antimicrobial.

Limitations of the study

The small dataset and the potential reporting bias due to the reliance on case notes; are the major two limitations. It is possible that drug adverse events were underreported. Even when tigecycline is mostly prescribed in line with the relevant marketing authorizations, patient outcomes are complex and influenced by many confounding factors and may not necessarily be generalizable to the entire KSA.

CONCLUSION

The study revealed that tigecycline prescription was in line with the marketing authorizations and national guidelines. While many patients had poor outcomes, these could not be attributed solely to a lack of tigecycline efficacy or toxicity; confounding factors, including infection severity/stage and comorbidities may influence patients' outcomes.

ACKNOWLEDGEMENT

The authors would like to acknowledge the Ministry of Health and the Directors of the Directorate of Health Affairs in KSA for supporting this research. We would also like to extend our thanks to all the participating hospitals.

REFERENCES

- Davies J, Dorothy D. Origins and evolution of antibiotic resistance. *Microbiol Mol Biol Rev* 2010; 74(3): 417-433.
- Costelloe C, Metcalfe C, Lovering A, Mant D, Hay AD. Effect of antibiotic prescription in primary care on antimicrobial resistance in individual patients: systematic review and meta-analysis. *Brit Med J* 2010; 340: c2096.
- Davey P, Brown E, Charani E, Fenelon L, Gould IM, Holmes A, Ramsay CR, Wiffen PJ, Wilcox M. Interventions to improve antibiotic prescription practices for hospital inpatients. *Cochrane Database Syst Rev* 2013; 4: CD003543.
- Leibovici L, Paul M, Ezra O. Ethical dilemmas in antibiotic treatment. *J Antimicrob Chemother* 2012; 67: 12-16.
- Paul M, Shani V, Muchtar E, Kariv G, Robenshtok E, Leibovici L. Systematic review and meta-analysis of the efficacy of appropriate empiric antibiotic therapy for sepsis. *Antimicrob Agents Chemother* 2010; 54: 4851-4863.
- Fischbach MA, Walsh CT. Antibiotics for emerging pathogens. *Science* 2009; 325: 1089-1093.
- Food and Drug Administration. FDA Approved Drug Products. Available at: <http://www.accessdata.fda.gov/scripts/cder/drugsatfda/index.cfm?fuseaction=Search.DrugDetails> [Accessed Aug 2014].
- Stein, Gary E., and William A. Craig. "Tigecycline: a critical analysis." *Clin infect dis* 2006; 43.4 : 518-524.
- Dryden M. Tigecycline: an antibiotic for the twenty-first century. *J Antimicrob Chemother* 2013; 68(suppl 2): ii3-ii4.
- Brust K, Evans A, Plemmons R. Tigecycline in treatment of multidrug-resistant Gram-negative bacillus urinary tract infections: a systematic review. *J Antimicrob Chemother* 2014; 69(10): 2606-2610.
- Tasina E, Haidich AB, Kokkali S, Arvanitidou M. Efficacy and safety of tigecycline for the treatment of infectious diseases: a meta-analysis. *Lancet Infect Dis* 2011; 11: 834-844.
- Muralidharan G, Micalizzi M, Speth J, Raible D, Troy S. Pharmacokinetics of tigecycline after single and multiple doses in healthy subjects. *Antimicrob Agents Chemother* 2005; 49(1): 220-229.
- MacGowan AP. Tigecycline pharmacokinetic/pharmacodynamic update. *J Antimicrob Chemother* 2008; 62(suppl 1): i11-i16.
- Cai Y, Wang R, Liang B, Bai N, Liu Y. Systematic review and meta-analysis of the effectiveness and safety of tigecycline for treatment of infectious disease. *Antimicrob Agents Chemother* 2011; 55(3): 1162-1172.
- Sun Y, Cai Y, Liu X, Bai N, Liang B, Wang R. The emergence of clinical resistance to tigecycline. *Int J Antimicrob Agents* 2013; 41(2): 110-116.
- Food and Drugs Administration. Drug Safety Communication 2013. Increased risk of death with Tygacil (tigecycline) compared to other antibiotics used to treat similar infections. Available at: <http://www.fda.gov/Drugs/DrugSafety/ucm369580.htm> [Accessed Aug 2014].
- Yahav D, Lador A, Paul M, Leibovici L. Efficacy and safety of tigecycline: a systematic review and meta-analysis. *J Antimicrob Chemother* 2011; 66: 1963-1971.
- Prasad P, Sun J, Danner RL, Natanson C. Excess deaths associated with tigecycline after approval based on noninferiority trials. *Clin Infect Dis* 2012; 54(12): 1699-1709.
- Kuo SC, Wang FD, Fung CP, Chen LY, Chen SJ, Chiang MC, Hsu SH, Liu CY. Clinical experience with tigecycline as treatment for serious infections in elderly and critically ill patients. *J Microbiol Immunol Infect* 2011; 44(1): 45-51.
- Bassetti M, Eckmann C, Bodmann KF, Dupont H, Heizmann WR, Montravers P, Guirao X, Capparella MR, *Trop J Pharm Res*, October 2015; 14(10): 1925

- Simoneau D, García MS. Prescription behaviours for tigecycline in real-life clinical practice from five European observational studies. *J Antimicrob Chemother* 2013; 68(suppl 2): ii5-ii14.
21. Kingdom of Saudi Arabia Food & Drug Authority. Drug Bulletin 2011; 2(3) Available at: <http://www.sfda.gov.sa/ar/drug/resources/DocLib1/Volume%202/Issue3Hihgh.pdf>. [accessed Aug 2014].
 22. Knaus WA, Draper EA, Wagner DP, Zimmerman JE. "APACHE II: a severity of disease classification system". *Critical Care Medicine* 1985; 13 (10): 818–29.
 23. Shin JA, Chang YS, Kim HJ, Kim SK, Chang J, Ahn CM, Byun MK. Clinical outcomes of tigecycline in the treatment of multidrug-resistant *Acinetobacter baumannii* infection. *Yonsei Med J* 2012; 53(5): 974-84.
 24. Karageorgopoulos DE, Kelesedis T, Kelesedis I, Falagas ME. Tigecycline for the treatment of multidrug-resistant (including carbapenem resistant) *Acinetobacter* infections: a review of the scientific evidence. *J Antimicrob Chemother* 2008; 62: 45-55.
 25. Baadani AM, Thawadi SI, El-Khizzi NA, Omrani AS. Prevalence of colistin and tigecycline resistance in *Acinetobacter baumannii* clinical isolates from 2 hospitals in Riyadh Region over a 2-year period. *Saudi Med J* 2013; 34(3): 248-253.
 26. Al-Qadheeb NS, Althawadi S, Alkhalaf A, Hosaini S, Alrajhi AA. Evolution of tigecycline resistance in *Klebsiella pneumoniae* in a single patient. *Ann Saudi Med* 2010; 30(5): 404-407.
 27. Hornsey M, Ellington MJ, Doumith M, Hudson S, Livermore DM, Woodford N. Tigecycline resistance in *Serratia marcescens* associated with up-regulation of the SdeXY-HasF efflux system also active against ciprofloxacin and cefpirome. *J Antimicrob Chemother* 2010; 65(3): 479-482.
 28. Ruzin A, Visalli MA, Keeney D, Bradford PA. Influence of transcriptional activator RamA on expression of multidrug efflux pump AcrAB and tigecycline susceptibility in *Klebsiella pneumoniae*. *Antimicrob Agents Chemother* 2005; 49: 1017-1022.
 29. Roy S, Datta S, Viswanathan R, Singh AK, Basu S. Tigecycline susceptibility in *Klebsiella pneumoniae* and *Escherichia coli* causing neonatal septicaemia (2007–2010) and role of an efflux pump in tigecycline non-susceptibility. *J Antimicrob Chemother* 2013; 68(5): 1036-1042.