

Diversity and Stability of Demersal Species Assemblages in the Gulf of Guinea

K. A. Koranteng

Marine Fisheries Research Division, P.O. Box BT-62, Tema, Ghana

(Email: kwamek@africaonline.com.gh)

Abstract

The structure of demersal fish assemblages on the continental shelf and upper continental slope of the Gulf of Guinea is described. Community structure is determined primarily by depth and type of sediment on the seabed. Changes in the composition of the identified species assemblages over a 25-year period are examined. The dynamics of the assemblages are influenced by physico-chemical parameters of the water masses, mainly temperature, salinity and dissolved oxygen, which are periodically modified by the seasonal coastal upwelling that occurs in the western Gulf of Guinea. Increased industrial trawling in coastal waters and environmental forcing jointly influenced the changes in the composition and relative importance of species in the assemblages. Irresponsible fishing operations (like the use of explosives) that lead to habitat alteration and other anthropogenic activities (like oil and gas exploration) which have the potential to cause environmental changes pose a threat to biodiversity in continental shelf waters of the Gulf of Guinea.

Introduction

Knowledge of species aggregation and diversity is an important tenet of the ecosystem management concept and approach that are receiving favourable consideration in recent times in the management of living natural resources. In fisheries, defining the aggregation of species in the ecosystem is the basis for managing species by the management unit approach (Tyler, Gabriel & Overholtz, 1982).

The structure of species assemblages has been established for several exploited fish populations around the world. In the Gulf of Guinea, Fager & Longhurst (1968) and Longhurst (1969) worked out the assemblage structure of demersal species on the continental shelf using data from the Guinean Trawling Survey of 1963-64 (Williams, 1968). Villegas & Garcia (1983) mapped demersal fish communities on the continental shelf of selected countries in the sub-region using data and information from a number of trawl surveys. Assemblage is

defined here as "an association of coexisting species with similar environmental tolerance, possibly trophic relationships, but not totally interdependent" (Bianchi, 1992).

In the last three decades, significant changes have occurred in the biological and physical components of the Gulf of Guinea marine ecosystem and in nearshore forcing factors that could have affected species associations and status of marine biodiversity in the sub-region (Koranteng, 1998). Koranteng (1998) showed that the period between 1963 and 1992 could be broken down into three climatic periods each of which had distinct environmental characteristics in the Gulf of Guinea. In the first period (i.e. 1963-1972), sea temperatures (surface and bottom) were relatively high, salinity was low and the thermocline was below its long-term average depth. Between 1972 and 1982 (the second climatic period), there was a global decline in sea temperatures and a rise in salinity. In the final period (1982-1992), temperatures

were high and salinity was low and erratic.

The seasonal coastal upwelling of cold, nutrient rich, subthermocline water drives the biology of the continental shelf waters of the Gulf of Guinea, especially off Ghana and La Côte d'Ivoire. The duration and intensity of the major upwelling (July-September) and minor upwelling (December-February) that occur annually in the sub-region have changed (Pezennec, 1995, Koranteng & Pezennec, 1998).

The building of dams on almost all major rivers in the sub-region for hydroelectric and irrigation purposes and the sub-Saharan drought in the 1970s and 1980s have resulted in reduced river discharge and sediment transport into the Gulf of Guinea (Lamb, 1982; Mahé, 1998).

Furthermore, siltation and draining of many coastal lagoons in the sub-region as a result of urbanisation and depletion of the mangrove cover around these lagoons have undermined their (the lagoons) contribution to the continental shelf ecosystem (Entsua-Mensah, 1998). Increased industrial fishing, especially in coastal waters, the use of fishing nets with small-sized meshes and other inappropriate fishing methods like dynamite and other explosives, have led to the decline in catch rates of many commercially exploited fish species and virtual extinction of others (Koranteng, 1998). According to Longhurst & Pauly (1987) these are some of the principal factors that cause ecological changes in tropical fish communities. Throughout the sub-region, oil and gas exploration and prospecting have been intensified in the last five decades or so. Large-scale drilling for oil is being undertaken on the continental shelf and in the delta areas and, occasionally, oil spills occur in the area from these activities

(Saenger *et al.*, 1998; Shell, 1998).

The most important recorded changes in fish population in the Gulf of Guinea are the decline and subsequent recovery of the round sardinella (*Sardinella aurita*) populations (Pezennec, 1995), the proliferation and subsequent decline of triggerfish (*Balistes capriscus*) (Koranteng, 1984, Caverivière, 1991) and increase in abundance of cuttlefish (*Sepia officinalis*) and globefish (*Lagocephalus laevigatus*) (Martos *et al.*, 1990; Koranteng, 1998). The proliferation of triggerfish in the Gulf of Guinea in the 1970s and its total domination of the ecosystem were described by Bakun (1995) as "one of the most phenomenal episodes in the history of fish population dynamics".

Studies of fish communities have shown that some of these natural and anthropogenic factors could induce changes in the structure of species assemblages, threaten the fish biodiversity and affect the general well-being of fishery resources (Brown *et al.*, 1976; Overholtz & Tyler, 1985; Greenstreet & Hall, 1996).

The primary objective of this study is to examine the structure of demersal species assemblages on the continental shelf and upper continental slope of the Gulf of Guinea and to examine the changes that have occurred in these assemblages in the last three decades. The study also seeks to isolate the factors that determine fish community structure in the study area and to assess the effects of environmental parameters and anthropogenic activities on the structure and dynamics of the assemblages.

Materials and methods

Between 1956 and 1992, a number of bottom

trawl surveys of the demersal fishery resources on the continental shelf and upper slope off Ghana were conducted (Koranteng, 1998). The Guinean Trawling Survey (GTS) organised by the Scientific, Technical and Research Commission of the Organisation of African Unity (OAU/STRC) in 1963-64 (Williams, 1968) was the first survey that covered the entire continental shelf of Ghana. From 1969 the Marine Fisheries Research Division (MFRD) of Ghana also conducted a number of bottom trawl surveys in Ghana's marine waters. The data from these surveys, made up of fish catches and measurements of some oceanographic parameters, were used in this study.

Each survey involved cruises in the upwelling and non-upwelling (or thermocline) seasons. The sampling procedures were almost identical in all the surveys. The catch from each haul was sorted according to species. Each species was weighed and the total number of individuals determined by total count or estimation from subsamples. During the trawl surveys, water temperature was measured using thermometers mounted on Nansen reversing bottles. Salinity was measured using a salinometer and dissolved oxygen by the Winkler titration method.

All catch data were inputted into the NAN-SIS computer program for trawl survey data logging and analysis (Stromme, 1992). The GTS catch data were subjected to multivariate statistical analyses to determine the groupings in each data set. A Two-Way Indicator Analysis method implemented by the TWINSpan computer program (Hill, 1979) was used. TWINSpan uses a divisive cluster analysis algorithm to classify the samples and correspondence analysis (CA) to perform an ordination which

is a multivariate statistical method of arranging samples and species in a two-dimensional graph in which similar samples or species are near each other and dissimilar entities are far apart. CA assumes that the response of species to environmental changes follows a unimodal Gaussian curve rather than the linear relationship assumed in other multivariate methods (Digby & Kempton, 1987; ter Braak, 1991; Jongman, ter Braak & van Tongeren, 1995). Pelagic species occurring in the trawls were included in the analysis as suggested by Longhurst & Pauly (1987) because several fish species classified as pelagic are also common in demersal trawls.

The community ecology program CANOCO (ter Braak, 1991) was used to assess the factors that influence the ordination. The hydrographic parameters used in the analysis are water temperature, salinity, and dissolved oxygen measured at depths trawled. Other environmental parameters are the depth sampled and type of bottom sediment at the stations trawled. Bottom type information was obtained from sediment maps produced during GTS and also from Martos *et al.* (1990). Bottom type was classified as follows: Hard (predominantly sand, shell, rock, gravel, grit or coral), Soft (predominantly mud) and Mixed (combination of hard and soft). The three bottom types were treated as three levels of a nominal variable (bottom type). The values of a nominal variable only represent the names of the levels and do not imply any order or magnitude of differences between them.

CANOCO performs a Detrended Correspondence Analysis (DCA) which is an improvement on CA and addresses its defects. The influence of each environmental

parameter on the ordination, hence assemblage structure was assessed through the significance of the correlation between the environmental parameter and the ordination axes. The DCA scores were plotted using the CANODRAW computer program (Smilauer, 1992).

To determine the most important species in each assemblage, the index of relative importance (IRI), modified by Pinkas, Oliphant & Iverson (1971) was used. The index is expressed as:

$$IRI = \%W \times \%F$$

where %W is the percentage composition by weight of each species in the assemblage and %F is the percentage frequency of occurrence of the species in hauls from the assemblage. Species with IRI values of 50 or more in each assemblage were considered as the most important species in the assemblage.

Published materials on species assemblages in other parts of the Gulf of Guinea, arising from surveys conducted in the sub-region, were examined with a view to comparing the assemblage structures derived from these surveys and the results of this work. The surveys included those conducted off La Côte d'Ivoire (Joanny & Menard, 1998), Togo and Benin (Crosnier & Berrit, 1966), Nigeria (Longhurst, 1964) and the Guinean Trawling Survey (GTS) conducted in the entire waters of the Gulf of Guinea (Williams, 1968).

The data sets from the GTS (1963-64) and MFRD surveys of 1981-82 (MFRD3) and 1989 (MFRD5) were used in the investigation on possible changes in demersal species assemblages on the continental shelf (depths down to 200 m) and upper continental slope (200 to about 600 m). The GTS was conducted during the first climatic

period described above, MFRD3 and MFRD5 were conducted during the second and third climatic periods respectively. The two MFRD surveys were similar in terms of spatial and temporal coverage and in survey methods. The GTS was the only survey of similar spatial coverage conducted in Ghanaian waters during the first climatic period. To obtain data comparable with those of the MFRD surveys, a subset of the GTS data from depths shallower than 100 m were re-analysed.

The first three assemblages on the continental shelf were used to examine fish community dynamics and stability and species diversity in the assemblages. The investigations centred on the nature and pattern of change and the possible factors and causes. For each survey and assemblage, a list of species with IRI > 50 was produced for the upwelling and thermocline seasons to determine the species representation at these periods. The nature and extent of changes at the community level were examined by delineating three depth ranges and following community succession within each depth range over time. The depth ranges are 10-30 m, 31-50 m and 51-100 m. For this analysis, data from trawl surveys conducted in Ghanaian waters in 1963/64, 1969/70, 1973-77 and 1979-90 were used.

In each depth range and climatic period, the following indices were calculated:

Shannon Diversity Index,

$$H' = -\sum_{i=1}^s \left(\frac{w_i}{w} \right) \ln \left(\frac{w_i}{w} \right),$$

Margelef's Richness Index,

$$R = \frac{s-1}{\ln(w)}, \text{ and}$$

$$\text{Pielou's Evenness Index } J' = \frac{H''}{\ln(s)}$$

where w_i is the sample weight of the i^{th} species, w is total weight and s is the number of species in a sample. Weights of the species were used in the calculations instead of numbers following the suggestion of Willhm (1968).

The reaction of important species groups to the observed environmental changes was examined. The average density (kg/ha) of each species group in each depth range and climatic period was calculated.

Results

Six species groups (or assemblages) were obtained from the GTS data sets. These groups, corresponding to those named by Longhurst (1969) are:

- Group 1: Sciaenid community
- Group 2: Lutjanid community
- Group 3: Sparid community (shallow part)
- Group 4: Sparid community (deep part)
- Group 5: Deep shelf community

Group 6: Upper slope community. Eurybathic or thermocline species (Longhurst, 1969) were not isolated in this work as they are generally part of the second assemblage. Also, the estuarine sciaenid community described by Longhurst (1969) was not adequately sampled in the GTS survey.

Fig.1 shows a CANODRAW bi-plot of sites and environmental parameters in the DCA axis 1 / axis 2 plane. Hauls from the same group (assemblage) are indicated by the same symbol and enclosed in an ellipse. Table 1 gives Pearson's product-moment correlation coefficient r of species ordination axes 1 and 2 with bottom environmental variables. The significantly high correlation between depth, temperature, salinity and dissolved oxygen and the first two DCA axes show the importance of these parameters in the determination of the structure of the demersal species assemblages in the study area. These physico-chemical parameters are themselves closely related to depth in the oceans and usually changed by seasons




Fig. 1. Plot of DCA axis 1 versus axis 2 for the Guinean Trawl Survey data. Co-ordinates of environmental parameters in the ordination plane are also shown (with original scale reduced by 1/5)

TABLE 1

Pearson's product-moment correlation coefficient r of species ordination axes 1 and 2 with bottom environmental variables

Survey	Variable	Species axis 1	Species axis 2	Survey	Variable	Species axis 1	Species axis 2
GTS 1963/64 (all hauls)	Hard	-0.13	-0.29**	1981/82	Hard	-0.49***	0.09
	Mixed	0.13	-0.14		Mixed	0.30**	0.01
	Soft	-0.02	0.43**		Soft	0.30**	0.13
	Depth	0.95**	0.01		Depth	-0.29**	0.79***
	Temperature	-0.81**	0.10		Temperature	0.23*	-0.65***
	Salinity	-0.64**	-0.01		Salinity	-0.02	0.28
	Oxygen	-0.34**	0.11		Oxygen	0.13	-0.54***
	Eigenvalues	0.68	0.50		Eigenvalues	0.40	0.34
GTS 1963/64 (hauls at £ 100 m)	Hard	-0.31**	-0.02	1989	Hard	-0.29**	-0.30**
	Mixed	-0.11	-0.29*		Mixed	0.16	0.12
	Soft	0.44**	0.37**		Soft	0.20*	0.26**
	Depth	-0.51***	-0.71***		Depth	-0.42***	0.60***
	Temperature	0.47***	-0.47***		Temperature	0.18	-0.54***
	Salinity	0.18*	-0.07		Salinity	0.08	-0.07
	Oxygen	0.21*	-0.19		Oxygen	-0.02	-0.47***
	Eigenvalues	0.54	0.40		Eigenvalues	0.46	0.36

r significant at * 5% level, ** at 1% level, *** at 0.1% level.

(Riley & Chester, 1971, Mann & Lazier, 1996).

The number of fish species caught during each cruise of the three surveys and classified according to assemblage, is given

in Table 2. The number varied between 47 in the first assemblage obtained from the data of the May 1982 cruise and 150 for the second assemblage of the September cruise of the GTS (1963). As an example of

TABLE 2

Number of species and species groups recorded during surveys in thermocline and upwelling seasons. Species are not limited exclusively to one assemblage or the other

Survey	Season	Number of species in assemblage		
		Sciaenid	Lutjanid	Sparid
1963/64	Upwelling	110	150	93
	Thermocline	143	110	83
1981/82	Upwelling	81	51	48
	Thermocline	47	65	77
1989	Upwelling	69	80	63
	Thermocline	55	74	51

differences that occur in species abundance during the two seasons, the IRI values greater than 50 in the sciaenid assemblage derived from the data in the three surveys (GTS - 1963/64, MFRD3 - 1981/82 and MFRD5 - 1989) are presented in Table 3. The Table shows species that are present

mainly during the upwelling season, those in the thermocline season and those that are available all the time in the assemblage (referred to as resident species).

The calculated Shannon Diversity Index (H'), Margalef's Richness Index (R) and Pielou's Evenness Index (J') for the three

TABLE 3
The Sciaenid assemblage in the Gulf of Guinea at different periods

1963/64	1981/82	1989
Upwelling season species	Upwelling season species	Upwelling season species
<i>Drepane africana</i>	<i>Decapterus rhonchus</i>	<i>Lagocephalus laevigatus</i>
<i>Epenephelus aeneus</i>	<i>Pagellus bellottii</i>	<i>Pagellus bellottii</i>
<i>Ilisha africana</i>	<i>Pomadasys incisus</i>	<i>Pentheroscion mbizi</i>
<i>Loligo</i> sp.	<i>Priacanthus arenatus</i>	<i>Pomadasys incisus</i>
<i>Pomadasys jubelini</i>	<i>Pseudupeneus prayensis</i>	<i>Priacanthus arenatus</i>
<i>Pseudolithus brachygnathus</i>	<i>Sepia</i> sp.	<i>Pteroscion peli</i>
<i>Pseudolithus typus</i>		<i>Rhizoprionodon acutus</i>
<i>Sparus caeruleostictus</i>		<i>Trachinocephalus myops</i>
		<i>Trachurus trecae</i>
Resident species	Resident species	Resident species
<i>Brachydeuterus auritus</i>	<i>Balistes capriscus</i>	<i>Brachydeuterus auritus</i>
<i>Galeoides decadactylus</i>	<i>Brachydeuterus auritus</i>	<i>Galeoides decadactylus</i>
<i>Pagellus bellottii</i>	<i>Chloroscombrus chrysurus</i>	<i>Penaeus notialis</i>
<i>Raja miraletus</i>	<i>Dentex canariensis</i>	<i>Pomadasys jubelini</i>
<i>Trichiurus lepturus</i>	<i>Epenephelus aeneus</i>	<i>Pseudupeneus prayensis</i>
<i>Pseudolithus senegalensis</i>	<i>Galeoides decadactylus</i>	<i>Sepia officinalis</i>
<i>Pteroscion peli</i>	<i>Selene dorsalis</i>	<i>Selene dorsalis</i>
	<i>Sparus caeruleostictus</i>	<i>Sparus caeruleostictus</i>
		<i>Trichiurus lepturus</i>
Thermocline season species	Thermocline season species	Thermocline season species
<i>Chilomycterus spinosus</i>	<i>Sphyraena</i> sp.	<i>Elops senegalensis</i>
	<i>Engraulis encrasicolus</i>	<i>Chloroscombrus chrysurus</i>
	<i>Ilisha africana</i>	<i>Dasyatis</i> sp.
	<i>Pseudolithus senegalensis</i>	<i>Dentex canariensis</i>
	<i>Pseudolithus</i> sp.	<i>Drepane africana</i>
	<i>Pteroscion peli</i>	<i>Elops senegalensis</i>
	<i>Scyacia micrurum</i>	<i>Epenephelus aeneus</i>
	<i>Sphyraena sphyraena</i>	<i>Eucinostomus melanopterus</i>
		<i>Grammolites gruvelli</i>
		<i>Lutjanus fulgens</i>
		<i>Sardinella maderensis</i>
		<i>Sphyraena sphyraena</i>
		<i>Torpedo</i> sp.

depth ranges and different climatic periods are presented in Table 4. The Table shows a decline in species diversity between the first and second climatic periods in the 10-30 m and 31-50 m depth ranges and a slight increase in the 51-100 m depth range. Species diversity increased in all three depth ranges in 1982-92. Species richness and evenness also showed some variation as indicated in Table 4.

The pie charts in Fig. 2 show percentage

composition by weight of three important groups of species (lutjanids, sciaenids, sparids) and triggerfish in the total weight of the four groups only. The other species caught in the survey were ignored in these comparisons. There is one pie chart for each climatic period and depth range.

Discussion

Structure of demersal fish communities in the Gulf of Guinea

The continental shelf of the Gulf of Guinea

TABLE 4

Calculated values of Shannon Diversity Index, Margalef's Richness Index and Pielou's Evenness Index at different depths and climatic periods

Period	Shannon Diversity Index				Margalef's Richness Index				Pielou's Evenness Index			
	10-30	51-50	51-100	Mean	10-30	51-50	51-100	Mean	10-30	51-50	51-100	Mean
1963-71	2.69	3.04	2.59	2.77	12.23	12.47	6.97	10.56	0.64	0.66	0.64	0.65
1972-82	2.16	2.03	2.63	2.27	8.14	8.77	8.47	8.46	0.54	0.47	0.62	0.54
1983-90	2.80	3.21	2.92	2.98	8.75	9.90	7.89	8.85	0.68	0.74	0.70	0.71




Fig. 2. Percentage composition of sciaenids, lutjanids, sparids and *B. capriscus* at three depth ranges and time periods

has distinct biotopes comprising mud, hard rocks and mixed deposits (Longhurst, 1964; Crosnier & Berrit, 1966; Williams, 1968; Martos et al., 1990). Off Ghana the shelf is characterised by a belt of soft, muddy substrate in shallow waters down to about 30 m depth, followed by a wide area of mixed to hard bottom type.

The demersal fauna is quite similar in the entire Gulf of Guinea (Williams, 1968; Longhurst, 1969),

however, the occurrence of isolated patches of rocky bottoms, estuaries and lagoons result in some differences in the distribution and abundance of demersal fish species. Similar species assemblages occur over similar biotopes and water depths throughout the region (Longhurst, 1964; Crosnier & Berrit, 1966; Longhurst, 1969; Williams, 1968; Villegas & Garcia, 1983; Longhurst & Pauly, 1987; Koranteng, 1998; Joanny & Menard, 1998). Longhurst (1969) noted the remarkable degree of agreement between the various authors of earlier works with regard to the organisation of the assemblages.

The six principal species assemblages were named by Longhurst (1969) as sciaenid, lutjanid, sparid (shallow component), sparid (deep component), deep shelf and upper slope assemblages. The species assemblages occurring throughout the Gulf of Guinea are described below.

- i. The sciaenid assemblage occurs in shallow-waters (15-50 m) on soft, sandy and muddy bottoms and close to estuaries. Very extensive sciaenid assemblages occur off La Côte d'Ivoire, eastern part of Ghana, Nigeria and Togo-Benin. The characteristic members include *Brachydeuterus auritus*, *Galeoides* spp., *Pseudolithus* spp. *Arius* spp., *Selene dorsalis* and *Ilisha africana*.
- ii. The lutjanid assemblage is usually found next to the sciaenid assemblage on mixed to hard bottoms. It could also occur close to shore, as is the case off Togo-Benin, or alternate with the sciaenid assemblage as the first assemblage encountered from shore (e.g. off Nigeria-Cameroon). The lutjanid assemblage is very

prominent off Liberia. Some of the important members of the assemblage are *Lethrinus atlanticus*, *Lutjanus* spp., *Acanthurus monroviae*, *Balistes forcipatus*, *Pagellus bellottii*, *Brachydeuterus auritus* and *Trachurus* sp.

- iii. The sparid assemblage comprises shallow and deep parts. The former is found within 15 and 70 m water depth on soft/mixed bottoms and extends into very shallow waters during some parts of the year. *Penaeus notialis*, *Brachydeuterus auritus*, *Balistes capriscus*, *Trachurus* sp., *Raja miraletus* and *Pagellus bellottii* are some of the indicator species of the shallow part. The deep part prefers mixed to hard bottoms and may be found in water depths down to 200 m. *Dentex* spp., *Boops boops*, *Squatina oculata*, *Trachurus* sp., *Scomber japonicus*, *Pagellus bellottii*, *Pseudupeneus prayensis*, *Epinephelus* spp. and *Lagocephalus laevigatus* represent this assemblage.
- iv. The deep continental shelf assemblage is found at water depths of 200 m or more on mixed/soft bottoms and have *Paracubiceps ledanoisi*, *Trachurus* sp., *Loligo* sp. as some of its important members.
- v. The upper continental slope assemblage usually found on mixed/soft bottoms at depths of 400 m or more has *Peristedion cataphractum*, *Antigonia capros*, *Chlorophthalmus* spp., *Epigonus* spp., *Trigla* spp., *Centrophorus uyato* as the representative species.

Variability and trends in fish community structure in the Gulf of Guinea

There were some differences in the composition of the assemblages between surveys and seasons. Certain species occurred repeatedly in some assemblages over the 25-year period examined in this work. These included *Brachydeuterus auritus*, *Galeoides decadactylus* and *Sparus caeruleostictus* in the sciaenid assemblage, *Dentex canariensis*, *Pagellus bellottii*, *Priacanthus arenatus*, *Pseudupeneus prayensis* and *Sparus caeruleostictus* in the lutjanid assemblage, and *Epinephelus aeneus* and *P. bellottii* in the sparid assemblage.

The pie charts in Fig. 2 display clearly significant shifts in relative importance of the various groups between the climatic periods and depth ranges. At 10-30 m, Sparidae accounted for 69% of the total density recorded for the three families (Sparidae, Sciaenidae and Lutjanidae) and *Balistes capriscus* in 1963-71. This was followed by Lutjanidae (26%), Sciaenidae (3%) and *B. capriscus* (2%). Between 1972 and 1982 when the population of *B. capriscus* increased in the Gulf of Guinea, the contribution of Sparidae in the 10-30 m depth range reduced substantially to 16%; Lutjanidae accounted for 7%, Sciaenidae 1% and *B. capriscus* 76%. In 1982-1990, the *Balistes capriscus* stock declined and accounted for only 4% of the total catch of the species and the other three groups. The percentage composition of Sparidae increased to 52%, Lutjanidae to 40%, and Sciaenidae to 2%.

In the 31-50 m depth range, the percentage composition of all three families reduced from the first climatic period to the second (81% to 20% for Sparidae; 11% to

4% for Lutjanidae and 2% to 1% for Sciaenidae). On the other hand *B. capriscus* increased in percentage composition from 6% to 75%. In 1982-1990, the situation changed to become similar to that in 1963-71, with the percentage contribution of *B. capriscus* substantially reduced and that of the other families increased.

In the 51-100 m depth range, the effect of *B. capriscus* was least shown. The percentage contribution of the species increased from 0% in 1963-71 to 15% in 1972-82. The contributions of the three families changed only slightly to accommodate this modest rise and subsequent decline of *Balistes capriscus*. In the third climatic period, the contribution of the species reduced to 3%. Accounting for 52-89% of catches of the four groups, Sparidae were the most important fishes at all depths for the three study periods, except between 1972 and 1982 where *Balistes capriscus* was dominant in the 10-30 m and 31-50 m depth ranges.

Possibility factors affecting fish community structure in the Gulf of Guinea

The significantly high correlation between depth, bottom temperature, bottom salinity and bottom dissolved oxygen and the first ordination axis and between sediment type and the second ordination axis, show the importance of these parameters in the determination of the structure and dynamics of the demersal species assemblages in the study area. It has been observed that changes could occur in the structure and composition of species assemblages as a result of differential reaction of species to seasonal environmental changes (Villegas & Garcia, 1983; Overholtz & Tyler, 1985). Thus the upwelling which appears to change the

properties of water masses could affect the structure and dynamics of demersal species assemblages and species diversity. Similarly, pollution which could arise from large-scale oil spillage, or chemical effluents from land-based sources, could have similar effect. Reduced river input into coastal waters due to the building of dams affect salinity regimes in coastal waters and could also affect the stability of the sciaenid assemblage.

Koranteng (1998) showed that during the first (1963-71) and third (1983-90) climatic periods the species assemblages were relatively easy to identify using ordination techniques, but in the second period (1972-1982), the assemblages were difficult to separate (especially the lutjanid and sparid assemblages). The author noted that the apparent disorganisation of the assemblages during the second climatic period could be attributed mainly to the sudden increase in abundance of *Balistes capriscus*.

During that period, the continental shelf waters were relatively cold and more saline; an environment preferred by the sparid community to which *B. capriscus* belongs (Gulland & Garcia, 1984). It is possible also that irresponsible fishing methods like the use of dynamite could also alter fish assemblage structure as it destroys habitats of the fish. For example, lutjanid species, the abundance of which appears to have declined in the study area, have been associated with corals (Longhurst, 1969) and is possible that these corals have been destroyed by dynamite or large industrial trawlers over the years.

It is important to note, however, that whereas environmental forcing could lead to temporal disruption of assemblages, habitat destruction could result in permanent change. However, it is often difficult, if not

impossible, to separate fluctuations due to natural factors from those caused by anthropogenic factors.

Diversity of continental shelf species assemblages in the Gulf of Guinea

An analysis of variance showed that H'' was significant ($P < 0.02$) between the climatic periods (Koranteng, 1998). Species richness and evenness also changed within the study period (Table 4). In 1963-71, the Margelef's Richness Index (R) was relatively higher in 10-30 m and 31-50 m, but these reduced significantly between 1972 and 1982. The Margelef's Richness Index increased again in both depth ranges between 1982 and 1990. The direction of change of R in the 51-100 m depth range was opposite that in the shallow areas, increasing between the first and second climatic periods but decreasing slightly after 1982. Pielou's Evenness Index J' also decreased between 1963-71 and 1972-82 but increased in 1983-90.

Because of differential reaction of individual species to environmental and ecosystem changes and to exploitation, species diversity is likely to change as a result of changes in the environment and/or exploitation. The recorded changes in the fish diversity index could be due to changes in the marine ecosystem relating to the environment, decline of the sardinella resources and proliferation of triggerfish. In the 10-30 m and 31-50 m depth ranges, the diversity index was lowest during 1972-82. In the 51-100 m depth range the diversity increased throughout the period under investigation. The increase of H'' in this depth range, especially in 1972-82, may have been due to migration of species predominantly from 31-50 m depth which

had quite similar sediment type. This is confirmed by the corresponding increase in species richness (R) at these depths during 1972-82. This situation further explains why assemblage discrimination was difficult during the 1972-82 period. The drastic reduction of evenness in 1972-82 at depths where triggerfish initially occupied is a further evidence of the effect of the proliferation of the species on the ecosystem.

It has been argued that diversity indices can be used to indicate the general 'health' of an ecosystem (Magurran, 1991); thus the decline in H' in the 1970s can be seen as an indication that the shelf ecosystem in the Gulf of Guinea was stressed at that time. The stress may be due to the declining temperature, increased salinity, changes in other near-shore forcing factors, increased exploitation, the 'invasion' of triggerfish, decrease of the *Sardinella aurita* stock in the western Gulf of Guinea or the combined effects of these. The invasion by triggerfish is perhaps a more likely cause considering the aggressive behaviour of the species. The magnitude of change in biomass recorded in the Gulf of Guinea, especially involving a comparatively hardy species like triggerfish, could induce structural changes in species assemblages.

Persistence in fish species assemblages

Long-term changes in fish assemblages have been attributed to a number of factors including over-exploitation of some species within the assemblage (Brown *et al.*, 1976; Overholtz & Tyler, 1985) and climatic variations (Sutcliffe & Muir 1977). Dominance or decline of some species in the assemblage may result from exploitation or differential response to changes in environmental forcing factors (Gulland & Garcia, 1984; Macpherson & Gordo, 1992).

Similarly, spatial changes in the assemblage can occur with the whole assemblage performing inshore-offshore temporal migrations in response to environmental forcing and pollution (e.g. Macdonald *et al.*, 1984; Gomes *et al.*, 1995).

The literature is rather inconsistent with respect to the persistence of fish species assemblages through time. One school of thought holds the view that marine fish communities are in constant motion and hold no memory that will enable them to return to a previous state (Tyler *et al.*, 1982; Beddington, 1984). On the other side of the argument, Longhurst & Pauly (1987) are of the view that "...one can assume that given sensitive management, or the relaxing of fishing pressure for some reason, tropical fish communities would tend to revert to the natural state...". McGlade (1989) also believes that "...natural systems do recover from large environmental perturbations". Greenstreet & Hall (1996) noted that differences in fish species assemblages over time are subtle and most apparent in the dominance structure. This was further underlined in an analysis of the effects of the industrial fisheries on the North Sea ecosystem (Robertson, McGlade & Leaver, 1996).

The results of this study appear to confirm the theory that the proliferation of *B. capricus* seriously disrupted the assemblage structure and behaviour of demersal fish species on the continental shelf of the Gulf of Guinea. Towards the end of the observation period when the species had almost disappeared from this ecosystem, the assemblages appeared similar to those before the dominance of *B. capricus*. However, it is uncertain whether the system was returning or had already returned to its original or 'natural' state.

Conclusion

The structure of species assemblages on the continental shelf in the Gulf of Guinea is determined primarily by depth and sediment type and the dynamics are influenced by physico-chemical parameters of the water masses. Structural changes in the assemblages and in species diversity in the area were probably due mainly to the sudden increase in abundance of *Balistes capriscus* and also environmental forcing. The relative importance of sciaenids, lutjanids and sparids on the continental shelf, as well as species richness, evenness and diversity changed in response to these alterations in the ecosystem. It appears that activities that lead to habitat alteration could bring about permanent changes in fish species assemblages, whereas temporal environmental changes could lead to short-term changes in the structure and dynamics of the assemblages.

Acknowledgement

The author is grateful to Dr Philippe Cury (IRD, Cape Town) and Prof. Jacqueline McGlade (UCL, London) and the reviewers for comments on the manuscript. The financial assistance provided by IRD (ex-ORSTOM) of France (through a research grant) and the European Union (through the INCO-DC Research and Technology for Development Programme) for this work is gratefully acknowledged.

References

- Bakun, A.** (1995). Global Climate Variations and Potential Impacts on the Gulf of Guinea Sardinella Fishery. In *Dynamics and Use of Sardinella Resources from Upwelling off Ghana and Ivory Coast* (F.X. Bard and K.A. Koranteng, ed.), pp. 60-74. Paris: ORSTOM Editions.
- Beddington, J. R.** (1984). The response of multispecies systems to perturbations. In *Exploitation of Marine Communities. Life Sciences Research 32* (R. M. May, ed.), pp. 209-225. New York: Springer-Verlag.
- Bianchi, G.** (1992). Demersal assemblages of tropical continental shelves. A study based on the data collected through the surveys of the R/V 'Dr. Fridtjof Nansen' (Thesis for Dr. Sci. degree), University of Bergen, Norway.
- Brown, B. E., Brennan, J. A., Grosslein, M. D., Heyerdahl, E. G. & Hennemuth, R. C.** (1976). The effect of fishing on the marine finfish biomass in the Northwest Atlantic from the Gulf of Maine to Cape Haterras. *International Commission for Northwest Atlantic Fisheries Research Bulletin 12*, 49-68.
- Caverivière, A.** (1991). L'explosion démographique du baliste (*Balistes carolinensis*) en Afrique de l'ouest et son évolution en relation avec les tendances climatiques. In *Variabilité, instabilité et changement dans les pêcheries ouest africaines* (P. Cury and C. Roy, ed.), pp. 354-367. Paris: ORSTOM Editions.
- Crossnier, A. & Berrit, G. R.** (1966). Fonds de pêche le long de la côte des Républiques du Dahomey et du Togo. ORSTOM (Miméo), pp. 1-94.
- Digby, P. G. N. & Kempton, R. A.** (1987). Multivariate analysis of ecological communities. London: Chapman and Hall, 206 pp.
- Entsua-Mensah, M.** (1998). Coastal lagoons and the continental shelf ecosystem of Ghana. *Paper presented at the Workshop on Sustainable Management of the Fishery resources in the Gulf of Guinea, 27-29 July 1998, Accra, Ghana.*
- Fager, E. W. & Longhurst, A. R.** (1968). Recurrent Group Analysis of Species Assemblages of Demersal Fish in the Gulf of Guinea. *Journal Fisheries Research Board of Canada 25* (7), 1405-1421.
- Gomes, M. C., Haedrich, R. L. & Villagarcia, M.G.** (1995). Spatial and temporal changes in

- the groundfish assemblages on the north-east Newfoundland/Labrador Shelf, North-west Atlantic, 1978-1991. *Fisheries Oceanography* 4(2), 85-101.
- Greenstreet, S. P. R. & Hall, S. J.** (1996). Fishing and the groundfish assemblage structure in the north-western North Sea: An analysis of long-term spatial trends. *J. Anim. Ecol.* 65, 577-598.
- Gulland, J. A. & Garcia, S.** (1984). Observed patterns in multispecies fisheries. In *Exploitation of marine communities. A report of the Dahlem Workshop on Exploitation of Marine Communities* (R.M. May, ed.), pp. 155-190. Berlin: Springer-Verlag.
- Hill, M. O.** (1979). *TWINSpan, A FORTRAN Program for Arranging Multivariate Data in an ordered Two-way Table by Classification of the Individuals and Attributes*. Ithaca, N.Y.: Cornell University.
- Joanny, T. & Menard, F.** (1998). Analysis of the spatial and temporal variability of demersal communities of the continental shelf of Côte d'Ivoire. *Paper presented at the Workshop on Sustainable Management of the Fishery resources in the Gulf of Guinea, 27-29 July 1998, Accra, Ghana*.
- Jongman, R. G. H., ter Braak, C. J. F. & van Tongeren, O. F. R.** (1995). *Data analysis in community and landscape ecology*. Cambridge: Cambridge University Press. 299 pp.
- Koranteng, K. A.** (1984). A trawling survey off Ghana, CECAF/TECH/84/63 CECAF Project, Dakar: (FAO), 72 pp.
- Koranteng, K. A.** (1998). The impacts of environmental forcing on the dynamics of demersal fishery resources of Ghana (PhD Thesis), University of Warwick. 377 pp.
- Koranteng, K. A. & Pezennec, O.** (1998). Variability and Trends in Environmental Time Series along the Ivorian and Ghanaian Coasts, In *Global versus Local Changes in Upwelling Systems* (M.-H. Durand, ed), pp. 167-177. Paris: ORSTOM Editions.
- Lamb, P. J.** (1982). Persistence on Saharan drought. *Nature* 299, 46-48.
- Longhurst, A. R.** (1964). The fish resources of the eastern Gulf of Guinea. *J. Cons.* 29, 302-334.
- Longhurst, A. R.** (1969). Species assemblages in tropical demersal fisheries. *Proceedings of the symposium on the oceanography and fisheries resources of the tropical Atlantic. Results of ICITA and GTS, Abidjan, Ivory Coast, 20-28 October 1966*, pp. 147-168. Paris: UNESCO Publications.
- Longhurst, A. R. & Pauly, D.** (1987). *Ecology of tropical oceans*. London: Academic Press, 407 pp.
- Macdonald, J. S., Dadswell, M. J., Appy, R. G., Melvin, G. D. & Methven, D. A.** (1984). Fishes, fish assemblages, and their seasonal movements in the lower Bay of Fundy and Passamaquoddy Bay, Canada. *Fishery Bulletin* 82 (1), 121 - 139.
- Macpherson, E. & Gordo, A.** (1992). Trends in the demersal fish community off Namibia from 1983 to 1990. *South African J. Marine Sci.* 12, 635-649.
- Magurran, A. E.** (1991). *Ecological diversity and its measurement*. London: Chapman and Hall.
- Mahé, G.** (1998). Freshwater yields to the Atlantic Ocean: local and regional variations from Senegal to Angola. In *Global versus Local Changes in Upwelling Systems* (M. H. Durand, ed.), pp. 127-138. Paris: ORSTOM Editions.
- Mann, K. H. & Lazier** (1996). *Dynamics of Marine Ecosystems, Biological-Physical Interactions in the Oceans*, 2nd edition. London: Blackwell Science: 394 pp.
- Martos, A. R., Yraola, I. S., Peralta, L. F. & Jimenez, J. F. G.** (1990). Informe de la Campana Guinea 90. Malaga, Spain: Instituto Español de Oceanografía.
- McGlade, J. M.** (1989). Fisheries Management: A review from science. In *Proceedings of the 3rd North Sea Seminar*, pp. 103-109.
- Overholtz, W. J. & Tyler, A. V.** (1985). Long-term Response of the Demersal Fish

- Assemblages of Georges Bank. *Fishery Bulletin* 83(4), 507-520.
- Pezenec, O.** (1995). Ecological importance of the Ivorian and Ghanaian minor upwelling season. In *Dynamics and Use of Sardinella Resources from Upwelling off Ghana and Ivory Coast*, (F.X. Bard and K.A. Koranteng, ed.), pp. 324-345. Paris: ORSTOM Editions.
- Pinkas, L., Oliphant, M. S. & Iverson, I. L. K.** (1971). Food habits of albacore, bluefin tuna and bonito in California waters. *Fishery Bulletin of the California Department of Fish and Game* 152, 1-105.
- Riley, J. P. & Chester, R.** (1971). Introduction to marine chemistry. London: Academic Press. 465 pp.
- Robertson, J., McGlade, J. & Leaver, I.** (1996). *Ecological effects of the North Sea fishing industry on the availability of human consumption species*. Univation, The Robert Gordon University, Aberdeen.
- Saenger, P., Santiane, Y., Baglo, M. Isebor, C., Armah, A. K. & Nganje, M.** (1998). *The Gulf of Guinea Project: Managing mangroves to protect biodiversity in West Africa*. <http://www.brooktrout.gso.uri.edu/ICmangaenger.html>.
- Shell** (1998). *Shell in Nigeria*. <http://www.shellnigeria.com/issues/environ>.
- Smilauer, P.** (1992). *CANODRAW*. Ithaca, N.Y.: Microcomputer Power.
- Stromme, T.** (1992). NAN-SIS, Software for fisheries survey data logging and analysis. User's manual. FAO Computerized Information Series (Fisheries). No. 4. Rome: FAO, 103 pp.
- Sutcliffe, W. H. Jr. & Muir, B. S.** (1977). Correlations of fish catch and environment factors in the Gulf of Maine. *J. Fisheries Res. Board of Canada* 34, 19-30.
- terBraak, C. J. F.** (1991). CANOCO-a FORTRAN program for canonical community ordination by [partial] [detrended] [canonical] correspondence analysis, principal component analysis and redundancy analysis (version 3.10). ITI-TNO, Wageningen: The Netherlands. (Manual and Update Notes).
- Tyler, A. V., Gabriel, W. L. & Overholtz, W. J.** (1982). Adaptive management based on structure of fish assemblages of northern continental shelves. In *Multispecies Approaches to Fisheries management Advice* (M. C. Mercer, ed.), pp. 149-156. Canadian Special Publication on Fisheries and Aquatic Sciences 59.
- Villegas, L. & Garcia, S.** (1983). *Demersal fish assemblages in Liberia, Ghana, Togo, Benin and Cameroon*. CECAF/ECAF Series 83/26 (En). Rome: FAO. 16 pp.
- Wilhm, J. L.** (1968). Use of biomass units in Shannon's formula. *Ecology* 49(1), 153-156.
- Williams, F.** (1968). *Report on the Guinean Trawling Survey*. Organisation of African Unity Scientific and Technical Research Commission (99).